



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Data on rhizosphere pH, phosphorus uptake and wheat growth responses upon TiO₂ nanoparticles application



Rafia Rafique^{a,e}, Zahra Zahra^a, Nasar Virk^b,
Muhammad Shahid^c, Eric Pinelli^d, Jean Kallerhoff^d,
Tae Jung Park^{e,*}, Muhammad Arshad^{a,*}

^a Institute of Environmental Sciences and Engineering, School of Civil and Environmental Engineering, National University of Sciences and Technology, Sector H-12, Islamabad 44000, Pakistan

^b Atta-ur-Rehman School of Biological Sciences, National University of Sciences and Technology Islamabad, 44000, Pakistan

^c Department of Environmental Science, COMSATS Institute of Information Technology, Vehari 61100, Pakistan

^d EcoLab, Université de Toulouse, CNRS, l'Agrobiopôle, Toulouse 31326, France

^e Department of Chemistry, Chung-Ang University, 84 Heukseok-ro, Dongjak-gu, Seoul 06974, Republic of Korea

ARTICLE INFO

Article history:

Received 14 December 2017

Received in revised form

30 January 2018

Accepted 1 February 2018

Available online 6 February 2018

Keywords:

Rhizosphere pH

TiO₂ NPs nanoparticles

Wheat

Phosphorus

Uptake

ABSTRACT

In this study, the data sets and analyses provided the information on the characterization of titanium dioxide nanoparticles (TiO₂ NPs), and their impacts on rhizosphere pH, and soil-bound phosphorus (P) availability to plants together with relevant parameters. For this purpose, wheat (*Triticum aestivum* L.) was cultivated in the TiO₂ NPs amended soil over a period of 60 days. After harvesting, the soil and plants were analyzed to examine the rhizosphere pH, P availability in rhizosphere soil, uptake in roots and shoots, biomass produced, chlorophyll content and translocation to different plant parts monitored by SEM and EDX techniques in response to different dosages of TiO₂ NPs. The strong relationship can be found among TiO₂ NPs application, P availability, and plant growth.

© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

DOI of original article: <https://doi.org/10.1016/j.agee.2017.12.010>

* Corresponding authors.

E-mail addresses: tjpark@cau.ac.kr (T.J. Park), marshad@iese.nust.edu.pk (M. Arshad).

<https://doi.org/10.1016/j.dib.2018.02.002>

2352-3409/© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	<i>Environmental and agricultural applications</i>
More specific sub- ject area	<i>Material synthesis, Effects of nanoparticles on soil-plant system, Nanobiotechnology</i>
Type of data	<i>Tables of TiO₂ effect on P concentration and plant biomass SEM and EDX images of TiO₂ NPs and their uptake by roots and leaves of plant Graphs of effect of TiO₂ NPs on rhizosphere soil, and P and chlorophyll content of leaves</i>
How data was acquired	<i>SEM, EDX, XRD, UV/Visible double beam spectrophotometer, chlorophyll meter (CCM 200-plus, Opti-Sciences, England, measurement area 0.7 cm²)</i>
Data format	<i>Raw, analyzed</i>
Experimental factors	<i>Wheat seeds were grown in sandy loam soil containing different concentrations of TiO₂ NPs. The experiments were held in a greenhouse for 60 days under ambient conditions.</i>
Experimental features	<i>Effects of soil application of TiO₂ NPs were measured on rhizosphere soil, roots, and shoots of plant</i>
Data source location	<i>Islamabad, Pakistan</i>
Data accessibility	<i>Data is available with this manuscript</i>

Value of the data

- The data provides information of TiO₂ NPs effects on wheat over a period of 60 days for better understanding of their long-term impacts on plant growth.
- The data can help to understand the relationship between TiO₂ NPs application and phytoavailability of P for farm and field level applications to ensure nutrient management.
- The data suggested the scientific community to extend the exposure time and comparison with other plant species instead of very short term bioassays.
- Future experiments can be compared with this data to predict the optimum concentrations of NPs for better plant development for different plant species.

1. Data

The datasets and analyses described the impacts of soil applied titanium dioxide nanoparticles (TiO₂ NPs) on wheat (*Triticum aestivum* L.) plants. Corresponding figures, graphs, and images are provided with this article.

2. Experimental design, materials, and methods

2.1. Synthesis and characterization of TiO₂ NPs

TiO₂ general purpose reagent was obtained from Sigma Aldrich Inc. (purity > 99%, St. Louis, MO, USA) and further processed, and calcined at 500 °C to synthesize pure anatase crystal structure of TiO₂ NPs as described in Zahra et al. [1]. Scanning electron microscope (SEM, Jeol, JSM 6490A, Tokyo, Japan), energy-dispersive X-ray spectroscopy (EDX, Jeol, JED 2300), and X-ray diffraction (XRD) analyses of as-prepared TiO₂ NPs were performed as shown in Fig 1.




Fig. 1. Characterization results of TiO₂ NPs. (a) SEM image, (b) EDX and, (c) XRD spectrum of TiO₂ NPs.




Fig. 2. Effect of TiO₂ NPs treatments on rhizosphere soil pH. Different alphabets correspond to statistically significant results at $p < 0.05$.

2.2. Soil application of TiO₂ NPs

TiO₂ NPs suspensions were prepared by mixing their various concentrations (0, 20, 40, 60, 80, and 100 mg L⁻¹) in deionized water and sonicated for 30 min. Four replicates of each treatment level, and the control group (without TiO₂ NPs) were maintained. Healthy seeds of wheat (Galaxy 2013) were obtained from the Ayub Agricultural Research Institute, Faisalabad, Pakistan. Three seeds were sown in each pot with as-prepared concentrations of TiO₂ NPs. The experiments were conducted in a greenhouse for 60




FIG. 3. Effect of TiO₂ NPs treatments on phytoavailability of P in rhizosphere soil. Different alphabets correspond to statistically significant results at $p < 0.05$.

Table 1

Effect of TiO₂ NPs treatments on plant shoot and root dry biomass of wheat.

TiO ₂ NPs Concentration (mg kg ⁻¹)	Shoot dry biomass (mg)	Root dry biomass (mg)	Total dry biomass (mg)
0	0.73 ± 0.09a	1.11 ± 0.16a	1.37 ± 0.06a
20	0.89 ± 0.05b	1.47 ± 0.12b	1.70 ± 0.05b
40	0.73 ± 0.42b	1.30 ± 0.74b	1.71 ± 0.17b
60	0.63 ± 0.59b	1.19 ± 1.14b	1.81 ± 0.39b
80	0.95 ± 0.05b	1.65 ± 0.13b	1.84 ± 0.04b
100	0.91 ± 0.05b	1.52 ± 0.1b	1.71 ± 0.07b

The values are the means of four replicates ± Standard Deviation (SD). The means followed by similar letter (a) in the same column are not significantly different whereas (b) represents statistically significant difference at $p < 0.05$.

days following randomized block design where the position of the pots was altered to avoid environmental bias effects.

2.3. Analysis of soil and plants

After 60 days of TiO₂ NPs exposure, the plants were uprooted and shaken carefully to remove soil at harvesting. The loosely bound soil adhered to the roots was collected with gentle washing in distilled water (100 mL) to investigate the rhizosphere pH (Fig. 2) and P (Fig. 3) using Olsen's method [2]. The roots and shoots were cut and dried in hot air oven for 48 h at 70 °C. After that, the dry biomass was recorded (Table 1) and stored for P analysis. For plant P content analysis, 100 mg of ground plant samples were added to acid mixture containing 5 mL of HNO₃/HClO₄ (2:1). This was digested on a hot plate followed by filtration through Whatman filter paper no. 42 to get clear aliquots for P content analysis (Table 2) using the vanado-molybdo-phosphoric acid colorimetric method [3].

2.4. Estimation of leaf chlorophyll content

A hand-held chlorophyll meter was used to measure the chlorophyll content index (CCI). The CCI readings were taken after the 30th day of NPs exposure for 16 alternate days until harvest. The everyday measurements are the mean of 32–48 readings for each treatment (Fig. 4). Following

Table 2Effect of TiO₂ NPs treatments on plant shoot and root P concentration of wheat.

TiO ₂ NPs Concentration (mg kg ⁻¹)	Shoot P concentration (mg)	Root P concentration (mg)	Total P concentration (mg)
0	1.52 ± 0.17a	1.48 ± 0.13a	3.00 ± 0.03a
20	1.65 ± 0.08b	1.84 ± 0.12b	3.49 ± 0.14b
40	1.77 ± 0.06b	2.08 ± 0.16b	3.85 ± 0.22b
60	0.85 ± 0.08b	2.52 ± 0.19b	4.37 ± 0.47b
80	1.73 ± 0.07b	1.99 ± 0.18b	3.72 ± 0.18b
100	1.68 ± 0.06b	1.90 ± 0.16b	3.58 ± 0.15b

Data is the mean of four replicates ± Standard Deviation (SD). Means followed by different letters in the same column indicate significantly significant results at $p < 0.05$.




Fig. 4. Effect of TiO₂ NPs treatments on foliar chlorophyll content of wheat on daily basis.

calibration Eq. (1) was used to process the raw data and convert the CCI index values to chlorophyll content expressed as m cm^{-2} [4].

$$y = -2.20e^{-03} + 3.09e^{-03}x - 5.63e^{-05}x^2 \quad (1)$$

where y = Total chlorophyll content

x = Chlorophyllmeter value

2.5. Microscopic analysis of plant

To investigate the uptake of TiO₂ NPs, plant samples were observed under SEM equipped with EDX to demonstrate the elemental composition of the control (0 mg kg⁻¹ TiO₂ NPs) and treated (60 mg kg⁻¹ TiO₂ NPs) samples of roots (Fig. 5), and shoots (Fig. 6).

2.6. Statistical analysis

The statistical significance analysis was done using Student's *T*-Test available in the Microsoft Excel analysis tool box. One-way ANOVA test was performed to identify statistically significant differences between the treatments. Statistix 8.1 was used to identify the least significant differences (LSD) at $p < 0.05$. All the data presented here supports the findings and discussion in Rafique et al. [5].




Fig. 5. SEM and EDX analysis of wheat roots at (a) 0 mg kg⁻¹, and (b) 60 mg kg⁻¹. The EDX spectrum was measured at 20 keV.




Fig. 6. SEM and EDX analysis of wheat leaves at (a) 0 mg kg⁻¹, and (b) 60 mg kg⁻¹ of TiO₂ NPs treatment. The EDX spectrum was measured at 20 keV.

Acknowledgements

The authors would like to thank Higher Education Commission (HEC) of Pakistan for financial support under Project No. 20-3060/NRPU/R&D/HEC/13. This work was also supported by Basic Science Research Program, National Research Foundation of Korea (NRF), Ministry of Science, ICT and Future Planning (NRF-2017R1A2B4009581).

Transparency document. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2018.02.002>.

References

- [1] Z. Zahra, N. Waseem, R. Zahra, H. Lee, M.A. Badshah, A. Mehmood, H.K. Choi, M. Arshad, Growth and metabolic responses of rice (*Oryza sativa* L.) cultivated in phosphorus-deficient soil amended with TiO₂ nanoparticles, *J Agric. Food Chem.* 65 (2017) 5598–5606.
- [2] S.R. Olsen, C.V. Cole, F.S. Watanabe, L.A. Dean, Estimation of Available Phosphorus in Soil by Extraction with Sodium Bicarbonate, 939, United States Department of Agriculture, Washington, D.C. (1954) 18–19.
- [3] J. Ryan, G. Estefan, A. Rashid, *Soil and Plant Analysis: Laboratory Manual*, 2nd ed., International Center for Agricultural Research in the Dry Areas (ICARDA), Aleppo, Syria, 2001.
- [4] A.D. Richardson, S.P. Duigan, G.P. Berlyn, An evaluation of noninvasive methods to estimate foliar chlorophyll content, *New Phytol.* 153 (2002) 185–194.
- [5] R. Rafique, Z. Zahra, N. Virk, M. Shahid, E. Pinelli, J. Kallerhoff, T.J. Park, M. Arshad, Dose-dependent physiological responses of *Triticum aestivum* L. to soil applied TiO₂ nanoparticles: alterations in chlorophyll content, H₂O₂ production, and genotoxicity, *Agric. Ecosyst. Environ.* 255 (2018) 95–101.