

Hypothermia Inhibits Endothelium-Independent Vascular Contractility via Rho-kinase Inhibition

Yoon Hee Chung^{1,†}, Keon Woong Oh^{2,†}, Sung Tae Kim³, Eon Sub Park², Hyun Dong Je⁴, Hyuk-Jun Yoon⁴,
Uy Dong Sohn⁵, Ji Hoon Jeong^{3,*} and Hyen-Oh La^{6,*}

Departments of ¹Anatomy, ²Pathology, ³Pharmacology, College of Medicine, Chung-Ang University, Seoul 06974,

⁴Department of Pharmacology, College of Pharmacy, Catholic University of Daegu, Gyeongsan 38430,

⁵Department of Pharmacology, College of Pharmacy, Chung Ang University, Seoul 06974,

⁶Department of Pharmacology, College of Pharmacy, The Catholic University of Korea, Bucheon 14662, Republic of Korea

Abstract

The present study was undertaken to investigate the influence of hypothermia on endothelium-independent vascular smooth muscle contractility and to determine the mechanism underlying the relaxation. Denuded aortic rings from male rats were used and isometric contractions were recorded and combined with molecular experiments. Hypothermia significantly inhibited fluoride-, thromboxane A₂-, phenylephrine-, and phorbol ester-induced vascular contractions regardless of endothelial nitric oxide synthesis, suggesting that another pathway had a direct effect on vascular smooth muscle. Hypothermia significantly inhibited the fluoride-induced increase in pMYPT1 level and phorbol ester-induced increase in pERK1/2 level, suggesting inhibition of Rho-kinase and MEK activity and subsequent phosphorylation of MYPT1 and ERK1/2. These results suggest that the relaxing effect of moderate hypothermia on agonist-induced vascular contraction regardless of endothelial function involves inhibition of Rho-kinase and MEK activities.

Key Words: ERK1/2, fluoride, hypothermia, MYPT1, phorbol ester, Rho-kinase

INTRODUCTION

Hypothermia which is defined as a body core temperature below 35.0°C, is characterized by shivering, mental confusion, muscle miscoordination, unstable physiological systems, and decreases in heart rate (Tankersley *et al.*, 2003), respiratory rate, and blood pressure. Although hypothermia can be beneficial in cardiac arrest (Oh *et al.*, 2015) and traumatic brain injury (Jensen *et al.*, 2016), it is a risk factor for arrhythmia, bleeding, thrombosis, sepsis, and intraventricular hemorrhage. Hypothermia can induce nitric oxide-dependent vasodilatation, which can be prevented by atropine in isolated canine coronary, femoral, and renal arteries (Evora *et al.*, 2007), suggesting that endothelium-dependent relaxation involves the release of nitric oxide and other vasoactive mediators (Zou *et al.*, 2015). Here, we investigated the mechanisms underlying the endothelium-independent relaxing effect of hypothermia on vascular smooth muscle contractility. Intact or denuded

aortic rings from male Sprague-Dawley rats were used, and isometric contractions were recorded using a computerized data acquisition system. These data were combined with molecular experiments.

Hypertension may increase morbidity and mortality due to other diseases in the presence of other risk factors such as genetics, diet, and lifestyle. Strategies for the prevention and management of hypertension and associated adverse consequences are based on lifestyle modification as well as pharmacological interventions (Appel *et al.*, 2006). Contraction of vascular smooth muscle is regulated by both Ca²⁺-dependent and Ca²⁺-sensitization mechanisms. The extent of MLC₂₀ phosphorylation or force of contraction induced by agonist stimulation is usually higher than that caused by an increase in the cytosolic Ca²⁺ concentration, which is referred to as Ca²⁺ sensitization (Akata, 2007; Kim *et al.*, 2011). The mechanism responsible for Ca²⁺ sensitization involves inhibition of myosin light chain phosphatase, leading to increased phos-

Open Access <https://doi.org/10.4062/biomolther.2016.233>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received Oct 20, 2016 Revised Dec 11, 2016 Accepted Dec 27, 2016
Published Online Feb 17, 2017

*Corresponding Authors

E-mail: jhjeong3@cau.ac.kr (Jeong JH), hola@catholic.ac.kr (La HO)
Tel: +82-2-820-5688 (Jeong JH), +82-2-2164-4049 (La HO)
Fax: +82-2-826-8752 (Jeong JH), +82-2-2164-4059 (La HO)

[†]The first two authors contributed equally to this work.

phorylation of the 20-kDa myosin light chain and subsequent enhanced contraction. Inhibition of MLCP in vascular smooth muscle is mediated by phosphorylation of the myosin phosphatase target subunit of MLCP via Rho-kinase, which leads to sustained phosphorylation of MLC₂₀. Subsequent studies have suggested that the inhibition of myosin phosphatase by Rho-kinase (Uehata *et al.*, 1997; Sakurada *et al.*, 2003; Akata, 2007) or thin filament regulation including the activation of protein kinase C (PKC), mitogen-activated protein kinase kinases (MEK), and extracellular signal regulated kinase (ERK) 1/2, as well as phosphorylation of the actin binding protein caldesmon (Wier and Morgan, 2003) may precipitate Ca²⁺ sensitization.

Activation of ERK1/2, in addition to regulating vascular contractility, is also connected to pathologic hypertrophy, hyperplasia, hypertension, and atherosclerosis (Ruppert *et al.*, 2013; Brietz *et al.*, 2016). ERK1/2 is activated by threonine and tyrosine phosphorylation by the Raf-activated kinase MEK. Fluoride, phorbol esters, and thromboxane A₂ mimetics have been shown to induce contractions of smooth muscles, which may be due primarily to enhanced Ca²⁺ sensitivity or partially to increased Ca²⁺ concentration. ERK1/2 activation was induced by thromboxane A₂ mimetic (Gallet *et al.*, 2003) or phorbol ester. Activation of ERK1/2 triggers ERK1/2-dependent cytoskeletal remodeling and formation of podosomes (Gu *et al.*, 2007). However, it is possible that the contractions induced by fluoride or thromboxane A₂ mimetics involve the RhoA/Rho-kinase pathway (Jeon *et al.*, 2006). However, it is not known whether this pathway is inhibited during hypothermia-induced vascular smooth muscle relaxation in aortic rings contracted with a Rho-kinase activator (fluoride), a MEK activator (phorbol ester), or with dual activators of both Rho-kinase and MEK (thromboxane A₂ or phenylephrine). Our aim in the present study was to investigate the possible roles of Rho-kinase or MEK inhibition on Ca²⁺ desensitization during hypothermia-induced relaxation of isolated, denuded rat aortas using Rho-kinase (fluoride) or MEK (phorbol ester) activators.

MATERIALS AND METHODS

Tissue preparation

Male Sprague-Dawley rats weighing 200-250 g were anesthetized with sodium pentobarbital (50 mg/kg i.p.) and then subjected to cervical dislocation, in accordance with procedures approved by the Institutional Animal Care and Use Committee of Chung-Ang University (Seoul, Korea) and Catholic University of Daegu (Daegu, Korea; IACUC-2014-041). Thoracic aortas were quickly removed and immersed in oxygenated (95% O₂/5% CO₂) physiological saline solution composed of (mM): 115.0 NaCl, 4.7 KCl, 2.5 CaCl₂, 1.2 MgCl₂, 25.0 NaHCO₃, 1.2 KH₂PO₄, and 10.0 dextrose (pH 7.4). They were then freed of all adherent connective tissue, and aortic endothelia were removed by gentle abrasion using a cell scraper if necessary.

Contraction measurements

Two stainless-steel triangles were inserted through each vessel ring, and each aortic ring was suspended in a water-jacketed organ bath (10 ml) that was adjusted from 37°C to 28°C (moderate hypothermia) using a thermostat and water baths and was aerated with a mixture of 95% O₂ and 5% CO₂. One triangle was anchored to a stationary support, and the

other was connected to an isometric force transducer (Grass FT03C, Quincy, MA, USA). Rings were stretched passively by applying an optimal resting tension of 2.0 g, which was maintained throughout the experiment. Each ring was equilibrated in the organ bath solution for 60 min before contractile responses to 50 mM KCl or 1 μM phenylephrine were measured. Isometric contractions where the muscle length does not change during contraction (Ito *et al.*, 2015), were recorded using one triangle anchored to a stationary support and the other connected to an isometric force transducer, an amplifier, and a computerized data acquisition system (PowerLab/8SP, AD Instruments, Castle Hill, NSW, Australia).

The indirect effect of hypothermia was determined by application of a temperature of 28°C using a thermoregulator and a water bath, 10 min before the application of fluoride (6 mM), thromboxane A₂ mimetic (0.1 μM), phenylephrine (1 μM), or phorbol ester (1 μM) in normal Krebs' solution.

Western blot analysis

Muscle strips were quick-frozen by immersion in a dry ice/acetone slurry containing 10% trichloroacetic acid (TCA) and 10 mM dithiothreitol (DTT). Muscles were stored at -80°C until use. Tissues were brought up to room temperature in a dry ice/acetone/TCA/DTT mixture and then homogenized in a buffer containing 20 mM MOPS, 4% SDS, 10% glycerol, 10 mM DTT, 20 mM β-glycerophosphate, 5.5 μM leupeptin, 5.5 μM pepstatin, 20 kIU aprotinin, 2 mM Na₃VO₄, 1 mM NaF, 100 μM ZnCl₂, 20 μM 4-(2-aminoethyl) benzenesulphonyl fluoride (AEBSF), and 5 mM EGTA. Protein-matched samples (modified Lowry protein assay, DC Protein Assay Kit, Bio-Rad, Hercules, CA, USA) were subjected to sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) (Protogel, National Diagnostics, Atlanta, Georgia, USA), transferred to polyvinylidene fluoride (PVDF) membranes, and subjected to immunostaining and densitometry using primary and secondary antibodies. Lane loading variations were corrected by normalization versus β-actin. Sets of samples produced during individual experiments were run in the same gel, and densitometry was performed on the same image.

Chemicals and antibodies

Sodium fluoride, KCl, phenylephrine, acetylcholine, U46619, Y-27632, and phorbol 12,13-dibutyrate were purchased from Sigma (St. Louis, MO, USA). DTT, TCA, and acetone were obtained from Fisher Scientific (Hampton, NH, USA). Enhanced chemiluminescence (ECL) kits were from Pierce (Rockford, IL, USA). Antibodies against phospho-myosin phosphatase targeting subunit protein 1 (phospho-MYPT1) at Thr855 (1:5,000), MYPT1, ERK, or phosphoERK at Thr202/Tyr204 (Cell Signaling Technology, Danvers, MA, USA or Upstate Biotechnology, Lake Placid, NY, USA) were used to determine levels of RhoA/Rho-kinase activity (Wooldridge *et al.*, 2004; Wilson *et al.*, 2005) or MEK activity. Anti-mouse IgM (goat) and anti-rabbit IgG (goat) conjugated with horseradish peroxidase were used as secondary antibodies (1:2,000 dilutions for both, Upstate). A specific MLC₂₀ antibody (1:1500, Sigma) and anti-mouse IgG (goat) conjugated to horseradish peroxidase (1:2000, Upstate) were used to determine the level of LC₂₀ phosphorylation.

Statistics

Data are expressed as mean ± standard error of the mean




Fig. 1. Effect of hypothermia or normothermia on fluoride-induced vascular contraction in denuded (A) or intact (B) muscles. Each ring was equilibrated in organ bath solution for 30-60 min before relaxation responses to hypothermia were measured. Data are expressed as the mean of 3-5 experiments with the vertical line representing SEM. * $p < 0.05$, ** $p < 0.01$, presence versus absence of hypothermia.




Fig. 2. Effect of hypothermia on fluoride-induced increases in phospho-MYPT1 level. Phospho-MYPT1 protein level decreased in rapidly-frozen hypothermia-treated rat aorta in the absence of endothelium compared to normothermia-treated rat aortas contracted with fluoride. The upper panel shows a typical blot, and the lower panel shows average densitometry results for relative levels of phospho-MYPT1. Data are expressed as the mean of 3-5 experiments with the vertical line representing SEM. * $p < 0.05$, ## $p < 0.01$, versus control or normal group respectively. Hypothermia: 28°C; Fluoride: 6 mM sodium fluoride.

(SEM). Student's unpaired *t* test or ANOVA was used to determine the statistical significance of differences in means between two groups using SPSS 12.0 (SPSS Inc., Chicago, IL, USA). *p*-values < 0.05 were regarded as statistically significant.

RESULTS

Effect of hypothermia on contractions of endothelium-denuded aortas induced by the full RhoA/Rho-kinase activator fluoride

Endothelium was removed by gentle abrasion with a cell




Fig. 3. Effect of hypothermia or normothermia on thromboxane A₂-induced vascular contraction in denuded muscles. Each ring was equilibrated in the organ bath solution for 30-60 min before relaxation responses to hypothermia were measured. Data are expressed as the mean of 3-5 experiments with the vertical line representing SEM. * $p < 0.05$, presence versus absence of hypothermia.

scraper to identify the direct effect of hypothermia on vascular smooth muscle. The absence of endothelium was confirmed by a lack of relaxation after treating contracted ring segments with acetylcholine (1 μM). Moderate hypothermia (28°C) had no significant effect on basal tension (data not shown), but significantly inhibited contraction induced by the Rho-kinase activator fluoride, regardless of the absence of endothelial nitric oxide synthesis (Fig. 1A) or intact (Fig. 1B) muscles. This suggests that the relaxation mechanism of hypothermia might involve the inhibition of Rho-kinase activity in addition to endothelial nitric oxide synthesis and the subsequent activation of guanylyl cyclase.

Effect of hypothermia on the level of MYPT1 phosphorylation at Thr-855

To confirm the role of hypothermia on thick filament regulation of smooth muscle contractility, we measured levels of myosin phosphatase targeting subunit 1 (MYPT1) and phospho-MYPT1 in muscles quick-frozen after 60 min exposure to hypothermia for equilibration. Each relaxing ring was contract-

ed with 6 mM fluoride. This work was conducted using quick frozen moderate hypothermia (28°C)-treated rat aortas in the absence of endothelium, and levels were compared to those




Fig. 4. Effect of hypothermia or normothermia on phenylephrine-induced vascular contraction in denuded muscles. Each ring was equilibrated in the organ bath solution for 30-60 min before relaxation responses to hypothermia were measured. Data are expressed as the mean of 3-5 experiments with the vertical line representing SEM. * $p < 0.05$, presence versus absence of hypothermia.

of normothermia-treated rat aortas (Fig. 2). Interestingly, a significant decrease in fluoride-induced MYPT1 phosphorylation at Thr855 in response to hypothermia treatment was observed (Fig. 2). Thus, thick or myosin filament regulation, including myosin phosphatase activation via RhoA/Rho-kinase inactivation, might be involved in the reduced contractility of hypothermia-treated rat aortas.

Effect of hypothermia on contraction of denuded aortas induced by the dual Rho kinase and MEK activators thromboxane A₂ and phenylephrine

Moderate hypothermia (28°C) inhibited thromboxane A₂ mimetic U46619- (Fig. 3) or phenylephrine-induced contraction (Fig. 4) of denuded muscles, suggesting that dual activators (thromboxane A₂ mimetic and phenylephrine) act similarly to full activators targeting Rho-kinase.

Effect of hypothermia on contractions of denuded aortas induced by the MEK activator phorbol 12, 13-dibutyrate

Phorbol esters are primarily MEK activators and partial Rho-kinase activators (Goyal *et al.*, 2009; Je and Sohn, 2009). Interestingly, phorbol 12,13-dibutyrate (PDBu)-induced contraction was inhibited by moderate hypothermia (28°C), regardless of endothelial nitric oxide synthesis in denuded muscles (Fig. 5), which suggested that thin or actin filament regulation including MEK/ERK was inhibited.




Fig. 5. An original tracing of phorbol ester-induced vascular contraction in a normothermia (A) or hypothermia (B)-treated denuded rat aorta.




Fig. 6. Effect of hypothermia on phorbol ester-induced increases in phospho-ERK1/2 levels. Phospho-ERK1/2 protein levels were decreased in rapidly-frozen hypothermia-treated rat aortas in the absence of endothelium compared to normothermia-treated rat aortas contracted with a phorbol ester. The upper panel shows a typical blot, and the lower panel shows average densitometry results for relative levels of phospho-ERK1/2. Data are expressed as the mean of 3-5 experiments with the vertical line representing SEM. * $p < 0.05$, ## $p < 0.01$, versus control or normal group respectively. Hypothermia: 28°C; Phorbol ester: 1 μ M phorbol 12,13-dibutyrate.

Effect of hypothermia on levels of ERK1/2 phosphorylated at Thr-202/Tyr-204

To confirm the role of hypothermia on thin filament regulation of smooth muscle contractility, we measured levels of ERK1/2 and phospho-ERK1/2 in muscles quick-frozen after 60 minutes of exposure to hypothermia for equilibration. Each relaxing ring was contracted with 1 μ M phorbol 12,13-dibutyrate. As compared with normothermia-treated rat aortas, a significant decrease in ERK 1/2 phosphorylation at Thr202/Tyr204 was observed in moderate hypothermia (28°C)-treated rat aortas in the absence of endothelium (Fig. 6); full vasorelaxation (Fig. 5) and thin filament regulation were observed. These findings indicate that thin or actin filament regulation, including ERK1/2 phosphorylation via MEK activation, plays an important role in hypothermia-induced relaxation.

Effect of hypothermia on the level of LC₂₀ phosphorylation

To investigate if hypothermia affected fluoride-induced MLC₂₀ phosphorylation, aortic rings were treated with hypothermia 10 min before the addition of fluoride. Each relaxing ring was contracted with 6 mM sodium fluoride. This work was performed using quick-frozen hypothermia (28°C)-treated rat aortas in the absence of endothelium, and levels were compared to those of normothermia-treated rat aortas (Fig. 7). A significant decrease in fluoride-induced LC₂₀ phosphorylation was found in response to hypothermia treatment (Fig. 7) or Y27632 (Rho kinase inhibitor) treatment. Thus, the reduced contractility of hypothermia-treated rat aorta may involve thick or myosin filament regulation via mechanisms including calcium immobilization and MLCK inactivation.




Fig. 7. Effect of hypothermia on the fluoride-induced increase in phospho-MLC₂₀ level. Rat aortas were treated with 1 μ M Y-27632 and exposed to hypothermic conditions. Phospho-MLC₂₀ level expressed as a percentage of total MLC₂₀ was significantly lower in rapidly-frozen hypothermia-treated rat aortas in the absence of endothelium than in normothermia-treated rat aortas contracted with fluoride (6 mM). Data are expressed as the mean of 3-5 experiments with the vertical line representing SEM. * $p < 0.05$, ## $p < 0.01$, versus control or normal group respectively. Fluoride: 6 mM sodium fluoride; Hypothermia: 28°C; Y: 1 μ M Y27632.

DISCUSSION

The present study demonstrates that moderate hypothermia can modulate vascular contractility in an endothelium-independent manner. Interestingly, the mechanism involved seems not only to be endothelium-dependent, but also to involve the inhibition of MEK and Rho-kinase activity. Hypothermia resulting from extreme exposure to cold, alcohol intoxication, low blood sugar, anorexia, and advanced age has been previously recognized for its sympathetic nervous system excitation and hepatic dysfunction. Therefore, we investigated whether the inhibition of RhoA/Rho-kinase or MEK activity contributed to hypothermia-induced vascular relaxation in denuded rat aortas contracted by a Rho-kinase activator (fluoride), a MEK activator (phorbol 12,13-dibutyrate), or a dual activator (thromboxane A₂ and phenylephrine).

The mechanism by which phorbol esters or thromboxane A₂ mimetics activate MEK/ERK has been established (Gu *et al.*, 2007). In contrast, previous studies that examined the mechanisms underlying arterial contractions induced by fluoride, thromboxane A₂, or phenylephrine have reported conflicting findings with regard to contractions triggered by Rho-kinase activation (Wilson *et al.*, 2005; Tsai and Jiang, 2006). These findings are consistent with the notion that hypothermia can decrease fluoride-, thromboxane A₂ mimetic-, and phorbol ester-induced contraction by inhibiting MEK or Rho-kinase activity.

The mechanisms by which MEK or Rho-kinase activation cause vascular contraction are being studied intensively, and several possibilities exist. Rho-kinase phosphorylates myosin light chain phosphatase, which decreases phosphatase activity and causes a buildup of phosphorylated myosin light chains (Pfitzer, 2001; Akata, 2007). Rho-kinase has also been dem-

onstrated to phosphorylate myosin light chains directly and independently of myosin light chain kinase and phosphatase activity (Amano *et al.*, 1996). Recently, Rho-kinase was found to be involved in vascular contractions evoked by fluoride, phorbol esters or thromboxane A₂ (Wilson *et al.*, 2005; Jeon *et al.*, 2006; Tsai and Jiang, 2006).

The present study demonstrated that hypothermia ameliorated contraction induced by vasoconstrictors (fluoride or phorbol ester) in an endothelium-independent manner (Fig. 1, 3, 4, 5), and that the mechanism involved the MEK/ERK and RhoA/Rho-kinase pathways. Previously, most vasodilation was believed to be caused by endothelial nitric oxide synthesis and the subsequent activation of guanylyl cyclase (Taubert *et al.*, 2002; Ajay *et al.*, 2003; Zou *et al.*, 2015). In the present study, moderate hypothermia significantly inhibited fluoride- or phorbol ester-induced contraction, regardless of endothelial function (Fig. 1, 5). Furthermore, hypothermia significantly decreased the phosphorylation of MYPT1 at Thr855 induced by fluoride (Fig. 2) with full relaxation (Fig. 1) and ERK 1/2 phosphorylation at Thr202/Tyr204 induced by a phorbol ester (Fig. 6), suggesting that inhibition of MEK or Rho-kinase activity is a major mechanism underlying the effects of hypothermia on smooth muscle contractility. MLC₂₀ is known to be phosphorylated by both MLCK and Rho-kinase (Somlyo and Somlyo, 2003). Activation of Rho-kinase by phenylephrine or fluoride inhibits the activity of myosin light chain phosphatase through phosphorylation of MYPT1, leading to an increase in MLC₂₀ phosphorylation as well as contractions (Sakurada *et al.*, 2003; Wilson *et al.*, 2005; Akata, 2007). Both phenylephrine and fluoride induced phosphorylation of MLC₂₀, and this was inhibited by Y27632 (Ferland *et al.*, 2016) and hypothermia (Fig. 7).

In summary, moderate hypothermia significantly attenuated contractions induced by the RhoA/Rho-kinase activator fluoride independent of endothelial function. Furthermore, moderate hypothermia significantly inhibited phorbol ester-induced contraction due to MEK activation. Thus, the mechanism underlying the relaxation induced by moderate hypothermia of fluoride- or phorbol ester-induced contractions involves inhibition of MEK and Rho-kinase activity. Inhibition of Rho-kinase activity and subsequent MYPT1 phosphorylation induced by hypothermia during fluoride-induced contraction suggests that Rho-kinase inactivation is required for relaxation. In conclusion, in addition to endothelial nitric oxide synthesis in intact muscle, both MEK and Rho-kinase inhibition contribute synergistically to hypothermia-induced vasorelaxation of denuded smooth muscle.

ACKNOWLEDGMENTS

This research was supported by a Chung-Ang University Research Scholarship Grant in 2015.

REFERENCES

Ajay, M., Gilani, A. U. and Mustafa, M. R. (2003) Effects of flavonoids on vascular smooth muscle of the isolated rat thoracic aorta. *Life Sci.* **74**, 603-612.
 Akata, T. (2007) Cellular and molecular mechanisms regulating vascular tone. Part 2: regulatory mechanisms modulating Ca²⁺ mobiliza-

tion and/or myofilament Ca²⁺ sensitivity in vascular smooth muscle cells. *J. Anesth.* **21**, 232-242.
 Amano, M., Ito, M., Kimura, K., Fukata, Y., Chihara, K., Nakano, T., Matsuura, Y. and Kaibuchi, K. (1996) Phosphorylation and activation of myosin by Rho-associated kinase (Rho-kinase). *J. Biol. Chem.* **271**, 20246-20249.
 Appel, L. J., Brands, M. W., Daniels, S. R., Karanja, N., Elmer, P. J. and Sacks, F. M. (2006) Dietary approaches to prevent and treat hypertension: a scientific statement from the American Heart Association. *Hypertension* **47**, 296-308.
 Brietz, A., Schuch, K. V., Wangorsch, G., Lorenz, K. and Dandekar, T. (2016) Analyzing ERK 1/2 signalling and targets. *Mol. Biosyst.* **12**, 2436-2446.
 Evora, P. R., Cable, D. G., Chua, Y. L., Rodrigues, A. J., Pearson, P. J. and Schaff, H. V. (2007) Nitric oxide and prostacyclin-dependent pathways involvement on *in vitro* induced hypothermia. *Cryobiology* **54**, 106-113.
 Ferland, D. J., Darios, E. S., Neubig, R. R., Sjögren, B., Truong, N., Torres, R., Dexheimer, T. S., Thompson, J. M. and Watts, S. W. (2016) Chemerin-induced arterial contraction is Gi- and calcium-dependent. *Vascul. Pharmacol.* **88**, 30-41.
 Gallet, C., Blaie, S., Lévy-Toledano, S. and Habib, A. (2003) Thromboxane-induced ERK phosphorylation in human aortic smooth muscle cells. *Adv. Exp. Med. Biol.* **525**, 71-73.
 Goyal, R., Mittal, A., Chu, N., Shi, L., Zhang, L. and Longo L. D. (2009) Maturation and the role of PKC-mediated contractility in ovine cerebral arteries. *Am. J. Physiol. Heart Circ. Physiol.* **297**, H2242-H2252.
 Gu, Z., Kordowska, J., Williams, G. L., Wang, C. L. and Hai, C. M. (2007) Erk1/2 MAPK and caldesmon differentially regulate podosome dynamics in A7r5 vascular smooth muscle cells. *Exp. Cell Res.* **313**, 849-866.
 Ito, K., Matsuzaki, M., Sasahara, T., Shin, M. and Yayama, K. (2015) Orthovanadate-induced vasoconstriction of rat mesenteric arteries is mediated by rho kinase dependent inhibition of myosin light chain phosphatase. *Biol. Pharm. Bull.* **38**, 1809-1816.
 Je, H. D. and Sohn, U. D. (2009) Inhibitory effect of genistein on agonist-induced modulation of vascular contractility. *Mol. Cells* **27**, 191-198.
 Jensen, K. O., Held, L., Kraus, A., Hildebrand, F., Mommsen, P., Mica, L., Wanner, G. A., Steiger, P., Moos, R. M., Simmen, H. P. and Sprengel, K. (2016) The impact of mild induced hypothermia on the rate of transfusion and the mortality in severely injured patients: a retrospective multi-centre study. *Eur. J. Med. Res.* **21**, 37.
 Jeon, S. B., Jin, F., Kim, J. I., Kim, S. H., Suk, K., Chae, S. C., Jun, J. E., Park, W. H. and Kim, I. K. (2006) A role for Rho kinase in vascular contraction evoked by sodium fluoride. *Biochem. Biophys. Res. Commun.* **343**, 27-33.
 Kim, J. G., Sung, H. J., Ok, S. H., Kwon, S. C., Cheon, K. S., Kim, H. J., Chang, K. C., Shin, I. W., Lee, H. K., Chung, Y. K. and Sohn, J. T. (2011) Calcium sensitization involved in dexmedetomidine-induced contraction of isolated rat aorta. *Can. J. Physiol. Pharmacol.* **89**, 681-689.
 Oh, T. G., Cha, W. C., Jo, I. J., Kang, M. J. and Lee, D. W. (2015) A survey-based study on the protocols for therapeutic hypothermia in cardiac arrest patients in Korea: focusing on the differences between level 1 and 2 centers. *Clin. Exp. Emerg. Med.* **2**, 210-216.
 Pfitzer, G. (2001) Invited review: regulation of myosin phosphorylation in smooth muscle. *J. Appl. Physiol.* **91**, 497-503.
 Ruppert, C., Deiss, K., Herrmann, S., Vidal, M., Oezkur, M., Gorski, A., Weidemann, F., Lohse, M. J. and Lorenz, K. (2013) Interference with ERK(Thr188) phosphorylation impairs pathological but not physiological cardiac hypertrophy. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 7440-7445.
 Sakurada, S., Takuwa, N., Sugimoto, N., Wang, Y., Seto, M., Sasaki, Y. and Takuwa, Y. (2003) Ca²⁺-dependent activation of Rho and Rho-kinase in membrane depolarization-induced and receptor stimulation-induced vascular smooth muscle contraction. *Circ. Res.* **93**, 548-556.
 Somlyo, A. P. and Somlyo, A. V. (2003) Ca²⁺ sensitivity of smooth muscle and nonmuscle myosin II: modulated by G proteins, kinases, and myosin phosphatase. *Physiol. Rev.* **83**, 1325-1358.

- Tankersley, C. G., Irizarry, R., Flanders, S. E., Rabold, R. and Frank, R. (2003) Unstable heart rate and temperature regulation predict mortality in AKR/J mice. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **284**, R742-R750.
- Taubert, D., Berkels, R., Klaus, W. and Roesen, R. (2002) Nitric oxide formation and corresponding relaxation of porcine coronary arteries induced by plant phenols: essential structural features. *J. Cardiovasc. Pharmacol.* **40**, 701-713.
- Tsai, M. H. and Jiang, M. J. (2006) Rho-kinase-mediated regulation of receptor-agonist-stimulated smooth muscle contraction. *Pflugers Arch.* **453**, 223-232.
- Uehata, M., Ishizaki, T., Satoh, H., Ono, T., Kawahara, T., Morishita, T., Tamakawa, H., Yamagami, K., Inui, J., Maekawa, M. and Narumiya, S. (1997) Calcium sensitization of smooth muscle mediated by a Rho-associated protein kinase in hypertension. *Nature* **389**, 990-994.
- Wier, W. G. and Morgan, K. G. (2003) α 1-Adrenergic signaling mechanisms in contraction of resistance arteries. *Rev. Physiol. Biochem. Pharmacol.* **150**, 91-139.
- Wilson, D. P., Susnjar, M., Kiss, E., Sutherland, C. and Walsh, M. P. (2005) Thromboxane A₂-induced contraction of rat caudal arterial smooth muscle involves activation of Ca²⁺ entry and Ca²⁺ sensitization: Rho-associated kinase-mediated phosphorylation of MYPT1 at Thr-855, but not Thr-697. *Biochem. J.* **389**, 763-774.
- Wooldridge, A. A., MacDonald, J. A., Erdodi, F., Ma, C., Borman, M. A., Hartshorne, D. J. and Haystead, T. A. (2004) Smooth muscle phosphatase is regulated *in vivo* by exclusion of phosphorylation of threonine 696 of MYPT1 by phosphorylation of Serine 695 in response to cyclic nucleotides. *J. Biol. Chem.* **279**, 34496-34504.
- Zou, Q., Leung, S. W. and Vanhoutte, P. M. (2015) Transient receptor potential channel opening releases endogenous acetylcholine, which contributes to endothelium-dependent relaxation induced by mild hypothermia in spontaneously hypertensive rat but not wistar-kyoto rat arteries. *J. Pharmacol. Exp. Ther.* **354**, 121-130.