



Research article

Ginsenoside Rg12, a new dammarane-type triterpene saponin from *Panax ginseng* root



Dong Gu Lee¹, Jaemin Lee¹, Ik-Hyun Cho², Hak-Jae Kim^{3, 4}, Sang-Won Lee⁴, Young-Ock Kim⁴, Chun-Gun Park⁴, Sanghyun Lee^{1,*}

¹ Department of Integrative Plant Science, Chung-Ang University, Anseong, Republic of Korea

² Department of Convergence Medical Science, Brain Korea 21 Plus Program, and Institute of Korean Medicine, College of Oriental Medicine, Kyung Hee University, Seoul, Republic of Korea

³ Department of Clinical Pharmacology, College of Medicine, Soonchunhyang University, Cheonan, Republic of Korea

⁴ Department of Medicinal Crop Research Institute, National Institute of Horticultural and Herbal Science, Rural Development Administration, Eumseong, Republic of Korea

ARTICLE INFO

Article history:

Received 15 April 2016

Received in Revised form

8 September 2016

Accepted 6 October 2016

Available online 11 October 2016

Keywords:

dammarane-type triterpene saponin

ginsenoside Rg12

Panax ginseng

white ginseng

ABSTRACT

Background: *Panax ginseng* has been used as Korean medicine for various diseases. It has antioxidant, hypotensive, sedative, analgesic, and endocrine activities. Dammarane-type triterpenes from the plant have various beneficial effects.

Methods: A dammarane-type triterpene saponin was isolated from *P. ginseng* root through chromatography such as repeated column chromatography and medium pressure liquid chromatography.

Results and conclusion: New dammarane-type triterpene saponin was isolated for the first time from nature. The structure was elucidated as ginsenoside Rg12 (1) based on spectral data. There may be good materials from *P. ginseng* for the development of industrial applications such as nutraceutical, pharmaceutical, and cosmeceutical purposes.

© 2016 The Korean Society of Ginseng, Published by Elsevier Korea LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Panax ginseng (Araliaceae plant) has been used as Korean medicine for several years to treat various diseases [1,2]. Dried ginseng has been used as medicine because it has various pharmacological effects on the central nervous and cardiovascular systems. It is also used for treating diabetes, inflammation aging, fatigue, oxidative damage, mutagenicity, and cancer. Finally, it is used as an antioxidant, hypotensive, sedative, analgesic, and endocrine [3–14].

The majority of *P. ginseng* contains protopanaxadiols (PPDs) and protopanaxatriols (PPTs) as dammarane-type triterpene saponins [15]. The PPDs are ginsenosides-Rb1, -Rb2, -Rd, -Rc, and -Rg3 at the C-3 position sugar moieties, whereas the PPTs are ginsenosides-Rg1, -Re, and -Rg2 at the C-6 position [16].

There have been many recent reports on the conversion of major dammarane-type triterpene saponins to more active minor dammarane-type triterpene saponins, which are in small quantities in ginseng. Current studies demonstrate the beneficial effects of these ginsenosides in a wide range of pathological activities [16,17].

In our continued chemical investigation on *P. ginseng* and dammarane-type triterpene saponins, we isolated and identified phytochemicals from *P. ginseng* root. The compound is purified through repeated column chromatography (CC) and medium pressure liquid chromatography (MPLC).

2. Materials and methods

2.1. Plant materials

The plant of *P. ginseng* Meyer was obtained at Geumsan region, Korea in 2014. A voucher specimen (No. LEE 2011-03) of this plant was deposited at our department.

2.2. Apparatus and chemicals

n-Hexane, *n*-butanol (*n*-BuOH), ethyl acetate (EtOAc), chloroform (CHCl₃), ethanol (EtOH), and pyridine-*d*₅ (MA, USA) were obtained from SamChun Pure Chemical Co., Korea. Fast atom bombardment mass was conducted using a JEOL JMS-AX505WA

* Corresponding author. Department of Integrative Plant Science, Chung-Ang University, 4726, Seodongdaero, Daedukmyun, Anseong 17546, Republic of Korea. E-mail address: slee@cau.ac.kr (S. Lee).

(Jeol, Japan), mass spectrometer. A high-resolution LC/MS/MS analysis was done in a Xevo G2 Q-TOF LC/MS/MS system (Waters, USA) using an ACQUITY UPLC I Class system (Dionex). The ^1H - and ^{13}C -NMR spectra were checked with a Bruker Avance 500 NMR spectrophotometer (Bremen, Germany) with trimethylsilane (TMS), the internal standard. Thin-layer chromatography (TLC) was conducted on Kiesel gel 60 F₂₅₄ (250- μm) silica gel plate (Art. 5715, Merck Co., Darmstadt, Germany), and visualized by a 10% H_2SO_4 spraying in a methanol (MeOH) solution. Accordingly, CC was performed with a LiChroprep RP-18 (40–63 μm , Merck Co.). An MPLC system (Biotage, Uppsala, Sweden), which was equipped with cartridges (KP-SIL, 39 mm \times 225 mm), was used. The sugar determinations were conducted with an HP 5890 series II GC (Hewlett-Packard, Avondale, PA, USA) using an HP-5 capillary column (30 m \times 0.32 mm i.d., 0.25- μm film thickness; Agilent, J&W Scientific, Folsom, CA, USA; injector temperature: 200°C; detector temperature: 200°C; column temperature: 230°C; and flow rate of He gas: 1 mL/min).

2.3. Extraction and isolation

The extraction of *P. ginseng* root (10.0 kg) was performed with EtOH (3 \times 21 L) under reflux. The concentration of the combined extracts was proceeded to have a brown residue (139 g). And then, the residue melted in H_2O (7 L) was successively partitioned with *n*-hexane (3 \times 7 L), CHCl_3 (3 \times 7 L), EtOAc (3 \times 7 L), and *n*-BuOH (3 \times 7 L) to provide the *n*-hexane, CHCl_3 , EtOAc, and *n*-BuOH-soluble fractions. A portion of the *n*-BuOH extract (600 g) was subjected to MPLC for separation using $\text{CHCl}_3/\text{MeOH}$ (gradient: 100:0 \rightarrow 0:100). A total of 13 fractions were obtained by combining those with the same R_f value on the TLC pattern (1 \rightarrow 13). Fraction 3 was separated on a LiChroprep RP18 column (ϕ 1.0 \times 32 cm) using MeOH/ H_2O (gradient: 1:3 \rightarrow 1:0) to obtain 9 fractions (WGB 3.1–3.9). A portion of the combined fractions (WGB 3.8 and WGB 3.9) were separated on a LiChroprep RP18 column (ϕ 1.0 \times 32 cm) using MeOH/ H_2O (gradient: 1:2 \rightarrow 1:0) to obtain 16 fractions (WGB 3.9.1–3.9.16) yielding Compound **1** (WGB 3.9.14).

2.4. Acidic hydrolysis of Compound 1

Compound **1** (10 mg) was heated under reflux with a 5% HCl in 60% aqueous dioxane (10 mL) mixture for 2 h. Under reduced pressure, the mixed solution was concentrated. The residue was then extracted with ether. The H_2O layer was neutralized with Ag_2CO_3 . Subsequently, the remaining solid was removed by filtration. The residue from filtration and standard sugars were compared through cellulose TLC ($\text{C}_5\text{H}_5\text{N}:\text{EtOAc}:\text{HOAc}:\text{H}_2\text{O}$, 36:36:7:21). The sugars were elucidated as D-glucoside.

2.5. Absolute configuration of sugars in Compound 1

Compound **1** (10 mg) was tested as in the above method. The sugar mixture was melted in 0.1 mL $\text{C}_5\text{H}_5\text{N}$, and added to 0.1 mL $\text{C}_5\text{H}_5\text{N}$ solution of 2 mg L-cysteine methyl ester hydrochloride followed by warming at 60°C for 1 h. The solvent was evaporated under N_2 gas. The residue was then dried *in vacuo* and was trimethylsilylated with TMS-HT (0.1 mL) at 60°C for 30 min. The *n*-hexane layer was separated and analyzed by GC after adding *n*-hexane and H_2O to the trimethylsilylated residue. The retention time (t_R) of the peak was 22.03 min as D-glucoside.

3. Results and discussion

The *n*-BuOH fraction was chromatographed by CC and MPLC to yield Compound **1** (Fig. 1).




Fig. 1. Structure of Compound 1.

Compound **1** was gained as a white powder that has a molecular ion peak at m/z 815 $[\text{M}]^-$ in the negative LC-MS. Compound **1** was corresponded to a molecular formula of $\text{C}_{42}\text{H}_{72}\text{O}_{15}$ in HRLC-MS [m/z 861.4843 ($\text{M} + \text{HCOO})^-$]. The calculated value of **1** was m/z 861.4848. The ^1H -NMR spectrum indicated two olefinic (i.e., δ 6.16 and 6.25) and two anomeric (i.e., δ 4.92 and 5.33) proton signals.

Table 1
 ^1H - and ^{13}C -NMR spectral data for Compound 1 ($\text{C}_5\text{D}_5\text{N}$, 500 MHz)

No.	δ_{H}	δ_{C}	HMBC
1	1.55 (2H, m)	39.7	C-3,10,19
2	1.85 (2H, m)	25.9	C-1,3
3	3.27 (1H, dd, 12.0, 4.4)	89.5	C-1',1,28,29
4	—	40.2	C-28, 29
5	0.77 (1H, m)	56.9	—
6	1.49, 1.36 (2H, m)	18.4	—
7	1.21 (1H, m)	35.6	C-8,14,18
8	—	39.7	C-7,18
9	1.36 (1H, m)	49.9	C-11
10	—	36.7	—
11	1.38 (1H, m)	31.2	C-9
12	3.94 (1H, m)	70.7	C-13
13	1.99 (1H, m)	51.9	C-12
14	—	50.7	C-7
15	1.03, 1.57 (2H, m)	31.3	—
16	1.38, 1.80 (2H, m)	26.3	—
17	2.57 (1H, m)	52.2	C-20
18	0.97 (3H, s)	17.1	C-7,8,14
19	0.83 (3H, s)	17.9	C-1,5,10
20	—	83.8	C-17
21	1.59 (3H, s)	25.8	C-17,20,22
22	6.0 (1H, d, 15.9)	127.1	C-20,21,24
23	6.25 (1H, dd, 15.9, 8.4)	137.9	C-24
24	2.22, 2.54 (2H, m)	39.8	C-20,23
25	—	81.9	—
26	1.62 (3H, s)	27.2	—
27	1.57 (3H, s)	18.9	—
28	1.30 (3H, s)	28.6	C-3,4,5,29
29	1.19 (3H, s)	16.5	C-3,4
30	0.97 (3H, s)	16.7	C-8,13,14,15
3-O-glc-1'	4.92 (1H, d, 7.5)	105.6	C-3
2'	4.15 (1H, t)	83.7	C-1''
3'	4.22 (1H, t)	77.6	—
4'	4.05 (1H, t)	72.2	—
5'	3.93 (1H, d)	78.6	—
6'	4.18 (1H, dd, 11.6, 3.2)	63.2	—
	4.36 (1H, dd, 11.6, 6.0)		
2'-O-glc-1''	5.13 (1H, d, 7.5)	106.5	C-2'
2''	4.02 (1H, t)	77.6	—
3''	4.14 (1H, t)	78.6	—
4''	4.17 (1H, t)	72.0	—
5''	4.14 (1H, t)	79.3	—
6''	4.42 (1H, dd, 11.6, 3.2)	64.2	—
	4.50 (1H, dd, 11.6, 6.0)		

HMBC, Heteronuclear Multiple Bond Correlation; delta C is ppm of carbon. Chemical shifts are reported in parts per million (δ), and coupling constants (J) are expressed in Hertz.

The acidic hydrolysis of **1** gained D-glucose. The chemical shifts of the two anomeric carbons in the ^{13}C -NMR spectrum were recorded at δ 105.6 and 106.3 (Table 1). Accordingly, the signals of anomeric carbon showed two β -D-glucosyl moieties. The significant downfield shift of C-2' at δ 79.8 in the inner β -D-glucosyl moiety at C-3 position of aglycone in the ^{13}C -NMR spectrum of C-2' at δ 79.8 indicated the linkage of the terminal β -D-glucosyl moiety to the inner β -D-glucosyl moiety at C-3. The stark difference of the NMR data between the two isomers was the chemical shift values of C-20 and the stereogenic center in the side chain attached to the PPD scaffold and its adjacent carbons, namely, C-17, and 21. In the NMR spectrum of 20-hydroxy-dammarane derivatives, the C-17 and -21 chemical shift values of 20(S)-dammarane derivatives are \sim 52.2 ppm and \sim 25.8 ppm, respectively. From identification of the correlations between H-1' (δ 4.92) and C-3 (δ 89.3) and H-1'' (δ 5.33) and C-2' (δ 79.8) by the HMBC, it was suggested that monodesmosyl chain was linked to the aglycone C-3. Moreover, the correlations were detected between H-24 (i.e., δ 2.22 and 2.54) and C-22 and -23 (i.e., δ 127.0 and 138.7) and H-23 (i.e., δ 6.25) and C-25 (i.e., δ 81.9) by the HMBC [18–24].

Accordingly, Compound **1** is a 20(S)-protopanaxadiol 3-mono-desmoside containing two β -D-glucoside moieties. Therefore, the structure of **1** was elucidated as ginsenoside Rg12. The isolation was for the first time from nature. This result will have valuable effects for the industrial development of ginsenosides from *P. ginseng* in diverse applications.

Conflicts of interest

The authors have no conflicts of interest to declare.

Acknowledgments

This study was carried out with the support of Cooperative Research Program for Agriculture Science and Technology Development (Project No. PJ011582052016), Rural Development Administration, Korea. The authors specifically thank the staff and crew of the National Center for InterUniversity Research Facilities (Seoul National University, Seoul, Korea) for their assistance with the NMR and GC/MS experiments.

References

- [1] Chau CF, Wu SH. The development of regulations of Chinese herbal medicines for both medicinal and food uses. *Trends Food Sci Technol* 2006;17:313–23.
- [2] Coates PM, Blackman MR, Cragg GM, Levine M, Moss J, White JD. *Encyclopedia of dietary supplements*. New York: Marcel Dekker; 2005.
- [3] Kim SH, Park KS. Effects of *Panax ginseng* extract on lipid metabolism in humans. *Pharmacol Res* 2003;48:511–3.
- [4] Sun BS, Gu LJ, Fang ZM, Wang CY, Wang Z, Lee MR, Li Z, Li JJ, Sung CK. Simultaneous quantification of 19 ginsenosides in black ginseng developed from *Panax ginseng* by HPLC-ELSD. *J Pharm Biomed Anal* 2009;50:15–22.
- [5] Chen CF, Chiou WF, Zhang JT. Comparison of the pharmacological effects of *Panax ginseng* and *Panax quinquefolium*. *Acta Pharmacol Sin* 2008;29:1103–8.
- [6] Yue PY, Mak NK, Cheng YK, Leung KW, Ng TB, Fan DT, Yeung HW, Wong RN. Pharmacogenomics and the Yin/Yang actions of ginseng: antitumor, angiomodulating and steroid-like activities of ginsenoside. *Chin Med* 2007;2:1–21.
- [7] Hofseth LJ, Wargovich MJ. Inflammation, cancer, and targets of Ginseng. *J Nutr* 2007;137:183–5.
- [8] Jung CH, Seog HM, Choi IW, Cho HY. Antioxidant activities of cultivated and wild Korean ginseng leaves. *Food Chem* 2005;92:535–40.
- [9] Rai D, Bhatia G, Sen T, Palit GJ. Antistress effects of *Ginkgo biloba* and *Panax ginseng*: a comparative study. *Pharmacol Sci* 2003;93:458–64.
- [10] Surh YJ, Na HK, Lee JY, Keum YS. Molecular mechanisms underlying antitumor promoting activities of heat-processed *Panax ginseng* C. *Korean Med Sci* 2001;16:38–41.
- [11] Choi S. Epidermis proliferative effect of the *Panax ginseng* ginsenoside Rb2. *Arch Pharm Res* 2002;25:71–6.
- [12] Chang LK, Whitaker DC. The impact of herbal medicines on dermatologic surgery. *Dermatol Surg* 2001;27:759–63.
- [13] Keum YS, Park KK, Lee JM, Chun KS, Park JH, Lee SK, Kwon H, Surh YJ. Antioxidant and antitumor promoting activities of the methanol extract of heat-processed ginseng. *Cancer Lett* 2000;150:41–8.
- [14] Attele AS, Wu JA, Yuan C. Ginseng pharmacology: multiple constituents and multiple actions. *Biochem Pharmacol* 1999;58:1685–93.
- [15] Hong HD, Choi SY, Kim YC, Lee YC, Cho CW. Rapid determination of ginsenosides Rb 1, Rf, and Rg 1 in Korean ginseng using HPLC. *J Ginseng Res* 2009;33:8–12.
- [16] Sun J, Hu S, Song X. Adjuvant effects of protopanaxadiol and protopanaxatriol saponins from ginseng roots on the immune responses to ovalbumin in mice. *Vaccine* 2007;25:1114–20.
- [17] Wang JR, Yamasaki Y, Tanaka T, Kouno I, Jiang ZH. Dammarane-type triterpene saponins from the flowers of *Panax notoginseng*. *Molecules* 2009;14:2087–94.
- [18] Kim DS, Chang YJ, Zedk U, Zhao P, Liu YQ, Yang CR. Dammarane saponins from *Panax ginseng*. *Phytochemistry* 1995;40:1493–7.
- [19] Ping Z, Liu YQ, Yang CR. Minor dammarane saponins from *Panax notoginseng*. *Phytochemistry* 1996;41:1419–22.
- [20] Baek SH, Bae ON, Park JH. Recent methodology in ginseng analysis. *J Ginseng Res* 2012;36:119–34.
- [21] Cho JG, Lee MK, Lee JW, Park HJ, Lee DY, Lee YH, Yang DC, Baek NI. Physicochemical characterization and NMR assignments of ginsenosides Rb1, Rb2, Rc, and Rd isolated from *Panax ginseng*. *J Ginseng Res* 2010;34:113–21.
- [22] Yang CM, Seo DS, Hong SH, Kim CH, Lee KR. Ginsenosides from the roots of Korean cultivated-wild ginseng. *Nat Prod Sci* 2008;14:171–6.
- [23] Ahmad VU, Basha A. *Spectroscopic data of saponins: the triterpenoid glycosides*. Florida: CRC Press; 2000. p. 664–717.
- [24] Wang DQ, Feng BS, Wang XB, Yang CR, Zhou J. Further study on dammarane saponins of leaves of *Panax japonicus* var. *major* collected in Quinling Mountains China. *Yao Xue Xue Bao* 1989;24:633–8.