

ANIMAL

Determination of calcium and phosphorus utilization in various hatchery by-products for broiler chickens

Hyeon Seok Choi¹, Geun Hyeon Park¹, Jong Hyuk Kim¹, Sang Yun Ji², Dong Yong Kil^{1*}

¹Department of Animal Science and Technology, Chung-Ang University, Anseong 17546, Korea


²Animal Nutritional Physiology Team, National Institute of Animal Science, Rural Development Administration, Wanju 55365, Korea

*Corresponding author: dongyong@cau.ac.kr

Abstract

The objective of the current study was to determine calcium (Ca) and phosphorus (P) utilization in 4 different hatchery by-products (HBPs) for broiler chickens. The four different HBPs included infertile eggs (IFE), unhatched eggs (UHE), low grade and dead chicks (LDC), and a mixture (MIX) of 55% IFE, 10% UHE, and 10% LDC with 25% hatched eggshells. A total of sixty four 50-day-old Ross 308 broiler chickens were randomly allotted to 1 of 4 dietary treatments with 8 replicates per treatment. Two birds were placed together in one metabolic cage. Additional 16 birds were used to measure the endogenous losses of Ca and P. A force-feeding procedure (i.e., crop intubation) was used to measure the apparent and true total tract retention (ATTR and TTTR, respectively) of Ca and P in the 4 HBPs. The results showed that the TTTR of Ca in the UHE was less ($p < 0.05$) than that in the IFE, LDC, and MIX. The amounts of available Ca in the MIX were greater ($p < 0.05$) than those in the IFE and UHE, which were greater ($p < 0.05$) than those in the LDC. The TTTR of P was not different among the 4 HBPs. However, the amounts of available P in the LDC were greater ($p < 0.05$) than those in the IFE and UHE, which were greater ($p < 0.05$) than those in the MIX. In conclusion, HBPs contain high amounts of available Ca and P because of high concentrations of total Ca and P with a high utilization rate. Therefore, the use of HBPs in broiler diets can reduce costs on Ca and P supplements.

Keywords: broiler chicken, calcium, hatchery by-product, phosphorus



OPEN ACCESS

Citation: Choi HS, Park GH, Kim JH, Ji SY, Kil DY. 2019. Determination of calcium and phosphorus utilization in various hatchery by-products for broiler chickens. Korean Journal of Agricultural Science. <https://doi.org/10.7744/kjoas.20190023>

DOI: <https://doi.org/10.7744/kjoas.20190023>

Received: February 13, 2019

Revised: May 12, 2019

Accepted: May 27, 2019

Copyright: © 2019 Korean Journal of Agricultural Science



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

The rapid growth of poultry industry leads to a large production of poultry by-products (Das et al., 2002). Hatchery by-products (HBPs) are one of the poultry by-products. HBPs include all kinds of by-products produced in poultry hatcheries during the period of hatching and after hatching. Typically, HBPs can be classified into infertile eggs (IFE), unhatched eggs (UHE), low grade and dead chicks (LDC), or hatched eggshells (HE) (Hamm and Whitehead, 1982; Rasool et al., 1999; Shahriar et al.,

2008), although these HBPs are generally collected in a mixture (MIX) of IFE, UHE, LDC, and HE at the end of hatching systems. Currently, most of HBPs have been disposed by spreading on the pasture or dumping in the landfill. However, there is an increasing concern on environmental pollution by the disposal of HBPs. If a large amount of HBPs is applied to the soil, environmental pollutions in soil and water are increased (Tymczynska et al., 2000; Volterra and Conti, 2000). Therefore, the appropriate method of disposing HBPs without harming the environment should be developed.

The HBPs contain high amounts of protein (22 - 56%) and calcium (Ca) (1 - 29%) despite high variations in nutritional compositions. Accordingly, HBPs have gained an increasing attention as a potential ingredient in animal diets (Abiola and Onunkwor, 2004; Shahriar et al., 2008; Abiola et al., 2012). Especially, HBPs contain high amounts of total calcium (Ca) and phosphorus (P), such that inclusion of HBPs in animal diets can reduce the use of Ca and P supplements like limestones (i.e., calcium carbonate) and various calcium phosphate. However, few previous experiments have determined Ca and P utilization in the HBPs as an ingredient for animal diets. In addition, to our knowledge, no previous experiments have evaluated Ca and P utilization in individual HBPs including IFE, UHE, and LDC in animal diets. This lack of information regarding Ca and P utilization in HBPs has led animal nutritionists to hesitate to use HBPs in animal diets.

Therefore, the objective of the current experiment was to determine Ca and P utilization in 4 HBPs including IFE, UHE, LDC, and conventional MIX for broiler chickens as an animal model.

Materials and Methods

Preparation of hatchery by-products

The HBPs used in the current experiment were obtained from a local hatchery (Dongsan broiler hatchery, Cheonan, Korea). The collected HBP samples were separated to IFE, UHE, LDC, or HE. In addition, these HBP samples were mixed according to the proportion of production in the typical hatchery (55% of IFE, 10% of UHE and LDC, and 25% of eggshells) in South Korea to produce the MIX. Each of these 4 HBP samples (IFE, UHE, LDC, and MIX) was dried in a forced-air drying oven at 50°C for 24 h. All dried HBP samples were finely ground through a 1-mm screen grinder (CM 290 Cemotec™, FOSS, Hilleroed, Denmark) before the chemical analyses and metabolic trials.

The HBP samples were measured for gross energy (GE), dry matter (DM), crude ash, crude protein (CP), acid-hydrolyzed ether extract (AEE), Ca, P, and amino acids (AA) (Table 1). These samples were analyzed for DM (method 930.15; AOAC, 2007), crude ash (method 942.05; AOAC, 2007), CP (method 990.03; AOAC, 2007), AEE (method 996.01; AOAC, 2007), and GE using bomb calorimeter (Model 6400, Parr instruments Co., Moline, USA). Benzoic acid was used as the standard for calibration of GE analysis. The HBP samples were also measured for the concentrations of total Ca and P using an inductively coupled plasma spectrometer (ICP) (Optima 5300 DV, Perkin Elmer Inc., Shelton, USA). The concentrations of each AA in the HBP samples were also measured using the high performance liquid chromatography (HPLC; Oh et al., 2011) (Ultimate 3000, Thermo Dionex, Sunnyvale, USA).

Birds, diets, and experimental design

A total of 64 Ross 308 broiler chickens (50-d-old; initial BW = 2.36 ± 0.23 kg) were used. Birds were randomly allotted to 1 of 4 dietary treatments with 8 replicates per treatment. Two birds were placed together in one metabolic cage (35.2 cm × 45.0

cm × 55.3 cm, width × length × height). Room temperature was set at 20°C and the light was provided for 24 h during the entire experiment. Four treatments included 4 individual HBPs of IFE, UHE, LDC, or MIX.

A precision-fed chicken assay was conducted based on the methodology demonstrated in previous experiments (Kim et al., 2011; Lee et al., 2018). Briefly, broiler chickens were obtained at 1 d of age and were fed a commercial broiler diets until 50 d of age. All birds were provided with feed and water ad libitum before the precision-fed procedure began. At the start of the experiment (50 d of age), all birds were fasted for 12 h to empty the gastrointestinal tract (GIT). After 12 h-fasting, broiler chickens were fed 15 g of each HBP by crop intubation, and excreta were collected continuously for 48 h. Additional 16 birds were used to measure endogenous losses of Ca and P. Those birds were also fasted for 12 h and afterward excreta were collected for 48 h. All experimental procedures were reviewed and approved by the Institutional Animal Care and the Use Committee at Chung-Ang University.

Excreta collection and chemical analysis

Collected excreta samples were dried in a forced-air drying oven at 60°C for 48 h and finely ground for the further analysis.

Table 1. Energy and nutrient compositions of hatchery by-products (HBPs; as-fed basis).

Items	IFE	UHE	LDC	MIX
Dry matter (%)	98.3	97.8	96.6	98.0
GE (kcal/kg)	5,133	4,753	5,945	3,764
Crude protein (%)	34.2	39.4	60.0	31.2
AEE (%)	23.3	23.8	24.6	16.7
Crude ash (%)	27.6	27.6	6.6	40.6
Calcium (%)	8.88	10.47	1.25	17.16
Phosphorus (%)	0.49	0.50	0.77	0.31
Essential amino acid (%)				
Arginine	2.09	2.48	3.70	1.54
Histidine	0.84	0.99	1.37	0.64
Isoleucine	1.80	1.88	2.52	1.14
Leucine	3.01	3.25	4.55	1.94
Lysine	2.29	2.44	3.50	1.41
Methionine	1.10	1.10	1.37	0.70
Phenylalanine	2.00	2.03	2.86	1.26
Threonine	1.78	1.91	2.66	1.21
Tryptophan	0.39	0.51	0.63	0.37
Valine	2.27	2.41	3.26	1.51
Non-essential amino acid (%)				
Alanine	2.02	2.28	3.57	1.37
Aspartic acid	3.63	3.83	5.28	2.38
Cysteine	0.64	0.85	1.34	0.64
Glutamic acid	4.72	5.24	7.96	3.25
Glycine	1.28	1.88	4.13	1.27
Proline	1.44	1.85	3.41	1.24
Serine	2.59	2.67	3.48	1.70
Tyrosine	1.57	1.34	2.00	1.11

IFE, infertile eggs; UHE, unhatched eggs; LDC, low grade and dead chicks; MIX, mixture (55% IFE, 10% UHE, 10% LDC, and 25% eggshells); AEE, acid-hydrolyzed ether extract; GE, gross energy.

The excreta from additional 16 birds were also collected and pooled for measuring endogenous losses of Ca and P. Each of HBP samples and excreta samples were analyzed for Ca and P concentrations using ICP as demonstrated by Kurtoğlu et al. (2005).

Calculation

Apparent total tract retention (ATTR) and true total tract retention (TTTR) of Ca and P in 4 HBPs for broiler chickens were calculated as follows (Prola et al., 2013; Lee et al., 2018; Sun and Kim, 2018):

$$X \text{ ATTR} = [(total \text{ X ingested} - total \text{ X excreted}) / total \text{ X ingested}] \times 100,$$

$$X \text{ TTTR} = [(total \text{ X ingested} - total \text{ X excreted} + endogenous \text{ loss of X}) / total \text{ X ingested}] \times 100$$

, where X represents Ca or P.

Statistical analysis

All data were analyzed by ANOVA as a completely randomized design using the GLM procedure of SAS (SAS Institute, Cary, USA). The replicate was used as the experimental unit for all analyses. Outlier data were checked using the UNIVARIATE procedure of SAS (Steel et al., 1997), but no outlier was identified. The means for ATTR and TTTR of Ca and P in various HBP were separated by LSD in multiple comparison procedure of SAS. Significance for statistical tests was set at $p < 0.05$.

Results and Discussion

The energy and nutrient compositions in 4 HBPs are presented in Table 1. For total Ca and P concentrations, total Ca concentrations of HBPs ranged from 1.25 to 17.16%. Those values agreed with the values from 1.04 to 25.62% reported by previous experiments (Abiola and Onunkwor, 2004; Mehdipour et al., 2009; Abiola et al., 2012). Likewise, total P concentrations of HBPs ranged from 0.31 to 0.77%. Those values were placed within the values from 0.01 to 0.84% reported by previous experiments (Rasool et al., 1999; Abiola and Onunkwor, 2004). These results indicate that HBPs used in this experiment are similar to those used in previous experiments. Interestingly, the MIX contained the greatest amounts of Ca, whereas LDC contained the least amounts of Ca. Other HBPs (IFE and UHE) contained intermediate amounts of Ca. In contrast, the LDC contained the greatest amounts of P, whereas the MIX contained the least amounts of P. The reason for these differences in total Ca and P concentrations is likely related to inclusion or presence of eggshells containing high amounts of Ca (36.65%) but low amounts of P (0.16%) (Gongruttananun, 2011). In the current experiment, the MIX was produced with additional eggshells but the LDC included few eggshells.

For the determination of Ca utilization and amounts of available Ca in the HBPs, the ATTR and TTTR of Ca in 4 HBPs were measured (Table 2). The ATTR of Ca in the IFE and MIX (57.0 and 61.4%, respectively) were greater ($p < 0.05$) than in UHE and LDC (44.6 and 35.2%, respectively). The values for TTTR of Ca were calculated by correcting ATTR of Ca with measured endogenous losses of Ca (0.118 g). The TTTR of Ca in the UHE (48.4%) was less ($p < 0.05$) than in the IFE, LDC, and MIX (61.4, 66.5, and 63.7%, respectively). The amount of available Ca in the HBPs was calculated with total Ca concentrations and TTTR of Ca. The amount of available Ca in the MIX (109.2 g/kg) was greater ($p < 0.05$) than those in the IFE and UHE (54.6 and 50.6 g/kg, respectively), which were greater ($p < 0.05$) than those in the LDC (8.3 g/kg).

Our values for the TTTR of Ca ranged from 48.4 to 66.5%. To our knowledge, no previous data are available on the TTTR of