



Neuroprotective effect of *Aster yomena* (Kitam.) Honda against hydrogen peroxide-induced oxidative stress in SH-SY5Y cells

Min Jeong Kim¹ · Ji Hyun Kim² · Sanghyun Lee³ · Eun Ju Cho¹
· Hyun Young Kim²

Received: 6 August 2020 / Accepted: 17 September 2020 / Published Online: 30 September 2020
© The Korean Society for Applied Biological Chemistry 2020

Abstract Oxidative stress is one of the contributors of neurodegenerative disorders including Alzheimer's disease. According to previous studies, *Aster yomena* (Kitam.) Honda (AY) possesses variable pharmacological activities including anti-coagulant and anti-obesity effect. In this study, we aimed to determine the neuroprotective effect of ethyl acetate fraction from *Aster yomena* (Kitam.) Honda (EFAY) against oxidative stress. Therefore, we carried out 3-(4,5-dimethylthiazol-2-yl)-2,3-diphenyl tetrazolium bromide, lactate dehydrogenase (LDH), and 2',7'-dichlorofluorescein diacetate assays in SH-SY5Y neuronal cells treated with hydrogen peroxide (H₂O₂). H₂O₂-treated control cells exhibited reduced viability of cells, and increased LDH release and reactive oxygen species (ROS) production compared to normal cells. However, treatment with EFAY restored the cell viability and inhibited LDH release and ROS production. To investigate the underlying mechanisms by which EFAY attenuated neuronal oxidative damage, we measured protein expressions using Western blot analysis. Consequently, it was observed that EFAY down-regulated cyclooxygenase-2 and interleukin-1 β protein expressions in H₂O₂-treated SH-SY5Y cells that mediated inflammatory

reaction. In addition, apoptosis-related proteins including B-cell lymphoma-2-associated X protein/B-cell lymphoma-2 ratio, cleaved caspase-9, and cleaved-poly (ADP-ribose) polymerase protein expressions were suppressed when H₂O₂-treated cells were exposed to EFAY. Our results indicate that EFAY ameliorated H₂O₂-induced neuronal damage by regulating inflammation and apoptosis. Altogether, AY could be potential therapeutic agent for neurodegenerative diseases.

Keywords Apoptosis · Blotting · Cell culture techniques · Hydrogen peroxide · Oxidative stress · Western

Introduction

As the average life expectancy of the world population is increasing, the incidence of neurodegenerative diseases is also increasing in parallel. According to the World Health Organization, in 2016, about 50 million people lived with dementia and the number is estimated to reach 152 million in 2050 [1]. Neurodegenerative diseases are characterized by progressive dysfunction of neuronal system, in which damaged neurons lead to the decline of cognitive function [2]. Although the mechanism of pathogenesis in neurodegenerative diseases has not been fully revealed, oxidative stress is considered as an important risk factor [3]. Oxidative stress is generated in living organisms when the homeostasis between the production of free radicals and antioxidant defense system gets out of balance [4]. Previous studies demonstrated that oxidative stress exerts cellular damage including DNA injury, inflammation, and apoptosis of brain cells including neuronal, glial, and cerebrovascular cells [5-7]. High concentrations of reactive oxygen species (ROS) can cause damage to cellular lipids, proteins, and DNA because of strong reactivity [8]. Hydrogen peroxide (H₂O₂) is generated in mitochondria during oxygen consumption and reacts with ions of iron or copper to

Eun Ju Cho (✉)
E-mail: ejcho@pusan.ac.kr

Hyun Young Kim (✉)
E-mail: hykim@gntech.ac.kr

¹Department of Food Science and Nutrition, Pusan National University, Busan 46241, Republic of Korea

²Department of Food Science, Gyeongnam National University of Science and Technology, Jinju 52725, Republic of Korea

³Department of Plant Science and Technology, Chung-Ang University, Anseong 17546, Republic of Korea

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

form hydroxyl radical ($\cdot\text{OH}$), which can cause direct damage to proteins and DNA [9,10]. Furthermore, the brain is particularly susceptible to the action of ROS. The brain has high polyunsaturated fatty acids (PUFAs) content and ROS preferentially oxidizes PUFAs resulting in lipid peroxidation [11,12]. These facts imply that the brain produces a large amount of ROS compared to other organs and ROS can induce oxidative damage in the brain.

Aster yomena (Kitam.) Honda (AY) is a perennial plant, widely distributed in Asia. It is used as traditional medicine and edible herb in Korea. According to previous studies, AY possesses variable pharmacological activities such as anti-coagulant and anti-obesity effect [13,14]. It has been demonstrated that AY showed radical scavenging effect through *in vitro* assays against 1,1-diphenyl-2-picrylhydrazyl (DPPH), hydroxyl ($\cdot\text{OH}$), and superoxide ($\text{O}_2\cdot^-$) radicals [15]. According to Bae and Kim [16], ethyl acetate fraction of AY (EFAY) was the most effective fraction with antioxidative activity compared to other extract and fractions and it was believed that there were abundant active compounds in EFAY, with strong pharmacological activities [16]. Nevertheless, until date, none of the studies have examined the neuroprotective effect of EFAY against oxidative stress in neuronal cells. Hence, we aimed to investigate the protective effect of EFAY on H_2O_2 -induced oxidative damage in SH-SY5Y neuronal cells.

Materials and Methods

Materials

Dulbecco's modified eagle's medium (DMEM), penicillin-streptomycin solution, fetal bovine serum (FBS), and trypsin-EDTA solution were obtained from Welgene Inc. (Daegu, Korea). 3-(4,5-Dimethylthiazol-2-yl)-2,3-diphenyl tetrazolium bromide (MTT) was purchased from Bio Basic Inc. (Toronto, Canada). 2',7'-Dichlorofluorescein diacetate (DCFH-DA) was from Sigma Chemical Co. (Saint Louis, MO, USA). Radioimmunoprecipitation (RIPA) buffer was purchased from Elpis Biotech. (Daejeon, Korea) and polyvinylidene fluoride (PVDF) membrane was obtained from Millipore Co. (Billerica, MA, USA). Primary and secondary antibodies were purchased from Cell Signaling Tech. (Beverly, CA, USA), Calbiochem Co. (San Diego, CA, USA), Bioss Inc. (Beijing, China), and Santa Cruz Biotech. (Santa Cruz, CA, USA).

Preparation of sample

AY was obtained from Gurye-gun (Jeollanam-do, Korea) and verified by Dr. K. Choi, Korea National Arboretum, Korea. Voucher specimen (LEE 2016-01) was deposited at Department of Plant Science and Technology, Chung-Ang University, Anseong, Korea. The leaf part of AY (1,645 g) was extracted with methanol (MeOH) at 65-75 °C and refluxed at the same time. After 8 times of repeated extraction, 393.9 g of AY MeOH extract was obtained.

The partitioning of MeOH extract was conducted using n-hexane, dichloromethane, ethyl acetate, and n-butanol. Consequently, 4.2 g of EFAY was obtained.

Cell culture

The SH-SY5Y cell line was obtained from KCLB (Korean Cell Line Bank, Seoul, Korea). SH-SY5Y cells were maintained in DMEM supplemented with 10% FBS and 100 units/mL penicillin-streptomycin. The humidified incubator was set at 37 °C and 5% $\text{CO}_2/95\%$ air. When the adherent cells reached approximately 80% confluence, they were harvested using 0.05% trypsin-EDTA in phosphate buffered saline (PBS, pH 7.4).

MTT assay

Harvested SH-SY5Y cells from T-75 flask were seeded at the density of 5×10^4 cells/mL in 96-well plate. After incubation for 24 h, the cells were treated with EFAY (2.5, 5, and 10 $\mu\text{g}/\text{mL}$) for 2 h and H_2O_2 (300 μM) was added followed by 24 h incubation. Subsequently, the cells were incubated with MTT solution (1 mg/mL) for 3 h and later solubilized by DMSO. The absorbance of each well was read at 540 nm [17].

Lactate dehydrogenase (LDH) release assay

LDH release assay was carried out employing an LDH cytotoxicity detection kit (Takara Bio Inc., Shiga, Japan). SH-SY5Y cells were harvested from T-75 flask and plated at a density of 5×10^4 cells/mL in 96-well plate followed by 24 h incubation. The cells were treated with EFAY (2.5, 5, and 10 $\mu\text{g}/\text{mL}$) for 2 h. Subsequently, H_2O_2 (300 μM) was added and the cells were incubated for 24 h. The supernatant was mixed with a reaction solution in a 96-well plate, and then incubated for 30 min at room temperature. The amount of LDH release was measured using a microplate reader at 490 nm.

DCFH-DA assay

SH-SY5Y cells were plated in 96-well black plate at a density of 5×10^4 cells/mL and incubated for 24 h. Before treating with H_2O_2 (300 μM) for 24 h, EFAY (2.5, 5, and 10 $\mu\text{g}/\text{mL}$) was pretreated with the cells for 2 h. After incubation with DCFH-DA (80 μM) for 30 min, fluorescence was measured at excitation - 480 nm and emission - 535 nm for 60 min.

Western blot analysis

Proteins were extracted using RIPA buffer supplemented with 1X protease inhibitor cocktail. These proteins were electrophoresed on 10-13% sodium dodecyl sulfate-polyacrylamide gels and then transferred to PVDF membranes. The membranes were blocked with 5% skim milk in PBS with tween 20 for 50 min and incubated overnight at 4 °C with primary antibodies. The primary antibodies used were: β -actin (1:1000; Cell Signaling Tech.); COX-2 (1:1000; Calbiochem Co.); IL-1 β (1:1000; Bioss Inc.); cleaved caspase-9 (1:1000; Cell Signaling Tech.); cleaved PARP

(1:500; Cell Signaling Tech.); Bax (1:500; Santa Cruz Biotech.); and Bcl-2 (1:500; Santa Cruz Biotech.). After incubation with appropriate HRP-conjugated secondary antibodies for 1 h, immunoreactive proteins were visualized using a chemiluminescent imaging system (CoreBio, Seoul, Korea).

Statistical analysis

The results of this study are shown as the mean \pm standard deviation (SD). The significance of data was determined by one-way analysis of variance (ANOVA) and Duncan's multiple range test for multiple comparisons. Statistical significance was considered when p was less than 0.05.

Results

Effect of EFAY on cell viability in SH-SY5Y cells treated with H₂O₂

The cell viability was measured to determine the protective effect of EFAY on H₂O₂-treated SH-SY5Y cells by performing MTT assay. As illustrated in Fig. 1, H₂O₂ reduced cell viability to 40.75% compared with 100% of normal cells. However, when SH-SY5Y cells were treated with EFAY, cell viability was restored to up to 45.50% at the concentration of 10 μ g/mL.

Effect of EFAY on LDH release in SH-SY5Y cells treated with H₂O₂

LDH release assay was used to evaluate the protective effect of EFAY on neuronal damage induced by H₂O₂ in SH-SY5Y cells.

A significant increase in the amount of released LDH was observed in H₂O₂-treated cells compared with normal cells (57.37%), as shown in Fig. 2. Meanwhile, the treatment with EFAY significantly decreased the release of LDH compared to control cells, indicating inhibition of neuronal cytotoxicity induced by H₂O₂ in SH-SY5Y cells.

Effect of EFAY on ROS production in SH-SY5Y cells treated with H₂O₂

In order to investigate the neuroprotective effect of EFAY against ROS production, DCFH-DA assay was carried out employing H₂O₂-treated SH-SY5Y cells. Figure 3 shows the effect of EFAY on H₂O₂ mediated production of ROS in SH-SY5Y cells (100%) compared with the non-treated group (79.38%). Treatment with 10 μ g/mL of EFAY resulted in a decrease in ROS production by 85.99%. The results demonstrated that ROS production caused by H₂O₂ was blocked by EFAY with subsequent reduction in neuronal cell death.

Effects of EFAY on inflammation-related protein expressions in H₂O₂-induced SH-SY5Y cells

To evaluate the effect of EFAY on inflammatory processes in H₂O₂-treated SH-SY5Y cells, Western blot analysis was performed. As shown in Fig. 4, protein levels of interleukin-1 β (IL-1 β) and cyclooxygenase-2 (COX-2) were increased in the presence of H₂O₂ as compared to normal cells, indicating that H₂O₂ induced inflammation in neuronal cells. However, treatment with EFAY significantly down-regulated protein levels of IL-1 β and COX-2 compared with H₂O₂-treated control cells. In particular, EFAY




Fig. 1 Effect of ethyl acetate fraction from *Aster yomena* (Kitam.) Honda on cell viability in SH-SY5Y cells treated with H₂O₂. SH-SY5Y cells were treated with EFAY (2.5, 5, and 10 μ g/mL) for 2 h and H₂O₂ (300 μ M) was added followed by 24 h incubation. H₂O₂-untreated cells served as a normal group. H₂O₂-treated cells served as a control group. Values are mean \pm SD. ^{a-e}Means with the different letters are significantly different ($p < 0.05$) by Duncan's multiple range test




Fig. 2 Effect of ethyl acetate fraction from *Aster yomena* (Kitam.) Honda on LDH release in SH-SY5Y cells treated with H₂O₂. SH-SY5Y cells were treated with EFAY (2.5, 5, and 10 μ g/mL) for 2 h and H₂O₂ (300 μ M) was added followed by 24 h incubation. H₂O₂-untreated cells served as a normal group. H₂O₂-treated cells served as a control group. Values are mean \pm SD. ^{a-e}Means with the different letters are significantly different ($p < 0.05$) by Duncan's multiple range test




Fig. 3 Effects of ethyl acetate fraction from *Aster yomena* (Kitam.) Honda on ROS production in SH-SY5Y cells treated with H₂O₂. (A) Change of ROS fluorescence during 60 min; (B) The intensity of ROS fluorescence at 60 min. SH-SY5Y cells were treated with EFAY (2.5, 5, and 10 µg/mL) for 2 h and H₂O₂ (300 µM) was added followed by 24 h incubation. H₂O₂-untreated cells served as a normal group. H₂O₂-treated cells served as a control group. Values are mean ± SD. ^{a-d}Means with the different letters are significantly different ($p < 0.05$) by Duncan's multiple range test




Fig. 4 Effects of ethyl acetate fraction from *Aster yomena* (Kitam.) Honda on inflammation-related protein expressions in H₂O₂-treated SH-SY5Y cells. SH-SY5Y cells were treated with EFAY (2.5, 5, and 10 µg/mL) for 2 h and H₂O₂ (300 µM) was added followed by 24 h incubation. β-actin was used as the loading control. H₂O₂-untreated cells served as a normal group. H₂O₂-treated cells served as a control group. Values are mean ± SD. ^{a-e}Means with the different letters are significantly different ($p < 0.05$) by Duncan's multiple range test

effectively reduced COX-2 protein expression in H₂O₂-treated SH-SY5Y cells. These results suggested that EFAY protected H₂O₂-induced neuronal inflammation by down-regulating IL-1β and COX-2 expressions.




Fig. 5 Effect of ethyl acetate fraction from *Aster yomena* (Kitam.) Honda on the level of Bax/Bcl-2 in H₂O₂-treated SH-SY5Y cells. SH-SY5Y cells were treated with EFAY (2.5, 5, and 10 µg/mL) for 2 h and H₂O₂ (300 µM) was added followed by 24 h incubation. β-Actin was used as the loading control. H₂O₂-untreated cells served as a normal group. H₂O₂-treated cells served as a control group. Values are mean ± SD. ^{a-e}Means with the different letters are significantly different ($p < 0.05$) by Duncan's multiple range test

Effects of EFAY on apoptosis-related protein expressions in H₂O₂-induced SH-SY5Y cells

We investigated the effect of EFAY on apoptosis pathway in SH-




Fig. 6 Effects of ethyl acetate fraction from *Aster yomena* (Kitam.) Honda on apoptosis-related protein expressions in H₂O₂-treated SH-SY5Y cells. SH-SY5Y cells were treated with EFAY (2.5, 5, and 10 µg/mL) for 2 h and H₂O₂ (300 µM) was added followed by 24 h incubation. β-Actin was used as the loading control. H₂O₂-untreated cells served as a normal group. H₂O₂-treated cells served as a control group. Values are mean ± SD. ^{a-c}Means with the different letters are significantly different (*p* < 0.05) by Duncan’s multiple range test

SH-SY5Y cells under oxidative stress induced by H₂O₂. Figure 5 illustrates that H₂O₂ significantly up-regulated B-cell lymphoma-2-associated X protein (Bax)/B-cell lymphoma-2 (Bcl-2) ratio. Furthermore, the protein levels of cleaved caspase-9 and cleaved poly (ADP-ribose) polymerase (PARP) were down-regulated compared to the normal cells, as can be seen in Fig. 6. However, treatment with EFAY significantly down-regulated Bax/Bcl-2 ratio and up-regulated cleaved caspase-9 and cleaved PARP protein levels. These results indicated that the treatment of H₂O₂-induced damaged SH-SY5Y cells with EFAY suppressed apoptosis pathway by modulating Bax/Bcl-2 ratio, cleaved caspase-9, and cleaved PARP.

Discussion

Neurodegenerative diseases are characterized by abnormal deposition of specific proteins in the brain, which could cause neuronal dysfunction and cell death [18]. For example, in the brain of a patient with Alzheimer’s disease, amyloid beta plaque and the neurofibrillary tangles are commonly observed as pathological markers [19]. The major lesion of Parkinson’s disease is selective

degeneration of dopaminergic neurons around substantia nigra [20]. Previous studies demonstrated that the modification and abnormal deposition of proteins in the brain are closely related to oxidative stress resulting from overgeneration of ROS [21]. Therefore, the accumulation of oxidized protein is dependent upon the balance between protein oxidation and degradation of the oxidized protein. A battery of ROS, including free radicals (such as ·OH and O₂⁻) and non-radical oxygen derivatives (such as H₂O₂ and ONOO⁻), contributes in maintaining the balance by promoting protein oxidation [22]. H₂O₂, one of the oxygen derivatives, is generated from O₂⁻, catalyzed by superoxide dismutase in the mitochondria. When the concentration of H₂O₂ exceeds that of antioxidant enzymes, H₂O₂ undergoes ferrous ion-catalyzed cleavage process by the Fenton reaction to produce ·OH [23]. The ·OH is one of the critical free radicals in the body because it could indiscriminately damage targets and mediate tissue damage consequently [24]. H₂O₂ can also induce neuronal cell death in SH-SY5Y cells *via* inflammatory and apoptotic processes [25,26]. Thereby, SH-SY5Y cells are widely used to study the pathogenesis of neurodegeneration. It was reported that H₂O₂ activated nuclear translocation of nuclear factor-kappa B (NF-κB) in SH-SY5Y cells and it up-regulated COX-2 and inducible nitric oxide synthase (iNOS) in the inflammatory pathway [26]. Our previous study also demonstrated that H₂O₂ activated apoptosis pathway in SH-SY5Y cells and several therapeutic agents reduced the cell death [5].

In our previous study, AY exhibited antioxidant effects by suppressing generation of free radicals including DPPH, ·OH, and O₂⁻ radicals [15]. It has been demonstrated that phenolic compounds from AY also had antioxidant activity. The phenolic compounds in AY are esculetin, caffeic acid, and apigenin [27]. Esculetin showed antioxidant activity by regulating glutathione system and lipid peroxidation in liver supernatants from male C57BL/6J mice [28]. Caffeic acid had antioxidative effects on intestinal ischemia-reperfusion injury in rats [29]. Especially, apigenin showed neuroprotective effects from H₂O₂-induced oxidative damage in SH-SY5Y cells [30]. However, the protective effect of AY from oxidative stress in neuronal cells has not been revealed yet. Therefore, we examined whether EFAY has a neuroprotective effect against oxidative stress that is induced by treating SH-SY5Y cells with H₂O₂.

It has been observed that the treatment of SH-SY5Y cells with H₂O₂ induced cellular toxicity and subsequently cell death [31]. We confirmed that H₂O₂ suppressed cell viability of SH-SY5Y cells based on MTT assay. According to previous study, ethanol extract from AY (EEAY) showed no cytotoxicity at concentration up to 500 µg/mL [32]. EEAY decreased adipocyte differentiation in the 3T3-L1 cells by suppressing adipogenic transcriptional factors without cytotoxicity at the concentration of 200 µg/mL [13]. Furthermore, our previous study unveiled that EFAY also reduced adipogenesis at a concentration up to 100 µg/mL [33]. Therefore, EFAY was treated under 100 µg/mL SH-SY5Y cells,

that had both no cytotoxicity and beneficial effect against oxidative stress. Furthermore, cellular damage by the cytotoxic stimulus, results in leaking of LDH, an enzyme normally found in cytosol into the extracellular fluid because of the disturbed cell membrane [34]. Our data showed that the release of LDH was increased in H₂O₂-treated SH-SY5Y cells and reversed by EFAY treatment in oxidative stress-induced SH-SY5Y cells. These findings revealed that EFAY exhibited a cytoprotective effect in SH-SY5Y cells against neuronal damage induced by H₂O₂.

Previous studies stated that treatment with H₂O₂ generates ROS (e.g., ·OH), which is one of the triggers leading to apoptotic signaling in SH-SY5Y cells [35,36]. The production of ROS could be monitored by measuring fluorescence emission following the conversion of DCFH-DA into DCF in the presence of ROS in cells [37]. Our results showed that H₂O₂ increased the ROS production in SH-SY5Y cells. However, EFAY significantly decreased the levels of ROS that were generated by H₂O₂. We previously confirmed antioxidative effect of AY by DPPH, ·OH, and O₂⁻ assay [15]. These data indicated that EFAY inhibited H₂O₂-induced oxidative stress by reducing ROS production in SH-SY5Y cells.

In the central nervous system (CNS), the up-regulation of IL-1β is a patterned response after CNS insults such as infection and trauma [38]. IL-1β stimulates NF-κB translocation into the nuclear region, which increases pro-inflammatory gene expressions including COX-2 and iNOS [39]. COX-2 is considered as a major regulator of the inflammatory process, and neuronal inflammation, which is a hall mark of cognitive dysfunction [40]. Many studies reported that COX-2 participated in prostaglandin synthesis, which mediated pathogenic mechanism such as inflammatory response [41,42]. In the brain, COX-2 also produces prostaglandins and subsequently induces neuronal toxicity, which can lead to neurological diseases [43,44]. Previously, it has been demonstrated that treatment of H₂O₂ activated phosphorylation of NF-κB at p65 in cells induced the productions of COX-2 and IL-1β [26]. In the present study, it was confirmed that H₂O₂ up-regulated the protein expressions of IL-1β and COX-2 in SH-SY5Y cells, suggesting that H₂O₂ activated inflammatory reaction in SH-SY5Y cells. On the contrary, treatment with EFAY down-regulated the expressions of IL-1β and COX-2. Earlier, it has been revealed that several phenolic compounds present in EFAY dysregulated interleukin-6 production in TNF-α stimulated MG-63 cells, proposing their role as anti-inflammatory agents [45]. Furthermore, EFAY lowered ovalbumin- and LPS-induced inflammation by down-regulating NF-κB and iNOS in RAW 264.7 cells [46]. These findings suggested that EFAY ameliorated H₂O₂-induced neuronal inflammation by regulating IL-1β and COX-2 expression.

When cells are exposed to cellular stress, pro-apoptotic BH3-only proteins directly activate pro-apoptotic proteins (e.g., Bax and Bak) and inactivate anti-apoptotic proteins (e.g., prosurvival Bcl-2-like proteins), leading to indirect activation of Bax and Bak. The activated Bax is oligomerized and translocated to the

mitochondrial outer membrane, resulting in the release of cytochrome C [47]. Bcl-2 is one of the prosurvival Bcl-2-like proteins and protects apoptosis by interfering with the release of cytochrome C from mitochondria [48,49]. The released cytochrome C leads to caspases activation that plays a key role at the various stages of the apoptotic process [50]. Caspase-9, one of the caspase family, subsequently promotes downstream caspases and finally activates PARP leading to ultimate cell death [51]. Excessive oxidative stress leads to neuronal apoptosis and H₂O₂ has been used as an inducer of oxidative stress in SH-SY5Y cells [52]. Previous studies demonstrated that treatment with H₂O₂ led to the formation of apoptotic features such as decreased Bax/Bcl-2 ratio, activation of the caspase cascade, and DNA fragmentation in SH-SY5Y cells [31,53,54]. Our results showed that H₂O₂-mediated neuronal apoptosis was activated in SH-SY5Y cells by up-regulating Bax/Bcl-2 ratio, whereas EFAY treatment down-regulated Bax/Bcl-2 ratio. Furthermore, up-regulated protein expressions of cleaved caspase-9 and cleaved PARP were observed subsequent to treatment of SH-SY5Y cells with H₂O₂. However, treatment with EFAY also down-regulated cleaved caspase-9 and cleaved PARP expressions, suggesting that EFAY exerts a protective effect against apoptosis in H₂O₂-treated neuronal cells. In addition, isoquercitrin, one of the active compounds of EFAY acts as an antioxidant and regulates apoptosis-related protein expressions of Bcl-2 and cleaved PARP in H₂O₂-treated RGC-5 cells [55]. Isoquercitrin was also reported to inhibit H₂O₂-induced apoptosis of EA.hy926 cells by mediating the PI3K/Akt/GSK3β signaling [56]. These results indicated that EFAY exerts a protective effect on the apoptotic effect against H₂O₂-induced neuronal damage in SH-SY5Y cells by down-regulating Bax/Bcl-2 ratio and caspase cascades.

Our findings proposed that EFAY recovered the cell viability and diminished release of LDH by suppressing the production of ROS against H₂O₂-mediated neuronal damage in SH-SY5Y cells. EFAY attenuated inflammation-related protein expressions including IL-1β and COX-2. Moreover, EFAY alleviated apoptosis by regulating Bax/Bcl-2 ratio, cleaved caspase-9, and cleaved PARP protein expressions. Based on the obtained results, the inhibitory effect of EFAY on inflammation and apoptosis pathways is apparent. In this study, we investigated neuroprotective effect of AY *via* inflammatory and apoptosis pathway. According to previous study, AY inhibited HO-induced growth inhibition in RAW 264.7 cells, which was associated with expression of nuclear factor erythroid 2-related factor-2 (Nrf-2) and heme oxygenase-1 (HO-1) [56]. Nrf-2 and its downstream mediator, HO-1, are important molecules in host defense against oxidative injury. Nrf-2 is a cytoprotective factor regulating the expression of genes coding for antioxidant proteins such as HO-1 [57]. HO-1 exerts beneficial effects through the elimination of toxic heme and production of biliverdin, iron ions, and carbon monoxide [58]. Since Nrf-2/HO-1 pathway is confused under neurodegenerative disease such as AD, the improvement of Nrf-2/HO-1 by AY

treatment leads to neuronal protection. Taken together, AY protects neuronal cells against oxidative stress and is proposed as an effective natural agent for the prevention of neurodegenerative disorders.

Acknowledgments This research was funded by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education [2016R1D1A1B03931593].

References

- Alzheimer's Association (2017) 2017 Alzheimer's disease facts and figures. *Alzheimers Dement* 13: 325–373
- Radi E, Formichi P, Battisti C, Federico A (2004) Apoptosis and oxidative stress in neurodegenerative diseases. *J Alzheimers Dis* 42: S125–S152
- Uttara B, Singh AV, Zamboni P, Mahajan RT (2009) Oxidative stress and neurodegenerative diseases: a review of upstream and downstream antioxidant therapeutic options. *Curr Neuropharmacol* 7: 65–74
- Foyer CH, Noctor G (2005) Redox homeostasis and antioxidant signaling: a metabolic interface between stress perception and physiological responses. *Plant Cell* 17: 1866–1875
- Lee AY, Choi JM, Lee MH, Lee J, Lee S, Cho EJ (2018) Protective effects of perilla oil and alpha linolenic acid on SH-SY5Y neuronal cell death induced by hydrogen peroxide. *Nutr Res Pract* 12: 93–100
- Choi EO, Jeong JW, Park C, Hong SH, Kim GY, Hwang HJ, Cho EJ, Choi YH (2016) Baicalein protects C6 glial cells against hydrogen peroxide-induced oxidative stress and apoptosis through regulation of the Nrf2 signaling pathway. *Int J Mol Med* 37: 798–806
- Zhu X, Smith MA, Honda K, Aliev G, Moreira PI, Nunomura A, Casadesus G, Harris PL, Siedlak SL, Perry G (2007) Vascular oxidative stress in Alzheimer disease. *J Neurol Sci* 257: 240–246
- Finkel T, Holbrook NJ (2000) Oxidants, oxidative stress and the biology of ageing. *Nature* 408: 239–247
- Barbouti A, Doulias PT, Nouis L, Tenopoulou M, Galaris D (2002) DNA damage and apoptosis in hydrogen peroxide-exposed Jurkat cells: bolus addition versus continuous generation of H₂O₂. *Free Radic Biol Med* 33: 691–702
- Stohs SJ, Bagchi D (1995) Oxidative mechanisms in the toxicity of metal ions. *Free Radic Biol Med* 18: 321–336
- Shichiri M (2014) The role of lipid peroxidation in neurological disorders. *J Clin Biochem Nutr* 54: 151–160
- Gaschler MM, Stockwell BR (2017) Lipid peroxidation in cell death. *Biochem Biophys Res Commun* 482: 419–425
- Han MH, Jeong JS, Jeong JW, Choi SH, Kim SO, Hong SH, Park C, Kim BW, Choi YH (2017) Ethanol extracts of *Aster yomena* (Kitam.) Honda inhibit adipogenesis through the activation of the AMPK signaling pathway in 3T3-L1 preadipocytes. *Drug Discov Ther* 11: 281–287
- Choi JH, Kim DW, Park SE, Choi BS, Sapkota K, Kim S, Kim SJ (2014) Novel thrombolytic protease from edible and medicinal plant *Aster yomena* (Kitam.) Honda with anticoagulant activity: purification and partial characterization. *J Biosci Bioeng* 118: 372–377
- Kim MJ, Kim JH, Lee SH, Cho EJ, Kim HY (2018) Determination of radical scavenging activity of *Aster yomena* (Kitam.) Honda. *J Korea Acad Industr Coop Soc* 19: 402–407
- Bae JS, Kim TH (2009) Acetylcholinesterase inhibitory and antioxidant properties of *Aster yomena* extract. *Korea J Herbol* 24: 121–126
- Mosmann T (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 65: 55–63
- Sayre LM, Smith MA, Perry G (2001) Chemistry and biochemistry of oxidative stress in neurodegenerative disease. *Curr Med Chem* 8: 721–738
- Hardy JA, Higgins GA (1992) Alzheimer's disease: the amyloid cascade hypothesis. *Science* 256: 184–185
- Warner TT, Schapira AH (2003) Genetic and environmental factors in the cause of Parkinson's disease. *Ann Neurol* 53: S16–23 discussion S23–25
- Stadtman ER, Berlett BS (1997) Reactive oxygen-mediated protein oxidation in aging and disease. *Chem Res Toxicol* 10: 485–494
- Berlett BS, Stadtman ER (1997) Protein oxidation in aging, disease, and oxidative stress. *J Biol Chem* 272: 20313–20316
- Liu D, Liu J, Wen J (1999) Elevation of hydrogen peroxide after spinal cord injury detected by using the Fenton reaction. *Free Radic Biol Med* 27: 478–482
- Slivka A, Cohen G (1985) Hydroxyl radical attack on dopamine. *J Biol Chem* 260: 15466–15472
- Filomeni G, Piccirillo S, Rotilio G, Ciriolo MR (2012) p38(MAPK) and ERK1/2 dictate cell death/survival response to different pro-oxidant stimuli via p53 and Nrf2 in neuroblastoma cells SH-SY5Y. *Biochem Pharmacol* 83: 1349–1357
- Chetsawang B, Putthaprasart C, Phansuwan-Pujito P, Govitrapong P (2006) Melatonin protects against hydrogen peroxide-induced cell death signaling in SH-SY5Y cultured cells: involvement of nuclear factor kappa B, Bax and Bcl-2. *J Pineal Res* 41: 116–123
- Kim AR, Jin Q, Jin HG, Ko HJ, Woo ER (2014) Phenolic compounds with IL-6 inhibitory activity from *Aster yomena*. *Arch Pharm Res* 37: 845–851
- Martín-Aragón S, Benedí JM, Villar AM (1998) Effects of the antioxidant (6, 7-dihydroxycoumarin) esculetin on the glutathione system and lipid peroxidation in mice. *Gerontology* 44: 21–25
- Sato Y, Itagaki S, Kurokawa T, Ogura J, Kobayashi M, Hirano T, Sugawara M, Iseki K. (2011) *In vitro* and *in vivo* antioxidant properties of chlorogenic acid and caffeic acid. *Int J Pharm* 403: 136–138
- Kang SS, Lee JY, Choi YK, Kim GS, Han BH (2004) Neuroprotective effects of flavones on hydrogen peroxide-induced apoptosis in SH-SY5Y neuroblastoma cells. *Bioorg Med Chem Lett* 14: 2261–2264
- Kwon SH, Kim JA, Hong SI, Jung YH, Kim HC, Lee SY, Jang CG (2011) Loganin protects against hydrogen peroxide-induced apoptosis by inhibiting phosphorylation of JNK, p38, and ERK 1/2 MAPKs in SH-SY5Y cells. *Neurochem Int* 58: 533–541
- Kang HJ, Jeong JS, Park NJ, Go GB, Kim SO, Park C, Kim BW, Hong SH, Choi YH (2017). An ethanol extract of *Aster yomena* (Kitam.) Honda inhibits lipopolysaccharide-induced inflammatory responses in murine RAW 264.7 macrophages. *Biosci Trends* 11: 85–94
- Kim MJ, Kim JH, Lee S, Kim HY, Cho EJ (2019) *Aster yomena* (Kitam.) Honda inhibits adipocyte differentiation in 3T3-L1 cells. *Int J Gerontol* 5: S33–S38
- Arechabala B, Coiffard C, Rivalland P, Coiffard LJM, Roeckholtzhauer YD (1999) Comparison of cytotoxicity of various surfactants tested on normal human fibroblast cultures using the neutral red test, MTT assay and LDH release. *J Appl Toxicol* 19: 163–165
- Liu X, Shibata T, Hisaka S, Osawa T (2009) Astaxanthin inhibits reactive oxygen species-mediated cellular toxicity in dopaminergic SH-SY5Y cells via mitochondria-targeted protective mechanism. *Brain Res* 1254: 18–27
- Venuprasad MP, Kumar HK, Khanum F (2013) Neuroprotective effects of hydroalcoholic extract of *Ocimum sanctum* against H₂O₂ induced neuronal cell damage in SH-SY5Y cells via its antioxidative defence mechanism. *Neurochem Res* 38: 2190–2200
- Cohn CA, Simon SR, Schoonen MA (2008) Comparison of fluorescence-based techniques for the quantification of particle-induced hydroxyl radicals. *Part Fibre Toxicol* 5: 2
- Shafelt SS, Kyrkanides S, Olschowka JA, Jennie HM, Johnson RE, O'Banion MK (2007) Sustained hippocampal IL-1 β overexpression mediates chronic neuroinflammation and ameliorates Alzheimer plaque

- pathology. *J Clin Invest* 117: 1595–1604
39. Wang Z, Nakayama T (2010) Inflammation, a link between obesity and cardiovascular disease. *Mediators Inflamm* 2010: 535918
 40. Lee JW, Lee YK, Yuk DY, Choi DY, Ban SB, Oh KW, Hong JT (2008) Neuro-inflammation induced by lipopolysaccharide causes cognitive impairment through enhancement of beta-amyloid generation. *J Neuroinflammation* 5: 37
 41. Berg J, Fellier H, Christoph T, Grarup J, Stimmeder D (1999) The analgesic NSAID lornoxicam inhibits cyclooxygenase (COX)-1/-2, inducible nitric oxide synthase (iNOS), and the formation of interleukin (IL)-6 *in vitro*. *Inflamm Res* 48: 369–379
 42. Ricciotti E, FitzGerald GA (2011) Prostaglandins and inflammation. *Arterioscler Thromb Vasc Biol* 31: 986–1000
 43. Manabe Y, Anrather J, Kawano T, Niwa K, Zhou P, Ross ME, Iadecola C (2004) Prostanoids, not reactive oxygen species, mediate COX-2-dependent neurotoxicity. *Ann Neurol* 55: 668–675
 44. Liang X, Wu L, Wang Q, Hand T, Bilak M, McCullough L, Andreasson K (2007) Function of COX-2 and prostaglandins in neurological disease. *J Mol Neurosci* 33: 94–99
 45. Kim AY, Shin HM, Kim JS, Shim HJ, Nam KW, Hwang KA, Youn HS (2017) Anti-inflammatory effects of *Aster yomena* extracts by the suppression of inducible nitric oxide synthase expression. *The Biomed Sci Lett* 23: 104–110
 46. Westphal D, Dewson G, Czabotar PE, Kluck RM (2011) Molecular biology of Bax and Bak activation and action. *Biochim Biophys Acta* 1813: 521–531
 47. Rossé T, Olivier R, Monney L, Rager M, Conus S, Fellay I, Jansen B, Borner C (1998) Bcl-2 prolongs cell survival after Bax-induced release of cytochrome c. *Nature* 391: 496–499
 48. Yang J, Liu X, Bhalla K, Kim CN, Ibrado AM, Cai J, Peng TI, Jones DP, Wang X (1997) Prevention of apoptosis by Bcl-2: release of cytochrome c from mitochondria blocked. *Science* 275: 1129–1132
 49. Grütter MG (2000) Caspases: key players in programmed cell death. *Curr Opin Struct Biol* 10: 649–655
 50. Mattson MP (2000) Apoptosis in neurodegenerative disorders. *Nat Rev Mol Cell Biol* 1: 120–129
 51. Satoh T, Sakai N, Enokido Y, Uchiyama Y, Hatanaka H (1996) Free radical-independent protection by nerve growth factor and Bcl-2 of PC12 cells from hydrogen peroxide-triggered apoptosis. *J Biochem* 120: 540–546
 52. Stridh H, Kimland M, Jones DP, Orrenius S, Hampton MB (1998) Cytochrome c release and caspase activation in hydrogen peroxide- and tributyltin-induced apoptosis. *FEBS Lett* 429: 351–355
 53. Chen L, Liu L, Yin J, Luo Y, Huang S (2009) Hydrogen peroxide-induced neuronal apoptosis is associated with inhibition of protein phosphatase 2A and 5, leading to activation of MAPK pathway. *Int J Biochem Cell Biol* 41: 1284–1295
 54. Jung SH, Kim BJ, Lee EH, Osborne NN (2010) Isoquercitrin is the most effective antioxidant in the plant *Thuja orientalis* and able to counteract oxidative-induced damage to a transformed cell line (RGC-5 cells). *Neurochem Int* 57: 713–721
 55. Zhu M, Li J, Wang K, Hao X, Ge R, Li Q (2016) Isoquercitrin inhibits hydrogen peroxide-induced apoptosis of EA.hy926 cells via the PI3K/Akt/GSK3 β signaling pathway. *Molecules* 21: 356
 56. Loboda A, Damulewicz M, Pyza E, Jozkiewicz A, Dulak J (2016) Role of Nrf2/HO-1 system in development, oxidative stress response and diseases: an evolutionarily conserved mechanism. *Cell Mol Life Sci* 73: 3221–3247
 57. Li C, Wang R, Hu C, Wang H, Ma Q, Chen S, He Y (2018) Pyridoxine exerts antioxidant effects in cell model of Alzheimer's disease *via* the Nrf-2/HO-1 pathway. *Cell Mol Biol* 64: 119–124
 58. Kim SO, Jeong JS, Choi YH (2019) Antioxidant and anti-inflammatory effects of ethanol extract of *Aster yomena* in RAW 264.7 macrophages. *J Life Sci* 29: 977–985