

Prognostic factors for neurological outcomes in Korean targeted temperature management recipients with return of spontaneous circulation after out-of-hospital cardiac arrests

A nationwide observational study

Jae Guk Kim, MD^{a,b}, Hyungoo Shin, MD^c, Hyun Young Choi, MD, PhD^{a,*}, Wonhee Kim, MD, PhD^a, Jihoon Kim, MD, PhD^d, Shinje Moon, MD^e, Bongyoung Kim, MD, PhD^f, Chiwon Ahn, MD^g, Juncheol Lee, MD^h

Abstract

Targeted temperature management (TTM) is recommended for comatose patients after out-of-hospital cardiac arrests (OHCAs). Even after successful TTM, several factors could influence the neuroprotective effect of TTM. The aim of this study is to identify prognostic factors associated with good neurological outcomes in TTM recipients.

This study used nationwide data during 2012 to 2016 to investigate prognostic factors associated with good neurological outcomes in patients who received TTM after the return of spontaneous circulation (ROSC). Multivariate logistic regression analysis was conducted to analyse the factors that may affect the neurological outcomes in the TTM recipients.

The study included 1578 eligible patients, comprising 767 with good and 811 with poor neurological outcomes. Multivariable analyses showed that OHCA in public places (OR, 1.599; 95% CI, 1.100–2.323, $P=.014$), initial shockable rhythms (OR, 1.721; 95% CI, 1.191–2.486, $P=.004$), pre-hospital ROSCs (OR, 6.748; 95% CI, 4.703–9.682, $P<.001$), bystander cardiopulmonary resuscitation (CPR) (OR, 1.715; 95% CI, 1.200–2.450, $P=.003$), and primary coronary interventions (PCIs) (OR, 2.488; 95% CI, 1.639–3.778, $P<.001$) were statistically significantly associated with good neurological outcomes. Whereas, increase of age (OR, 0.962; 95% CI, 0.950–0.974, $P<.001$) and conventional cooling (OR, 0.478; 95% CI, 0.255–0.895, $P=.021$) were statistically significantly associated with poor neurological outcome.

This study suggests that being younger, experiencing OHCA in public places, having initial shockable rhythm, pre-hospital ROSC, and bystander CPR, implementing PCIs and applying intravascular or surface cooling devices compared to conventional cooling method could predict good neurological outcomes in post-cardiac arrest patients who received TTM.

Abbreviations: CI = confidence interval, CPC = cerebral performance category, CPR = cardiopulmonary resuscitation, ECMO = extracorporeal membrane oxygenation, EMS = emergency medical service, KCDC = Korean Centres for Disease Control and Prevention, OHCA = out-of-hospital cardiac arrest, OHCAS = Out-of-Hospital Cardiac Arrest Surveillance, OR = odds ratio, PCI = percutaneous coronary intervention, ROSC = return of spontaneous circulation, TTM = targeted temperature management, VF = ventricular fibrillation, VT = ventricular tachycardia.

Keywords: Nationwide observation study, neurological outcomes, out-of-hospital cardiac arrest, prognosis, targeted temperature management

Editor: Abdelouahab Bellou.

JGK and HS contributed equally to this work.

The authors have no funding to disclose.

The authors have no conflicts of interest to disclose.

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

^a Department of Emergency Medicine, Hallym University College of Medicine, ^b Department of Emergency Medicine, Graduate School of Medicine, Kangwon National University, Chuncheon, ^c Department of Emergency Medicine, Hanyang University College of Medicine, Hanyang University Guri Hospital, Guri, ^d Department of Thoracic and Cardiovascular Surgery, Hallym University College of Medicine, Chuncheon, ^e Department of Internal Medicine, Hallym University College of Medicine, ^f Department of Internal Medicine, Hanyang University College of Medicine, Seoul, ^g Department of Emergency Medicine, Armed Force Yangju Hospital, Yangju, ^h Department of Emergency Medicine, Armed Force Capital Hospital, Seongnam, Republic of Korea.

* Correspondence: Hyun Young Choi, 1, Singil-ro, Yeongdeungpo-gu, Seoul 07441, Republic of Korea (e-mail: chy6049@naver.com).

Copyright © 2020 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

How to cite this article: Kim JG, Shin H, Choi HY, Kim W, Kim J, Moon S, Kim B, Ahn C, Lee J. Prognostic factors for neurological outcomes in Korean targeted temperature management recipients with return of spontaneous circulation after out-of-hospital cardiac arrests: A nationwide observational study. *Medicine* 2020;99:15 (e19581).

Received: 31 October 2019 / Received in final form: 16 February 2020 / Accepted: 18 February 2020

<http://dx.doi.org/10.1097/MD.00000000000019581>

1. Introduction

Severe neurological injury is the most common cause of death among patients who experience out-of-hospital cardiac arrests (OHCAs).^[1] A minority of OHCA patients are successfully resuscitated and survive with good neurological outcomes.

Targeted temperature management (TTM) is widely recommended during the critical care of patients who are comatose post-cardiac arrest.^[2–4] TTM as a neuroprotective treatment improves outcomes in post-resuscitation care,^[4,5] and it minimizes neuronal loss and brain damage.^[6] The potential mechanisms of TTM that improve neurologic outcomes are as follows: a lower cerebral metabolic rate for oxygen, suppression of the chemical reaction associated with reperfusion injury, or activation of anti-apoptotic mechanisms via cooling or control.^[7,8] Since several factors, including advanced life support, access to emergent coronary angiography, optimising support for cerebral and organ perfusion, and preventing hyperglycemia and infections,^[5,9–13] influence the neuroprotective effects of TTM, few patients achieve good neurological outcomes post-cardiac arrest, even after successful cardiopulmonary resuscitation (CPR) and TTM.

Many studies of several prognostic factors have evaluated their associations with neurological outcomes after the implementation of TTM.^[14,15] However, predicting neurological outcomes is difficult, even if TTM is successful. In this study, we analysed data from a nationwide cohort of patients who experienced OHCAs to identify prognostic factors associated with good neurological outcomes in TTM recipients.

2. Methods

2.1. Study design and settings

This was a retrospective observational study that evaluated the prognostic factors associated with good neurological outcomes in patients who underwent TTM from January 2012 to December 2016, using nationwide data from the Out-of-Hospital Cardiac Arrest Surveillance (OHCAS) database that is managed by the Korean Centers for Disease Control and Prevention (KCDC) (<https://cdc.go.kr>). OHCAS is undertaken in 17 provinces within South Korea, which has a population of 50 million people, and the database contains information describing the timing of the patients' cardiac arrests and the patients' outcomes at hospital discharge. The local ethics committee approved this study (Kangnam Sacred Heart Hospital's Institutional Review Board No. 2018-07-025-001), and the need for informed consent was waived because of the retrospective nature of the study and the use of anonymous clinical data for the analysis. The KCDC approved the use of the data for this study.

The application of and the types of TTM methods used were decided by physicians in accordance with the hospitals' protocols. This study's methodology fulfilled the criteria in the Strengthening the Reporting of Observational Studies in Epidemiology checklist.^[16]

2.2. Data source

The OHCAS database comprises a population-based, EMS-assessed OHCA registry and a retrospective patient cohort. The data from patients who had experienced OHCAs were extracted from the EMS registry; these data are collected by emergency medical technicians using EMS run sheets (patient records), and

they are input by EMS providers immediately after the transportation of OHCA patients. The data describing the OHCA patients' hospital care and outcomes at hospital discharge were provided by the KCDC's medical record reviewers who visit all of the emergency departments (EDs) and hospitals to which OHCA patients are referred, and they review the medical records in relation to the patients' treatment and outcomes.

The OHCAS database contains information that describes individuals and settings, EMSs, ED care, hospital procedures, and the outcomes at discharge, including mortality and the neurological outcomes, using a customised survey form. The registry form was developed based on the Utstein-style guidelines^[17] and the Resuscitation Outcome Consortium Project.^[18]

2.3. Study population

A total of 142,905 patients had experienced OHCAs and were registered in the OHCAS database between January 2012 and December 2016, and, of these, patients aged ≥ 18 years who experienced OHCAs and received TTM in hospital settings after ROSC as a result of CPR were included in the study. We excluded patients whose OHCAs were a consequence of trauma, patients aged < 18 years, any patients whose data describing their neurological status were invalid.

2.4. Variables

Information was collected that described the patients' ages and sexes, the places where the cardiac arrests occurred (public vs non-public), the aetiological factors (cardiac vs non-cardiac), the initially monitored rhythm (shockable vs non-shockable), witnessed cardiac arrests, bystander CPR, TTM-related factors, including the cooling method, pre-induction time, and induction time, PCIs, extracorporeal membrane oxygenation (ECMO), and mechanical CPR. Regarding the places where the cardiac arrests occurred, public places were defined as any places that are generally accessible to people, including roads, public buildings, parks, and beaches. An initial shockable rhythm was defined as initially monitored ventricular fibrillation (VF) or ventricular tachycardia (VT). ROSC was defined for all rhythms as the restoration of a spontaneous rhythm that was sustained for > 20 min. The cardiac origins of the OHCAs, including ischemic heart disease, arrhythmias, and cardiac tamponade, were identified from reviews of the medical records. A cardiac arrest was presumed to have had a non-cardiac origin if it was caused by trauma, drowning, poisoning, burns, asphyxia, hanging, or any other non-cardiac factor.

PCI included ballooning and primary stenting. Regarding initiation time of TTM, the pre-induction time was defined as the interval between the cardiac arrest and TTM initiation, and the induction time was defined as the interval between TTM initiation and the achievement of the target temperature.

The external temperature management systems used comprised external pads that were applied to the body surface to reduce the body temperature, and they included the Arctic Sun system (Medivance Inc, Louisville, CO), Gaymar products (Gaymar Industries Inc, Orchard Park, NY), the Blanketrol III system (Cincinnati Sub-Zero Products, Cincinnati, OH), and the Emcools Flex.Pad device (Emcools Medical Cooling Systems GmbH, Traiskirchen, Austria). The intravascular cooling devices used comprised intravascular catheters inserted into large vessels to reduce body temperature, and they included the Coolgard

3000 Thermal Regulation System (Aelsius Medical Corporation, Irvine, CA). Conventional cooling methods were defined as basic external cooling devices that included fans and ice packs that were placed on the main vascular access points, or basic internal cooling methods, including stomach irrigation with cool saline administered through a Levin tube, urinary bladder irrigation using cold saline administered through a Foley catheter, and cold saline intravenous infusions.

Target temperature and maintenance time for TTM were not provided from the raw data of OHCAs in this study. However, all TTM protocol in the South Korea has been adhering to the international 2010 to 2015 American Heart Association guidelines (target temperature: 32 to 36°C, maintenance time: 12 to 24 h).^[19,20]

Patients with invalid or missing data on neurological outcome were excluded from the study population.

2.5. Outcome measures

The study's outcome was the neurological outcome at hospital discharge, which was assessed using the cerebral performance category (CPC) scoring system. CPC scores of 1 and 2 were considered good neurological outcomes, and CPC scores of 3, 4, and 5 were considered poor neurological outcomes. The neurological outcomes were determined according to the discharge summaries in the patients' medical records.

2.6. Statistical analyses

The continuous data are presented as the medians and interquartile ranges, and the categorical data are presented as numbers and percentages. The Kolmogorov–Smirnov test was used to assess the normality of the data. The continuous variables were analysed using the independent sample *t* test for the parametric data and the Mann–Whitney *U* test for the nonparametric data. Pearson's chi-square test and Fisher's exact test were used to analyse the categorical variables. Multivariate logistic regression analysis, which used a backward likelihood ratio model, was conducted to analyse the factors that may affect the neurological outcomes in the TTM recipients. Any variables with *P* values <.05 in the univariate analyses were included in the multivariate regression analysis. All of the statistical analyses were conducted using IBMSPSS software, version 24.0 (IBM Corporation, Armonk, NY) and R, version 3.3.2 (The R Foundation for Statistical Computing, Vienna, Austria). A value of *P* <.05 was considered statistically significant.

3. Results

3.1. Characteristics of the study subjects

Of the 142,905 patients who had experienced OHCAs and were registered during the study period, we excluded patients who did not receive TTM (*n* = 140,301), patients aged <18 years or those who had experienced traumatic OHCAs (*n* = 612), and patients whose CPC scores were unknown at hospital discharge (*n* = 414). The remaining 1578 patients were enrolled to participate in this study, and, of these, 767 (48.6%) comprised the good neurological outcome group, and 811 (51.4%) comprised the poor neurological outcome group (Fig. 1).

Table 1 summarizes the clinical characteristics of the patients in the two groups. The median age of the patients in the good neurological outcome group was significantly lower than that of




Figure 1. Study's flow diagram. CPC=cerebral performance category, OHCA=out-of-hospital cardiac arrest, TTM=target temperature management.

the patients in the poor neurological outcome group (*P* <.001). The numbers of male patients (*P* = .003), OHCAs in public places (*P* <.001), pre-hospital ROSCs (*P* <.001), witnessed cardiac arrests (*P* <.001), patients who underwent bystander CPR (*P* <.001), initial shockable rhythms (*P* <.001), OHCAs of cardiac origin (*P* <.001), and percutaneous coronary interventions (PCI) (*P* <.001) were significantly higher in the good neurological outcome group than those in the poor neurological outcomes group. The rate of mechanical CPR was significantly lower in the good neurological outcome group than that in the poor neurological outcome group (*P* = .004). The induction time was significantly longer in the good neurological outcome than that in the poor neurological outcome group (*P* <.001), and the TTM cooling methods differed significantly between the groups (*P* <.001).

3.2. Analysis of prognostic factors associated with neurological outcomes

We performed multivariate logistic regression analyses to identify factors associated with good neurological outcomes. The variables included in the analyses were age, sex, OHCA in a public place, pre-hospital ROSC, witnessed cardiac arrest, bystander CPR, initial shockable rhythm, cause of cardiac arrest, mechanical CPR, TTM cooling method and initiation time, PCI, and ECMO. Among these variables, OHCA in a public places (odds ratio [OR]: 1.599; 95% confidence interval [CI]: 1.100–2.323; *P* = .014), initial shockable rhythm (OR: 1.721; 95% CI: 1.191–2.486; *P* = .004), pre-hospital ROSC (OR: 6.748; 95% CI: 4.703–9.682; *P* <.001), bystander CPR (OR: 1.715; 95% CI: 1.200–2.450; *P* = .003), and PCI (OR: 2.489; 95% CI: 1.639–3.778; *P* <.001) were statistically significantly associated with good neurological outcomes (Table 2). Whereas, older age (OR: 0.962; 95% CI: 0.950–0.974; *P* <.001) and conventional cooling




Figure 2. Independent predictors of good neurological outcomes in recipients of target temperature management following out-of-hospital cardiac arrest. CPR = cardiopulmonary resuscitation, OHCA = out of hospital cardiac arrest, PCI = primary coronary intervention, ROSC = return of spontaneous circulation.

who are less likely to undergo ROSCs and who experience longer periods of CPR before the ROSC, could worsen before TTM.

A pre-hospital ROSC was a prognostic factor for good neurological outcomes associated with TTM in our analysis. Chest compressions only generate 25% to 30% of the normal cardiac output when they are performed under optimal conditions. Hence, prolonged CPR is related to cerebral damage and a low likelihood of a ROSC.^[36] Consequently, a pre-hospital ROSC could contribute to good neurological outcomes in TTM recipients by reducing levels of brain damage before TTM.

Bystander CPR was identified as a prognostic factor for good neurological outcomes associated with TTM in this study. Cerebral blood flow is extremely low during closed-chest CPR following a prolonged cardiac arrest.^[37–39] Low cerebral blood flow is deleterious during cerebral resuscitation.^[40] The AHA and European guidelines stress the importance of early CPR as part of the chain of survival.^[41] Kragholm et al^[42] reported that compared with no bystander resuscitation, bystander CPR significantly reduced the brain damage and nursing home admission rates. If bystander CPR is implemented properly, it could yield better cerebral blood flows, and, therefore, contribute improved neurological outcomes after TTM.^[43]

In post-cardiac arrest management, PCI was identified as a prognostic factor for good neurological outcomes in the TTM recipients who had experienced OHCA in this study. The 2015 AHA guidelines recommend emergency coronary angiography for adult patients who are comatose after OHCA with suspected cardiac origins.^[3] Given that TTM could worsen cardiac dysfunction and hemodynamic imbalances and that coronary artery disease is a major cause of adult OHCA, PCI could contribute to good neurological outcomes in TTM recipients by removing the presumed cause of the OHCA and supporting the hemodynamic status during post-resuscitation care.^[21,35,44] Previous studies' findings have also shown that PCI in addition to TTM could positively influence the neurological outcomes in TTM recipients.^[45–48]

Regarding cooling methods, this study's findings showed that compared with intravascular cooling systems, the neurological outcomes were adversely associated with conventional cooling methods, and that there were no differences between intravascular and external cooling devices. Conventional cooling devices,

for example, crushed ice or ice bags, are the easiest ways to apply TTM,^[49] but they are associated with high rates of unintentional overcooling below the target temperature, which could have deleterious effects, including atrial or ventricular arrhythmias, coagulopathy, and increased risks of infection.^[50–53] Furthermore, compared with external or intravascular cooling devices, conventional cooling methods are less effective at maintaining temperatures,^[54] and these disadvantages could weaken the neuroprotective effect of TTM and adversely affect the neurological outcomes after TTM.^[55]

We found an adverse association between older age and the neurological outcomes after TTM. Oh et al^[56] reported that age was a negative prognostic factor for neurological outcomes (OR: 0.94; 95% CI: 0.93–0.96; $P < .001$). Winther-Jensen et al^[57] reported that older age was associated with increased rates of unfavourable neurological outcomes. As people age, comorbidity, and disability rates increase,^[58] and neurological function and neuronal plasticity decline^[58,59]; hence, older patients may respond poorly to TTM.

5. Limitation

Our study has several limitations. First, since this was a retrospective observational study, it may have been subject to selection bias relating to the type of TTM method used and the post-cardiac arrest management procedure implemented such as PCI. Therefore, result of this study should be interpreted with caution.

Second, we could not assess the patients' neurological outcomes beyond hospital discharge, because the neurological outcomes were measured using data from the patients' medical records at hospital discharge. Early CPC scores measured at hospital discharge can change until 6 months post-discharge^[60,61]; therefore, prognostic factors in this study could be different compared with prognostic factors for long-term neurological outcome of TTM recipients.

Third, our retrospective registry did not contain data that described potential confounders, for example, the patients' underlying diseases, hemodynamic statuses, mental statuses before TTM, the laboratory findings, and descriptions of the cardiac arrest durations. These factors could affect patient survival and the patients' neurological outcomes. Thus, more

studies that include more variables related to the patients' statuses and TTM are required to corroborate our results.

Fourth, target temperature and maintenance time for TTM were not provided from the raw data of OHCAS in this study. However, all TTM protocol in the South Korea has been adhering to the international 2010 to 2015 American Heart Association guidelines (target temperature: 32–36°C, maintenance time: 12–24 h).^[19,20] Therefore, the TTM guideline change from 2010 to 2015 might affect the outcomes of patients.

Finally, whether our study's findings can be generalised to other countries with different medical systems is uncertain. The study was performed in the context of South Korea's emergency medical service (EMS), which does not permit the provision of advanced cardiac life support to patients. Hence, the findings from this study should be interpreted with caution regarding their generalisability to other countries' medical systems that provide advanced cardiac life support to patients. To generate more generalizable data, studies involving other races and countries are required.

6. Conclusion

Good neurological outcomes could be expected among TTM recipients if patients are younger, experience OHCAs in public places, have initial shockable rhythms, and have pre-hospital ROSCs, and bystander CPR is performed. In post-cardiac arrest care, the implementation of PCIs, and intravascular or surface cooling rather than conventional cooling methods could improve patients' neurological outcomes after TTM.

Author contributions

Conceptualization: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi, Wonhee Kim, Jihoon Kim, Shinje Moon, Bongyoung Kim, Chiwon Ahn, Juncheol Lee.

Data curation: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi, Wonhee Kim, Jihoon Kim, Shinje Moon, Bongyoung Kim, Chiwon Ahn, Juncheol Lee.

Formal analysis: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi, Wonhee Kim.

Investigation: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi, Wonhee Kim, Jihoon Kim, Shinje Moon, Bongyoung Kim, Chiwon Ahn, Juncheol Lee.

Methodology: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi, Wonhee Kim, Jihoon Kim, Shinje Moon, Bongyoung Kim, Chiwon Ahn, Juncheol Lee.

Project administration: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi.

Resources: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi.

Software: Jae Guk Kim, Hyun Young Choi, Wonhee Kim.

Supervision: Hyun Young Choi.

Validation: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi, Jihoon Kim, Shinje Moon.

Visualization: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi.

Writing – original draft: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi.

Writing – review & editing: Jae Guk Kim, Hyungoo Shin, Hyun Young Choi, Wonhee Kim, Jihoon Kim, Shinje Moon, Bongyoung Kim, Chiwon Ahn, Juncheol Lee.

Hyun Young Choi orcid: 0000-0002-4898-3224.

References

- Laver S, Farrow C, Turner D, et al. Mode of death after admission to an intensive care unit following cardiac arrest. *Intensive Care Med* 2004;30:2126–8.
- Bernard SA, Gray T, Buist W, et al. Treatment of Comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med* 2002;346:557–63.
- Callaway CW, Donnino MW, Fink EL, et al. Part 8: post-cardiac arrest care: 2015 American Heart Association Guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2015;132(18 Suppl 2):S465–82.
- Group H, aCAS. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med* 2002;346:549–56.
- Nielsen N, Wetterslev J, Cronberg T, et al. Targeted temperature management at 33 degrees C versus 36 degrees C after cardiac arrest. *N Engl J Med* 2013;369:2197–206.
- Tripathy S, Mahapatra AK. Targeted temperature management in brain protection: an evidence-based review. *Indian J Anaesth* 2015;59:9–14.
- Han HS, Park J, Kim JH, et al. Molecular and cellular pathways as a target of therapeutic hypothermia: pharmacological aspect. *Curr Neuropharmacol* 2012;10:80–7.
- Steen PA, Newberg L, Milde JH, et al. Hypothermia and barbiturates: individual and combined effects on canine cerebral oxygen consumption. *Anesthesiology* 1983;58:527–32.
- Cronberg T, Lilja G, Horn J, et al. Neurologic function and health-related quality of life in patients following targeted temperature management at 33 degrees C vs 36 degrees C after out-of-hospital cardiac arrest: a randomized clinical trial. *JAMA Neurol* 2015;72:634–41.
- Fugate JE, Moore SA, Knopman DS, et al. Cognitive outcomes of patients undergoing therapeutic hypothermia after cardiac arrest. *Neurology* 2013;81:40–5.
- Kim F, Nichol G, Maynard C, et al. Effect of prehospital induction of mild hypothermia on survival and neurological status among adults with cardiac arrest: a randomized clinical trial. *JAMA* 2014;311:45–52.
- McNally B, Robb R, Mehta M, et al. Out-of-hospital cardiac arrest surveillance—Cardiac Arrest Registry to Enhance Survival (CARES), United States, October 1, 2005–December 31, 2010. *MMWR Surveill Summ* 2011;60:1–9.
- Smith K, Andrew E, Lijovic M, et al. Quality of life and functional outcomes 12 months after out-of-hospital cardiac arrest. *Circulation* 2015;131:174–81.
- Moseby-Knappe M, Pellis , Tommaso , et al. Head computed tomography for prognostication of poor outcome in comatose patients after cardiac arrest and targeted temperature management. *Resuscitation* 2017;119:89–94.
- Rundgren M, Karlsson , Torbjörn , et al. Neuron specific enolase and S-100B as predictors of outcome after cardiac arrest and induced hypothermia. *Resuscitation* 2009;80:784–9.
- von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 2008;61:344–9.
- Jacobs I, Nadkarni V, Bahr J, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update and simplification of the Utstein templates for resuscitation registries: a statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian Resuscitation Council, New Zealand Resuscitation Council, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Councils of Southern Africa). *Circulation* 2004;110:3385–97.
- Daya MR, Schmicker RH, Zive DM, et al. Out-of-hospital cardiac arrest survival improving over time: results from the Resuscitation Outcomes Consortium (ROC). *Resuscitation* 2015;91:108–15.
- Donnino MW, Andersen L, Berg KM, et al. Temperature management after cardiac arrest: an advisory statement by the Advanced Life Support Task Force of the International Liaison Committee on Resuscitation and the American Heart Association Emergency Cardiovascular Care Committee and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation. *Circulation* 2015;132:2448–56.

- [20] Neumar RW, Otto CW, Link MS, et al. Part 8: adult advanced cardiovascular life support: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010;122(18 Suppl 3):S729–67.
- [21] Nolan JP, Neumar RW, Adrie C, et al. Post-cardiac arrest syndrome: epidemiology, pathophysiology, treatment, and prognostication. A Scientific Statement from the International Liaison Committee on Resuscitation; the American Heart Association Emergency Cardiovascular Care Committee; the Council on Cardiovascular Surgery and Anesthesia; the Council on Cardiopulmonary, Perioperative, and Critical Care; the Council on Clinical Cardiology; the Council on Stroke. *Resuscitation* 2008;79:350–79.
- [22] Okada K, Ohde S, Otani NET-AL>. Prediction protocol for neurological outcome for survivors of out-of-hospital cardiac arrest treated with targeted temperature management. *Resuscitation* 2012;83:734–9.
- [23] Safar P. Cerebral resuscitation after cardiac arrest: research initiatives and future directions. *Ann Emerg Med* 1993;22(2 Pt 2):324–49.
- [24] Soleimanpour H, Rahmani F, Golzari SE, et al. Main complications of mild induced hypothermia after cardiac arrest: a review article. *J Cardiovasc Thorac Res* 2014;6:1–8.
- [25] Adrie C, Cariou A, Mourvillier B, et al. Predicting survival with good neurological recovery at hospital admission after successful resuscitation of out-of-hospital cardiac arrest: the OHCA score. *Eur Heart J* 2006;27:2840–5.
- [26] Sunde K, Pytte M, Jacobsen D, et al. Implementation of a standardised treatment protocol for post resuscitation care after out-of-hospital cardiac arrest. *Resuscitation* 2007;73:29–39.
- [27] Valenzuela TD, Roe DJ, Cretin S, et al. Estimating effectiveness of cardiac arrest interventions: a logistic regression survival model. *Circulation* 1997;96:3308–13.
- [28] Institute of Medicine. *Strategies to Improve Cardiac Arrest Survival: A Time to Act*. Washington, DC: The National Academies Press. 2015.
- [29] Murakami Y, Iwami T, Kitamura T, et al. Outcomes of out-of-hospital cardiac arrest by public location in the public-access defibrillation era. *J Am Heart Assoc* 2014;3:e000533.
- [30] Folke F, Gislason GH, Lippert FK, et al. Differences between out-of-hospital cardiac arrest in residential and public locations and implications for public-access defibrillation. *Circulation* 2010;122: 623–30.
- [31] Iwami T, Hiraide A, Nakanishi N, et al. Outcome and characteristics of out-of-hospital cardiac arrest according to location of arrest: a report from a large-scale, population-based study in Osaka, Japan. *Resuscitation* 2006;69:221–8.
- [32] Muraoka H, Ohishi Y, Hazui H, et al. Location of out-of-hospital cardiac arrests in Takatsuki City: where should automated external defibrillator be placed. *Circ J* 2006;70:827–31.
- [33] Holzer M, Bernard SA, Hachimi-Idrissi S, et al. Hypothermia for neuroprotection after cardiac arrest: systematic review and individual patient data meta-analysis. *Crit Care Med* 2005;33:414–8.
- [34] Sakoh M, Gjedde A. Neuroprotection in hypothermia linked to redistribution of oxygen in brain. *Am J Physiol Heart Circ Physiol* 2003;285:H17–25.
- [35] Stub D, Bernard , Stephen , et al. Post cardiac arrest syndrome. *Circulation* 2011;123:1428–35.
- [36] Matos RI, Watson RS, Nadkarni VM, et al. Duration of cardiopulmonary resuscitation and illness category impact survival and neurologic outcomes for in-hospital pediatric cardiac arrests. *Circulation* 2013;127:442–51.
- [37] Chandra N, Weisfeldt ML, Tsitlik J, et al. Augmentation of carotid flow during cardiopulmonary resuscitation by ventilation at high airway pressure simultaneous with chest compression. *Am J Cardiol* 1981;48:1053–63.
- [38] Ditchey RV, Winkler JV, Rhodes CA. Relative lack of coronary blood flow during closed-chest resuscitation in dogs. *Circulation* 1982;66: 297–302.
- [39] Hoekstra JW, Rinnert K, Van Ligten P, et al. The effectiveness of bystander CPR in an animal model. *Ann Emerg Med* 1990;19:881–6.
- [40] White BC, Aust SD, Arfors KE, et al. Brain injury by ischemic anoxia: hypothesis extension—a tale of two ions? *Ann Emerg Med* 1984;13(9 Pt 2):862–7.
- [41] Kleinman ME, Brennan EE, Goldberger ZD, et al. Part 5: adult basic life support and cardiopulmonary resuscitation quality: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2015;132(18 Suppl 2):S414–35.
- [42] Kragholm K, Wissenberg M, Mortensen RN, et al. Bystander efforts and 1-year outcomes in out-of-hospital cardiac arrest. *N Engl J Med* 2017;376:1737–47.
- [43] Hoekstra J. Bystander CPR: a review. *Resuscitation* 1990;20:97–113.
- [44] Nolan JP, Deakin CD, Soar J, et al. European Resuscitation Council guidelines for resuscitation 2005. Section 4. Adult advanced life support. *Resuscitation* 2005;67(Suppl 1):S39–86.
- [45] Hovdenes J, Laake JH, Aaberge L, et al. Therapeutic hypothermia after out-of-hospital cardiac arrest: experiences with patients treated with percutaneous coronary intervention and cardiogenic shock. *Acta Anaesthesiol Scand* 2007;51:137–42.
- [46] Knafelj R, Radsel P, Ploj T, et al. Primary percutaneous coronary intervention and mild induced hypothermia in comatose survivors of ventricular fibrillation with ST-elevation acute myocardial infarction. *Resuscitation* 2007;74:227–34.
- [47] Noc M. Urgent coronary angiography and percutaneous coronary intervention as a part of postresuscitation management. *Crit Care Med* 2008;36(11 Suppl):S454–7.
- [48] Wolfrum S, Pierau C, Radke PW, et al. Mild therapeutic hypothermia in patients after out-of-hospital cardiac arrest due to acute ST-segment elevation myocardial infarction undergoing immediate percutaneous coronary intervention. *Crit Care Med* 2008;36:1780–6.
- [49] Kim F, Olsufka M, Carlbom D, et al. Pilot study of rapid infusion of 2 L of 4 degrees C normal saline for induction of mild hypothermia in hospitalized, comatose survivors of out-of-hospital cardiac arrest. *Circulation* 2005;112:715–9.
- [50] Merchant RM, Abella BS, Peberdy MA, et al. Therapeutic hypothermia after cardiac arrest: unintentional overcooling is common using ice packs and conventional cooling blankets. *Crit Care Med* 2006;34(12 Suppl): S490–4.
- [51] Mouritzen CV, Andersen MN. Mechanisms of ventricular fibrillation during hypothermia. Relative changes in myocardial refractory period and conduction velocity. *J Thorac Cardiovasc Surg* 1966;51:585–9.
- [52] Rohrer MJ, Natale AM. Effect of hypothermia on the coagulation cascade. *Crit Care Med* 1992;20:1402–5.
- [53] Sessler DI. Complications and treatment of mild hypothermia. *Anesthesiology* 2001;95:531–43.
- [54] Hoedemakers CW, Ezzahiti M, Gerritsen A, et al. Comparison of cooling methods to induce and maintain normo- and hypothermia in intensive care unit patients: a prospective intervention study. *Crit Care* 2007;11:R91.
- [55] Feuchtl A, Gockel B, Lawrenz T, et al. Endovascular cooling improves neurological short-term outcome after prehospital cardiac arrest. *Intensivmedizin und Notfallmedizin* 2007;44:37–42.
- [56] Oh SH, Park KN, Lim J, et al. The impact of sex and age on neurological outcomes in out-of-hospital cardiac arrest patients with targeted temperature management. *Crit Care* 2017;21: 272.
- [57] Winther-Jensen M, Pellis T, Kuiper M, et al. Mortality and neurological outcome in the elderly after target temperature management for out-of-hospital cardiac arrest. *Resuscitation* 2015;91:92–8.
- [58] van de Glind EM, van Munster , Barbara C, et al. Pre-arrest predictors of survival after resuscitation from out-of-hospital cardiac arrest in the elderly a systematic review. *BMC Geriatr* 2013;13:68.
- [59] Wallmüller C, Spiel , Alexander , et al. Age-dependent effect of targeted temperature management on outcome after cardiac arrest. *Eur J Clin Invest* 2018;48:e13026.
- [60] Na MK, Kim W, Lim TH, et al. Gray matter to white matter ratio for predicting neurological outcomes in patients treated with target temperature management after cardiac arrest: a systematic review and meta-analysis. *Resuscitation* 2018;132:21–8.
- [61] Scheel M, Storm , Christian , et al. The prognostic value of gray-white-matter ratio in cardiac arrest patients treated with hypothermia. *Scand J Trauma Resusc Emerg Med* 2013;21:23.