

J Bone Metab 2016;23:49 http://dx.doi.org/10.11005/jbm.2016.23.1.49 pISSN 2287-6375 eISSN 2287-7029

## Erratum to: Vitamin D Deficiency Is Highly Concomitant but Not Strong Risk Factor for Mortality in Patients Aged 50 Year and Older with Hip Fracture

## Gyeong-Hak Lee<sup>1</sup>, Jung-Won Lim<sup>2</sup>, Yong-Gum Park<sup>3</sup>, Yong-Chan Ha<sup>2</sup>

<sup>1</sup>Department of Orthopaedic Surgery, National Medical Center, Seoul; <sup>2</sup>Department of Orthopaedic Surgery, Chung-Ang University College of Medicine, Seoul; <sup>3</sup>Department of Surgery, Chung-Ang University College of Medicine, Seoul, Korea

J Bone Metab 2015;22(4):205-209

In this article, on page 205, the author's name has been spelled incorrectly in this article: "Gyeong-Hak Lee" should have appeared as "Kyung-Hag Lee".

The author's name should be corrected as the following: from "Gyeong-Hak Lee" to "Kyung-Hag Lee"

In this article, on page 206, since the publication of this article, a small error has been discovered on Table 1.

The Table 1 should be corrected as the following:

## Copyright © 2016 The Korean Society for Bone and Mineral Research

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/3.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.



## Table 1. Demographics of patients

Tuble 1. Demographics of patients	
Parameters	Findings
Number of patients	489
Age at the time of admission (yr)	76.5 (50 to 101)
Male:Female	146:343
Body mass index (kg/m <sup>2</sup> )	22.0 (13.3 to 36.2)
Mean of 25-hydroxy-vitamin D <sub>3</sub> levels (ng/mL)	15.7 (0.0 to 112.8)
Diagnosis Neck fracture Intertrochanteric fracture	194 (39.7%) 295 (60.3%)
Type of management Internal fixation Arthroplasty Conservative	160 (32.7%) 312 (63.8%) 17 (3.5%)
ASA score 1 2 3 4	1 (0.2%) 137 (28.0%) 346 (70.8%) 5 (1.0%)
Osteoporosis Normal Osteopenia Osteoporosis	19 (3.9%) 111 (22.7%) 359 (73.4%)
Koval's grade by pre-fracture I III IV V VI VII	257 (52.6%) 90 (18.4%) 31 (6.3%) 42 (8.6%) 39 (8.0%) 22 (4.5%) 8 (1.6%)

ASA, American Society of Anesthesiologist.

We apologize for any inconvenience caused.