

Original Article

# Cardioprotection via mitochondrial transplantation supports fatty acid metabolism in ischemia-reperfusion injured rat heart

Jehee Jang<sup>1,#</sup>, Ki-Woon Kang<sup>2,#</sup>, Young-Won Kim<sup>1</sup>, Seohyun Jeong<sup>1</sup>, Jaeyoon Park<sup>3</sup>, Jihoon Park<sup>3</sup>, Jisung Moon<sup>3</sup>, Junghyun Jang<sup>3</sup>, Seohyeon Kim<sup>3</sup>, Sunghun Kim<sup>3</sup>, Sungjoo Cho<sup>3</sup>, Yurim Lee<sup>3</sup>, Hyoung Kyu Kim<sup>4</sup>, Jin Han<sup>4</sup>, Eun-A Ko<sup>5</sup>, Sung-Cherl Jung<sup>5</sup>, Jung-Ha Kim<sup>6,\*</sup>, and Jae-Hong Ko<sup>1,\*</sup>

<sup>1</sup>Department of Physiology, College of Medicine, Chung-Ang University, Seoul 06974, <sup>2</sup>Division of Cardiology, Department of Internal Medicine, College of Medicine, Chung-Ang University Hospital, Seoul 06973, <sup>3</sup>Department of Medicine, College of Medicine, Chung-Ang University, Seoul 06974, <sup>4</sup>Cardiovascular and Metabolic Disease Center, SMART Marine Therapeutics Center, Inje University, Busan 47392, <sup>5</sup>Department of Physiology, School of Medicine, Jeju National University, Jeju 63243, <sup>6</sup>Department of Family Medicine, College of Medicine, Chung-Ang University Hospital, Seoul 06973, Korea

## ARTICLE INFO

Received February 21, 2024

Revised February 27, 2024

Accepted February 28, 2024

## \*Correspondence

Jung-Ha Kim

E-mail: girlpower219@cau.ac.kr

Jae-Hong Ko

E-mail: akdongyi01@cau.ac.kr

## Key Words

Autografts

Mitochondria

Myocardial ischemia

Myocardial reperfusion

Oxygen consumption

Transplantation

#These authors contributed equally to this work.

**ABSTRACT** In addition to cellular damage, ischemia-reperfusion (IR) injury induces substantial damage to the mitochondria and endoplasmic reticulum. In this study, we sought to determine whether impaired mitochondrial function owing to IR could be restored by transplanting mitochondria into the heart under *ex vivo* IR states. Additionally, we aimed to provide preliminary results to inform therapeutic options for ischemic heart disease (IHD). Healthy mitochondria isolated from autologous gluteus maximus muscle were transplanted into the hearts of Sprague–Dawley rats damaged by IR using the Langendorff system, and the heart rate and oxygen consumption capacity of the mitochondria were measured to confirm whether heart function was restored. In addition, relative expression levels were measured to identify the genes related to IR injury. Mitochondrial oxygen consumption capacity was found to be lower in the IR group than in the group that underwent mitochondrial transplantation after IR injury ( $p < 0.05$ ), and the control group showed a tendency toward increased oxygen consumption capacity compared with the IR group. Among the genes related to fatty acid metabolism, *Cpt1b* ( $p < 0.05$ ) and *Fads1* ( $p < 0.01$ ) showed significant expression in the following order: IR group, IR + transplantation group, and control group. These results suggest that mitochondrial transplantation protects the heart from IR damage and may be feasible as a therapeutic option for IHD.

## INTRODUCTION

Ischemic heart disease (IHD) has been recognized as a primary cause of death in patients with cardiovascular disorders; its pathophysiology is the inability of the left ventricular (LV) myocardium to pump the required blood flow to meet metabolic demands [1].

Despite both pharmacological and interventional advances, the current therapeutic strategy for IHD provides limited prevention of heart failure after the occurrence of IHD. This is especially true for patients who have been treated with reperfusion within a short period following IHD and have experienced extensive LV myocardial damage subsequent to advanced heart failure and even



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. Copyright © Korean J Physiol Pharmacol, pISSN 1226-4512, eISSN 2093-3827

**Author contributions:** All authors read and approved the final manuscript. J.J., K.W.K., and Y.W.K.: Conceptualization, Data curation, Formal analysis, Writing – original draft, Writing – review & editing. J.J. and K.W.K.: Visualization. J.J., S.J., J.Y.P., J.H.P., J.M., J.J.<sup>3</sup>, S.Hy.K., S.Hu.K., S.J.C., and Y.L.: Investigation, Writing – review & editing. K.W.K. and Y.W.K.: Resources, Project administration. H.K.K., J.H., E.A.K., and S.C.J.: Project administration, Formal analysis, Writing – review & editing. J.H.K.<sup>6</sup> and J.H.K.: Conceptualization, Funding acquisition, Supervision, Writing – original draft, Writing – review & editing.

cardiovascular death [2]. More effective therapeutic options have long been required to protect and improve damaged LV myocardial function after the occurrence of IHD. Furthermore, recent research has shown that when oxygen and substrates required for respiration are lacking due to ischemia, the mitochondrial tricarboxylic acid cycle and oxidative phosphorylation are interrupted and the mitochondrial membrane voltage collapses, causing oxidative damage and apoptosis [3-5]. Ischemia-reperfusion (IR) leads to  $\text{Ca}^{2+}$  overload and the formation of reactive oxygen species, which open the mitochondrial permeability transition pore, release cytochrome c into the cytoplasm, and activate caspase-3 [6].

Normal mitochondrial function supplies the energy necessary for cell survival through the production of adenosine triphosphate (ATP), which is abundant in the liver, brain, heart, kidney, and muscle cells [7,8]. In particular, mitochondria occupy more than 30% of the volume of cardiomyocytes, provide 95% of the ATP required for heartbeat, and serve as metabolic hubs for oxidative phosphorylation, the citric acid cycle, and fatty acid  $\beta$ -oxidation [9]. Cyclosporine, MTP-131, and TRO4030 are used in the pharmacological treatment of mitochondrial damage, inhibiting mitochondrial membrane permeability and preventing cell death. However, the precise mechanism underlying this phenomenon is not yet known [6]. Another option may be mitochondrial transplantation, which involves the transfer of mitochondria into LV myocardial cells. Mitochondrial transplantation is a method of separating mitochondria from externally derived tissues or cells and transplanting them into new tissues or cells [10]. When mitochondria enter IR-damaged myocardial cells through transplantation, energy production is increased by replenishing mitochondrial DNA (mtDNA), and the cells are restored to their proper function [10,11]. To date, methods for transplanting mitochondria *in vitro* have included co-incubation, direct microinjection, cell-penetrating peptides, mitoception, photothermal nanoblade use, magnetomitotransfer, Mitopunch, Fluid FM, and mitochondrial delivery of the extracellular vesicles, among others [10]. However, these methods have limitations in reflecting *in vivo* situations because they represent phenomena that occur in a test tube. In addition, although research on direct mitochondrial transfer was conducted before cell-mediated mitochondrial transfer, its mechanism and strategy remain controversial. Therefore, we aimed to provide preliminary results for clinical therapeutic applications by directly transplanting mitochondria into LV myocardial cells using the Langendorff system.

The Langendorff system is an experimental method developed by Oscar Langendorff in 1895 that allows the heart to beat *in vitro* by performing retrograde perfusion of an isolated heart [12]. When an insertion tube is connected to the aorta and a solution containing oxygen and nutrients is perfused, the solution is perfused into the coronary artery by the aortic valve, supplying oxygen and nutrients to the myocardium, and the perfusate is discharged into the right atrium. By obstructing the oxygen sup-

ply to the myocardium, conditions such as ischemia, myocardial infarction, and reperfusion injury can be reproduced and studied [13].

In the present study, mitochondrial transplantation was performed using the Langendorff system after creating a situation similar to that of IR *in vitro*. We aimed to investigate the therapeutic feasibility of mitochondrial transplantation as a treatment for IHD *via* the following experiments. First, the rat heart was connected to the Langendorff system to create IR and post-IR mitochondrial transplantation states. Second, after the Langendorff perfusion experiment, the oxygen consumption capacity of the mitochondria in the myocardial tissue was measured, and the respiratory control rate (RCR) was calculated to confirm the ATP production capacity of the mitochondria. Third, the number of mitochondrial copies was measured using real-time PCR. The ATP production capacity per mitochondrial unit was confirmed by the second and third experiments. Fourth, to identify candidate genes related to IHD, gene expression levels were measured using real-time PCR.

## METHODS

### Laboratory animals

Sprague-Dawley rats (male, 7 weeks old, body weight  $200 \pm 10$  g) (Coatech) were used for all experiments. Animal testing was approved by the Chung-Ang University Animal Experiment Ethics Committee (approval number: 202301020113). The breeding environment was maintained at a temperature of  $22^\circ\text{C} \pm 3^\circ\text{C}$ , humidity of  $50\% \pm 10\%$ , ventilation of 14 to 18 times/h, lighting of 150 to 300 lx, and light/dark cycle of 12 h. Water and food were freely consumed during the experimental period. Twelve animals were randomly assigned to three groups: the normal group (CON, control group), ischemia-reperfusion group (IR), and ischemia-reperfusion-mitochondrial transplantation group (IR + transpl). Three animals in each group were utilized for assessing mitochondrial function, and the fourth was used for Evans staining. All animal experiments were performed in accordance with Institutional Animal Care and Use Committee (IACUC) guidelines, the Laboratory Animal Act, and animal testing regulations.

### Heart isolation

The animals were anesthetized by intravenous injection (1 ml/kg) of alfaxalone (5 mg/kg, Alfaxan; JUROX Pty Limited), xylazine hydrochloride (5 mg/kg, Rompun inj.; BAYER KOREA Ltd.), and heparin (300 IU/kg, Greencross Heparin Sodium Inj.; GC Biopharma Corporation) [14,15]. After confirming the absence of the flexion reflex, the rats were euthanized *via* heart excision. The aorta of each isolated rat heart and the cannula were then securely ligated with sutures. Subsequently, each heart was installed onto

the Langendorff system and retrogradely perfused with Normal Tyrode's (NT) solution. This solution contained 143 mM NaCl, 5.4 mM KCl, 5 mM HEPES, 0.5 mM MgCl<sub>2</sub>, 0.3 mM NaH<sub>2</sub>PO<sub>4</sub>, 1.8 mM CaCl<sub>2</sub>, and 5.5 mM glucose, and was adjusted to pH 7.4.

### Animal cardiac ischemic model

The animal cardiac ischemic model was modified as previously described [15]. The Langendorff systems were washed using flowing sterilized water and NT solution for 5 min each, and the experiment was performed after preheating the Langendorff systems to 37°C using a constant temperature circulating water tank (Changshin Science). All of the isolated hearts were stabilized via perfusion with oxygen-saturated NT solution for 20 min. The control group was perfused with an oxygen-saturated NT solution for 150 min, and the IR group was perfused with a nitrogen-saturated NT solution for 30 min to induce ischemia and then perfused with an oxygen-saturated NT solution for 60 min to induce reperfusion injury. In the IR + transpl group, IR damage was induced in the same manner as in the IR group, and the isolated mitochondria were perfused for 60 min, followed by transplantation (see Fig. 1A). The heart was perfused retrogradely in constant flow mode, and the flow rate was adjusted to  $7 \pm 1$  ml/min. The heart rate (HR) was measured via video observation lasting for 10 sec.

### Isolation and transplantation of mitochondria

After euthanasia, approximately 1 g of the gluteus maximus muscle was collected, transferred to a 50 ml tube, and mixed with 5–10 ml mitochondrial isolation buffer (MIB) 1 solution (180 mM KCl, 0.5 mM EDTA-Na<sub>2</sub>, 10 mM Tris-base, pH 7.4 at 4°C). The muscle was then cut into small pieces with medical scissors and


then homogenized with an overhead stirrer. The homogenate was centrifuged at 1,000 g for 10 min at 4°C, the supernatant was transferred to a new tube, and centrifugation (1,000 g, 10 min, 4°C) was repeated. Subsequently, the supernatant was decanted to a new tube and centrifuged (7,000 g, 10 min, 4°C), and the resulting supernatant was removed. The pellet was resuspended in 5–10 ml of MIB 2 solution (180 mM KCl, 0.5 mM EDTA-Na<sub>2</sub>, 10 mM Tris-base, 1 g/l BSA, pH 7.4 at 4°C) and centrifuged (7,000 g, 10 min, 4°C). The resulting supernatant was removed, resuspended with 5–10 ml of MIB 2 solution, centrifuged (7,000 g, 10 min, 4°C) once more, and the supernatant was removed to obtain a mitochondrial pellet. For quality control, the mitochondria content was validated through RCR, and quantification was based on protein levels using the Bradford assay. The transplantation solution was prepared with 230 ng of mitochondrial protein per 200 ml of NT solution and perfused for 60 min [16].

### Confirmation of the non-infarcted area

After completing the experimental protocol, 0.6 ml of 0.25% Evans blue dye was uniformly infused into the aortic cannula orifice. Subsequently, the heart was frozen at –20°C for 2 h and sliced into 5 mm thick transverse sections. The presence of a blue stain demarcated the non-ischemic area; conversely, a pale negative stain indicated the infarcted myocardium and/or the viable myocardium in the area at risk [17,18].

### Preparation of permeabilized fibers from heart tissue

The details of our experiment have been previously described by Kuznetsov *et al.* [19]. The LV endocardium of the perfused heart was immersed in isolation solution A (10 mM Ca-EGTA solution, 0.1 μM Ca, 20 mM imidazole, 20 mM taurine, 49 mM



**Fig. 1. Observation of non-ischemic area at endpoint of animal experiment.** (A) Schematic illustration of experimental protocol. (B) Representative cardiac sections stained with 0.25% Evans blue. The upper panel shows control group cardiac section stained with 0.25% Evans blue. The middle panel shows IR group cardiac section stained with 0.25% Evans blue. The lower shows IR + transpl group cardiac section stained with 0.25% Evans blue. CON (control), group without ischemia or reperfusion injury; IR, group with damage induced by ischemia and reperfusion; IR + transpl, group transplanted with isolated mitochondria after inducing IR damage.

K-MES, 3 mM K<sub>2</sub>HPO<sub>4</sub>, 9.5 mM MgCl<sub>2</sub>, 5.7 mM ATP, 15 mM phosphocreatine, 1 μM leupeptin, pH 7.1) containing 10 mM Ca-EGTA solution (2.77 mM CaK<sub>2</sub>EGTA stock solution, 7.23 mM K<sub>2</sub>EGTA stock solution) (CaK<sub>2</sub>EGTA stock solution: 100 mM CaCO<sub>3</sub>, 100 mM EGTA, 200 mM KOH, K<sub>2</sub>EGTA stock solution: 100 mM EGTA, 200 mM KOH). Then, 50 μg/ml saponin was added to 2 ml of the stripped tissue and vortexed (minimum speed, 20 min, 4°C) to create permeable tissue. Afterwards, to remove saponin and ATP, respiration medium B (0.5 mM EGTA, 3 mM MgCl<sub>2</sub>·6H<sub>2</sub>O, 20 mM taurine, 10 mM KH<sub>2</sub>PO<sub>4</sub>, 20 mM HEPES, 1 g/l BSA, 60 mM K-lactobionate, 110 mM mannitol, 0.3 mM DTT, pH 7.1) (60 mM K-lactobionate: 500 mM K-lactobionate stock solution) and 1 mg/ml fatty acid-free BSA was added to the tissue and vortexed (minimum speed, 5 min, 4°C). The procedure was repeated three to four times, followed by an experiment measuring the oxygen consumption capacity of the mitochondria.

### Measurement of mitochondrial oxygen consumption capacity

Tissues and solutions were placed inside a chamber (Instech) connected to a Neofox system (Ocean Optics) to measure oxygen consumption capacity. The oxygen consumption rate (OCR) was measured using the NeoFox viewer (version 2.30; Ocean Optics) program. After the chamber was preheated to 37°C using a constant temperature circulating water tank (Changshin Science), 100 μl of respiration medium B solution was added into the chamber, and the magnetic stir bar was operated for 10–15 min until the viewer's graph was stabilized. Subsequently, permeable tissue, G/M (10 mM glutamate + 5 mM malate), and 2 mM adenosine diphosphate were added (in that order), and oxygen consumption

capacity was measured by observing the change in oxygen inside the chamber. After the slopes of states 3 and 4 had been calculated, the RCR was calculated as follows [20–22] (Supplementary Fig. 1):

$$\text{RCR} = \text{state 3} / \text{state 4}$$

### Mitochondrial copy number (mtDNA-CN) measurement

The mtDNA-CN was measured using endo-myocardial DNA. DNA was obtained using a QIAamp DNA Mini Kit (Qiagen), and 8 ng was used for quantification using a UV spectrophotometer (Nanodrop 1000; Thermo Fisher Scientific). Real-time PCR was performed using a LightCycler 2.0 instrument (Roche Diagnostics). After the cycle of initial denaturation at 95°C for 10 min, denaturation at 95°C for 10 sec, annealing at 60°C for 10 sec, and extension at 72°C for 10 sec, the melting curve was performed 35 times under conditions of 65°C–95°C and 0.1°C/sec. The primers used in this study are listed in Table 1 [23,24]. Because the extracted DNA included genomic DNA (gDNA) and mtDNA, the expression level was quantified by comparing the mtDNA-CN and nuclear gene copy number. The formulas used to quantify the expression level and the RCR per unit of mitochondria were as follows [25]:

$$\Delta\text{Ct} = \text{mtDNA Ct} - \text{gDNA Ct}$$

$$\text{Relative DNA copy number} = 2 \times 2^{-\Delta\text{Ct}}$$

$$\text{OCR per unit mitochondria} = \text{RCR} / \text{mitochondria copy number}$$

**Table 1. Primers used for amplification**

Primer	Sequence 5'–3'	Annealing temp (°C)	Cycles	Ref.	
MT copy number					
MT	Forward	GCCACAAGTACACATCCACA	58	35	[23]
	Reverse	GGGGTAATGAAAGAGGCAA			
<i>Gapdh</i>	Forward	TGCTTACCACCTTCTTGAT	60	35	[23]
	Reverse	TGGAAAGCTGTGGCGTGAT			
Real-time PCR					
<i>Gapdh</i>	Forward	TGCTTACCACCTTCTTGAT	60	35	[23]
	Reverse	TGGAAAGCTGTGGCGTGAT			
<i>Cpt1b</i>	Forward	CAGCCATGCCACCAAGATC	61	40	[24]
	Reverse	AAGGGCCGCACAGAATCC			
<i>Elovl5</i>	Forward	AGAATAGTGGCATGGTGGTAGACG	61	40	Self design
	Reverse	ACGAGCGCAGGAGAATCAGA			
<i>Fads1</i>	Forward	ATGCCCTCTTCTAAATCTCGTTCC	61	40	Self design
	Reverse	TGGTAGGGGCAAATGGTGGTA			
<i>Gja1</i>	Forward	GCTTGTGTAATTGCCGCACGAGGAA	61	40	Self design
	Reverse	AAGGATCGCGTGAAGGAAGAAGC			

MT, mitochondrial genome; *Gapdh*, glyceraldehyde-3-phosphate dehydrogenase gene; *Cpt1b*, carnitine palmitoyltransferase 1B; *Elovl5*, ELOVL fatty acid elongase 5; *Fads1*, fatty acid desaturase 1; *Gja1*, gap junction alpha 1.

## Real-time quantitative PCR

Real-time PCR was performed to measure gene expression using a LightCycler 2.0 (Roche Diagnostics). After initial denaturation at 95°C for 10 min followed by 40 cycles of denaturation at 95°C for 10 sec, annealing for 10 sec at an appropriate temperature for each primer, and extension for 10 sec at 72°C, the melting curve was measured at 65°C–95°C and 0.1°C/sec. The primers and denaturation temperatures are listed in Table 1 [23,24]. The expression level of the candidate gene was normalized to the expression level of *Gapdh*, a reference gene, and the relative quantity was quantified using the  $2^{-\Delta\Delta Ct}$  method [26].

$$\Delta Ct = \text{target gene Ct} - \text{reference gene Ct}$$

$$\Delta\Delta Ct = \text{target sample group } \Delta Ct - \text{control group } \Delta Ct$$

$$\text{Relative expression} = 2^{-\Delta\Delta Ct}$$

## Statistical analysis

Values were expressed as mean  $\pm$  standard error of the mean. The Kruskal–Wallis test was selected because RCR does not require the groups to be normally distributed. Analysis of variance (ANOVA) was utilized to compare mtDNA-CN and gene expression levels. We employed repeated-measures ANOVA to analyze the time-dependent effects of HR. The Bonferroni test was used to adjust for multiple comparisons among the three groups in the *post-hoc* analysis. Statistical analyses were conducted using R version 4.3.2 [27], with a significance threshold set at 0.05.

## RESULTS

### Acute IR injury influenced HR in perfused rat heart

After stabilization, the isolated hearts exhibited a normal beating rate of  $145 \pm 4$  beats/min. Following mitochondrial transplantation, the IR + transpl group displayed a tendency toward a de-


creased HR compared to the other groups. However, we observed inconsistencies in this result, as all isolated hearts maintained a HR within the normal range [28] seen *ex vivo* from stabilization to the termination of the experiment, and no statistically significant differences were detected (Fig. 1).

### Mitochondrial transplantation did not induce the structural recovery of IR-damaged hearts


We observed areas of the heart that were stained blue (indicating perfused regions), as well as those that did not stain (indicating injured regions). The results revealed that the non-ischemic area in the IR group was distinctly smaller than that in the control group; this was not restored in the IR + transpl group (Fig. 2).

### Mitochondrial transplantation revealed a tendency to restore mitochondrial function in IR-damaged hearts

To evaluate mitochondrial function, the oxygen consumption



**Fig. 2. Changes in heart rate over time by group.** All isolated hearts showed normal *ex vivo* heart rates during the experiment. CON (control), group without ischemia or reperfusion injury; IR, group with damage induced by ischemia and reperfusion; IR + transpl, group transplanted with isolated mitochondria after inducing IR damage.



**Fig. 3. Mitochondrial function analysis.** (A) RCR tracked ATP production by assessing the oxygen consumption capacity of mitochondria. \* $p < 0.05$  indicates that the group means are not all statistically equal. # $p < 0.05$  denoted a comparison between IR and IR + transpl. (B) mtDNA-CN assessed cellular ATP requirements. (C) Oxygen consumption per unit mitochondria was confirmed. CON (control), group without ischemia or reperfusion injury; IR, injury due to ischemia and reperfusion group; IR + transpl, group transplanted with isolated mitochondria after inducing IR injury; mtDNA-CN, mitochondrial DNA copy number; RCR, respiratory control ratio; ATP, adenosine triphosphate.



capacity was measured twice for each individual, and the RCR was calculated. Compared with the control group, the IR group showed a tendency toward decreased oxygen consumption capacity (1.68 times lower). Conversely, compared with the IR group, the IR + transpl group showed a tendency toward increased oxygen consumption capacity (2.04 times higher), although this difference did not reach statistical significance (Fig. 3A). mtDNA-CN was observed to be similar in all groups (Fig. 3B). Notably, IR injury led to a decreased RCR to mtDNA-CN ratio, then recovered after mitochondrial transplantation (Fig. 3C).


### Mitochondrial transplantation regulated several fatty acid metabolism-related genes

A list of genes linked to fatty energy metabolism in rat skeletal and myocardial tissues, derived from our prior studies [15,23,29], was integrated with the relevant KEGG dataset [rno01212: Fatty acid metabolism - *Rattus norvegicus* (rat)] [30]. In total, 62 genes were identified, from which we randomly selected four. The relative expression levels of *Cpt1b* ( $p < 0.05$ ), *Gja1* ( $p < 0.01$ ), and *Fads1* ( $p < 0.01$ ) differed between the groups, although that of *Elovl5* did not. The expression level of *Fads1* was higher in the IR group than in the control group ( $p < 0.05$ ), and was lower in the IR + transpl group than in the IR group ( $p < 0.01$ ). The expression level of *Gja1* was lower in the IR + transpl group than in the control ( $p < 0.01$ ) or IR groups ( $p < 0.05$ ) (Fig. 4).

## DISCUSSION

There is a dearth of effective therapeutic options for IHD due to rising medical costs and cardiovascular-related mortality rates. Thus, in this study, we sought to determine whether damaged myocardial function could be restored by transplanting mitochondria into the heart under *ex vivo* IR states. It was previously reported that transplanted mitochondria can remain viable in the myocardium for 28 days although isolated mitochondrial function rapidly decreases within 1 h owing to external factors [31]. Isolated mitochondria from L6 cells were co-incubated with H9C2 cells to further determine the time dependence of mitochondrial transplantation efficiency. As no significant differences were observed within 3 h (data not shown), mitochondrial transplantation was performed for more than an hour in the Langendorff experiment.

The isolated hearts were connected to the Langendorff system, and an IR-damaged state and post-damage mitochondrial transplantation conditions were reproduced. As a result, the HR tended to slow during ischemic conditions compared to that under normoxic conditions, but tended to increase again during reperfusion and mitochondrial transplantation (Fig. 2). It has been reported that under ischemic conditions, the HR increases as a compensation for the lack of oxygen and nutrients being delivered to the myocardium [21,32,33]; however, in this experiment, this was predicted to be the result of the heart failing to interact with



**Fig. 4. Relative expression levels of target genes.** Comparison of cardiac gene expression levels associated with fatty acid energy metabolism in the mitochondrial transplantation. Statistically significant results indicated that the group means are not equal, with significance levels of \* $p < 0.05$  and \*\* $p < 0.01$ . The gene expression showed statistically significant differences, with \* $p < 0.05$  and \*\* $p < 0.01$  indicating levels of significance. CON (control), group without ischemia or reperfusion injury; IR, group with damage induced by ischemia and reperfusion; IR + transpl, group transplanted with isolated mitochondria after inducing IR damage; *Cpt1b*, carnitine palmitoyltransferase 1B; *Elovl5*, ELOVL fatty acid elongase 5; *Fads1*, fatty acid desaturase 1; *Gja1*, gap function alpha 1.

surrounding tissues under *ex vivo* conditions. The simple HR was a poor biomarker in our study because it remained within the normal biological range.

While our findings showed that oxygen consumption capacity was lower in the IR group than in the control group, the oxygen consumption capacity of the IR + transpl group was higher than that of the IR group, although it improved compared to that of the IR group, it did not improve sufficiently to match that of the control group (Fig. 1). Cowan *et al.* [32] reported that mitochondrial transplantation into cardiac cells, such as myocardial cells and fibroblasts, was possible by culturing cardiac cells under oxygen-deprived conditions to create conditions similar to those of ischemia, followed by co-incubating with a solution containing mitochondria. This has been reproduced using the Langendorff system [32,33]. Research has shown that most transplanted mitochondria fuse with existing mitochondria to form a network, and some are hydrolyzed [31]. In other words, IR damage can be induced using the Langendorff system, and the transplantation method of directly perfusing mitochondria into the heart using the Langendorff system appears to protect myocardial tissue from IR damage. Therefore, mitochondrial transplantation may emerge as a therapeutic option for IHD. Currently, the best-known mechanism by which mitochondria move between cells involves endosomes [11,31,34]. Specifically, mitochondria are transplanted into cardiac cells through actin-dependent endocytosis, which increases energy production and replenishes mtDNA to induce proper cell function [11,31,34]. Under normal conditions, the heart uses fatty acids as the main energy substrate, followed by carbohydrates, ketone bodies, and amino acids; however, it is known to dynamically switch substrate availability to ensure efficient cardiac function [35]. Among the three groups in this study, the *Cpt1b* and *Fads1* genes showed the highest expression in the IR group and the lowest expression in the IR + transpl group (Fig. 4). A previous study showed a reduction in fatty acid oxidation with lipid accumulation in *Cpt1b* knockout mice, demonstrating that this gene is related to fatty acid oxidation [36]. From this, it can be expected that normal fatty acid oxidation did not occur due to IR damage, and that the expression of *Cpt1b* increased as a compensation. In the IR + transpl group, the expression of *Cpt1b* was lower, suggesting the restoration of fatty acid metabolism and myocardial protection effects through mitochondrial transplantation. A previous study demonstrated that the *Fads1* gene induces apoptosis and increases reactive oxygen species when silenced, indicating that it is a gene related to the survival of healthy cells [37]. In the IR group, it could be expected that the expression of the *Fads1* gene increased due to the excessive production of reactive oxygen species in the myocardium and a protective effect against cell damage. In the IR + transpl group, the expression of the *Fads1* gene was reduced but was restored to normal by mitochondrial transplantation, and apoptosis-prevention effects were expected. The expression of connexin 43 (encoded by *Gja1*) increases under ischemic conditions [38]. Existing research has

shown that this gene protects the mitochondria by reducing mitochondrial membrane potential, respiration, and reactive oxygen species production [38]. In this study, the expression of *Gja1* was found to be lower in the IR + transpl group than in the IR group, which is thought to be due to the protective effect of mitochondrial transplantation rather than *Gja1* itself, resulting in relatively lower expression of *Gja1* (Fig. 4). Considering the mitochondrial respiratory capacity in myocardial tissue observed in the above results, it appears that cardiomyocyte dysfunction can be restored by replenishing mitochondria damaged by IHD through mitochondrial transplantation; however, it also functions as normal myocardial cells without IR damage and thus, can be interpreted as not recovering. In the future, based on the results of this study, additional interpretation of the functions of the *Cpt1b* and *Fads1* genes, which appear to be expressed as compensatory effects of IR, will be required. In addition, various transplantation methods will be developed by applying mitochondrial transplantation to an *in vivo* IR model and methods that can restore function to the levels observed in the normal myocardium will be studied.

Given that acute injury often manifests through metabolic alterations rather than immediate physical or structural changes, our study focused on assessing infarct size using Evans blue staining and detecting IHD-related gene markers *via* PCR. While additional investigations are warranted to address the limitations in the number of repetitions and to elucidate the underlying mechanisms and potential clinical applications, the findings of this study suggest that mitochondrial transplantation may confer protection against early IR injury through the modulation of cardiac energy metabolism. Therefore, mitochondrial transplantation holds promise as a potential therapeutic intervention for IHD.

## FUNDING

This study was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF), funded by the Ministry of Education (2017R1D1A1B06035273 and 2021R1A6A3A01088465), and by the Chung-Ang University Research Scholarship Grants in 2022.

## ACKNOWLEDGEMENTS

None.

## CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

## SUPPLEMENTARY MATERIALS

Supplementary data including one figure can be found with this article online at <https://doi.org/10.4196/kjpp.2024.28.3.209>

## REFERENCES

- Vedin O, Lam CSP, Koh AS, Benson L, Teng THK, Tay WT, Braun OÖ, Savarese G, Dahlström U, Lund LH. Significance of ischemic heart disease in patients with heart failure and preserved, midrange, and reduced ejection fraction: a nationwide cohort study. *Circ Heart Fail*. 2017;10:e003875.
- Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, de Ferranti SD, Floyd J, Fornage M, Gillespie C, Isasi CR, Jiménez MC, Jordan LC, Judd SE, Lackland D, Lichtman JH, Lisabeth L, Liu S, Longenecker CT, Mackey RH, et al; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics-2017 update: a report from the American Heart Association. *Circulation*. 2017;135:e146-e603. Erratum in: *Circulation*. 2017;135:e646. Erratum in: *Circulation*. 2017;136:e196.
- Fuhrmann DC, Brüne B. Mitochondrial composition and function under the control of hypoxia. *Redox Biol*. 2017;12:208-215.
- Leist M, Single B, Castoldi AF, Kühnle S, Nicotera P. Intracellular adenosine triphosphate (ATP) concentration: a switch in the decision between apoptosis and necrosis. *J Exp Med*. 1997;185:1481-1486.
- Lesnefsky EJ, Chen Q, Slabe TJ, Stoll MS, Minkler PE, Hassan MO, Tandler B, Hoppel CL. Ischemia, rather than reperfusion, inhibits respiration through cytochrome oxidase in the isolated, perfused rabbit heart: role of cardiolipin. *Am J Physiol Heart Circ Physiol*. 2004;287:H258-H267.
- Trankle C, Thurber CJ, Toldo S, Abbate A. Mitochondrial membrane permeability inhibitors in acute myocardial infarction: still awaiting translation. *JACC Basic Transl Sci*. 2016;1:524-535.
- Zoratti M, Szabò I. The mitochondrial permeability transition. *Biochim Biophys Acta*. 1995;1241:139-176.
- Nisoli E, Falcone S, Tonello C, Cozzi V, Palomba L, Fiorani M, Pisconti A, Brunelli S, Cardile A, Francolini M, Cantoni O, Caruba MO, Moncada S, Clementi E. Mitochondrial biogenesis by NO yields functionally active mitochondria in mammals. *Proc Natl Acad Sci U S A*. 2004;101:16507-16512. Erratum in: *Proc Natl Acad Sci U S A*. 2005;102:5635.
- Marin W, Marin D, Ao X, Liu Y. Mitochondria as a therapeutic target for cardiac ischemiareperfusion injury (Review). *Int J Mol Med*. 2021;47:485-499.
- D'Amato M, Morra F, Di Meo I, Tiranti V. Mitochondrial transplantation in mitochondrial medicine: current challenges and future perspectives. *Int J Mol Sci*. 2023;24:1969.
- Pacak CA, Preble JM, Kondo H, Seibel P, Levitsky S, Del Nido PJ, Cowan DB, McCully JD. Actin-dependent mitochondrial internalization in cardiomyocytes: evidence for rescue of mitochondrial function. *Biol Open*. 2015;4:622-626.
- Zimmer HG. The isolated perfused heart and its pioneers. *News Physiol Sci*. 1998;13:203-210.
- Bell RM, Mocanu MM, Yellon DM. Retrograde heart perfusion: the Langendorff technique of isolated heart perfusion. *J Mol Cell Cardiol*. 2011;50:940-950.
- Lau C, Ranasinghe MG, Shiels I, Keates H, Pasloske K, Bellingham MC. Plasma pharmacokinetics of alfaxalone after a single intraperitoneal or intravenous injection of Alfaxan(®) in rats. *J Vet Pharmacol Ther*. 2013;36:516-520.
- Kim HK, Kang SW, Jeong SH, Kim N, Ko JH, Bang H, Park WS, Choi TH, Ha YR, Lee YS, Youm JB, Ko KS, Rhee BD, Han J. Identification of potential target genes of cardioprotection against ischemia-reperfusion injury by express sequence tags analysis in rat hearts. *J Cardiol*. 2012;60:98-110.
- Frezza C, Cipolat S, Scorrano L. Organelle isolation: functional mitochondria from mouse liver, muscle and cultured fibroblasts. *Nat Protoc*. 2007;2:287-295.
- Suzuki Y, Lyons JK, Yeung AC, Ikeno F. In vivo porcine model of reperfused myocardial infarction: in situ double staining to measure precise infarct area/area at risk. *Catheter Cardiovasc Interv*. 2008;71:100-107.
- Liu J, Liu C, Chen H, Cen H, Yang H, Liu P, Liu F, Ma L, Chen Q, Wang L. Tongguan capsule for treating myocardial ischemia-reperfusion injury: integrating network pharmacology and mechanism study. *Pharm Biol*. 2023;61:437-448.
- Kuznetsov AV, Veksler V, Gellerich FN, Saks V, Margreiter R, Kunz WS. Analysis of mitochondrial function in situ in permeabilized muscle fibers, tissues and cells. *Nat Protoc*. 2008;3:965-976.
- Gilmer LK, Roberts KN, Joy K, Sullivan PG, Scheff SW. Early mitochondrial dysfunction after cortical contusion injury. *J Neurotrauma*. 2009;26:1271-1280.
- Odinokova IV, Shalbuyeva N, Gukovskaya AS, Mareninova OA. Isolation of pancreatic mitochondria and measurement of their functional parameters. *Pancreapedia: Exocrine Pancreas Knowledge Base*. 2011. doi: 10.3998/panc.2011.25
- Salin K, Villasevil EM, Anderson GJ, Selman C, Chinopoulos C, Metcalfe NB. The RCR and ATP/O indices can give contradictory messages about mitochondrial efficiency. *Integr Comp Biol*. 2018;58:486-494.
- Lee D, Kim YW, Kim JH, Yang M, Bae H, Lim I, Bang H, Go KC, Yang GW, Rho YH, Park HS, Park EH, Ko JH. Improvement characteristics of bio-active materials coated fabric on rat muscular mitochondria. *Korean J Physiol Pharmacol*. 2015;19:283-289.
- Kong HL, Hou AJ, Liu NN, Chen BH, Dai SN, Huang HT. The effects of ginsenoside Rb1 on fatty acid  $\beta$ -oxidation, mediated by AMPK, in the failing heart. *Iran J Basic Med Sci*. 2018;21:731-737.
- Rooney JP, Ryde IT, Sanders LH, Howlett EH, Colton MD, Germ KE, Mayer GD, Greenamyre JT, Meyer JN. PCR based determination of mitochondrial DNA copy number in multiple species. *Methods Mol Biol*. 2015;1241:23-38.
- Rao X, Huang X, Zhou Z, Lin X. An improvement of the 2<sup>-</sup>(-delta delta CT) method for quantitative real-time polymerase chain reaction data analysis. *Biostat Bioinforma Biomath*. 2013;3:71-85.
- R Core Team. R: a language and environment for statistical computing. R Foundation for Statistical Computing; 2023.
- Butova XA, Myachina TA, Khokhlova AD. A combined Langendorff-injection technique for simultaneous isolation of single cardiomyocytes from atria and ventricles of the rat heart. *MethodsX*. 2020;8:101189.
- Lee D, Seo Y, Kim YW, Kim S, Bae H, Choi J, Lim I, Bang H, Kim JH, Ko JH. Far-infrared radiation stimulates platelet-derived growth



- factor mediated skeletal muscle cell migration through extracellular matrix-integrin signaling. *Korean J Physiol Pharmacol.* 2019;23:141-150.
30. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* 2000;28:27-30.
  31. Kaza AK, Wamala I, Friehs I, Kuebler JD, Rathod RH, Berra I, Ericsson M, Yao R, Thedsanamoorthy JK, Zurakowski D, Levitsky S, Del Nido PJ, Cowan DB, McCully JD. Myocardial rescue with autologous mitochondrial transplantation in a porcine model of ischemia/reperfusion. *J Thorac Cardiovasc Surg.* 2017;153:934-943.
  32. Cowan DB, Yao R, Akurathi V, Snay ER, Thedsanamoorthy JK, Zurakowski D, Ericsson M, Friehs I, Wu Y, Levitsky S, Del Nido PJ, Packard AB, McCully JD. Intracoronary delivery of mitochondria to the ischemic heart for cardioprotection. *PLoS One.* 2016;11:e0160889.
  33. Masuzawa A, Black KM, Pacak CA, Ericsson M, Barnett RJ, Drumm C, Seth P, Bloch DB, Levitsky S, Cowan DB, McCully JD. Transplantation of autologously derived mitochondria protects the heart from ischemia-reperfusion injury. *Am J Physiol Heart Circ Physiol.* 2013;304:H966-H982.
  34. Cowan DB, Yao R, Thedsanamoorthy JK, Zurakowski D, Del Nido PJ, McCully JD. Transit and integration of extracellular mitochondria in human heart cells. *Sci Rep.* 2017;7:17450.
  35. Gambardella J, Lombardi A, Santulli G. Metabolic flexibility of mitochondria plays a key role in balancing glucose and fatty acid metabolism in the diabetic heart. *Diabetes.* 2020;69:2054-2057.
  36. Warfel JD, Bermudez EM, Mendoza TM, Ghosh S, Zhang J, Elks CM, Mynatt R, Vandanmagsar B. Mitochondrial fat oxidation is essential for lipid-induced inflammation in skeletal muscle in mice. *Sci Rep.* 2016;6:37941.
  37. Tang L, Li J, Fu W, Wu W, Xu J. Suppression of FADS1 induces ROS generation, cell cycle arrest, and apoptosis in melanocytes: implications for vitiligo. *Aging (Albany NY).* 2019;11:11829-11843.
  38. Basheer WA, Fu Y, Shimura D, Xiao S, Agvastian S, Hernandez DM, Hitzeman TC, Hong T, Shaw RM. Stress response protein GJA1-20k promotes mitochondrial biogenesis, metabolic quiescence, and cardioprotection against ischemia/reperfusion injury. *JCI Insight.* 2018;3:e121900.