

## Corrigendum to 'Retrospective analysis of 30-day unplanned readmission after major abdominal surgery with reversal by sugammadex or neostigmine' (Br J Anaesth 2019; 122: 370–378)

Tak Kyu Oh<sup>1</sup>, Ah-Young Oh<sup>1,2,\*</sup>, Jung-Hee Ryu<sup>1,2</sup>, Bon-Wook Koo<sup>1</sup>, In-Ae Song<sup>1</sup>, Sun Woo Nam<sup>1</sup> and Hee-Jung Jee<sup>3</sup>

<sup>1</sup>Department of Anesthesiology and Pain Medicine, Seoul National University Bundang Hospital, Seongnam, Republic of Korea, <sup>2</sup>Department of Anesthesiology and Pain Medicine, Seoul National University College of Medicine, Seoul, Republic of Korea and <sup>3</sup>Department of Biostatistics, Korea University College of Medicine, Seoul, Republic of Korea

\*Corresponding author. E-mail: [ohahyoung@hanmail.net](mailto:ohahyoung@hanmail.net)

DOI of original article: <https://doi.org/10.1016/j.bja.2018.11.028>.

The authors regret that an error was present in Figure 1. The correct version of the figure appears below:

The authors would like to apologise for any inconvenience caused.




Fig 1. Flow chart of patient selection.