



Search for Neutrino Emission from Hard X-Ray AGN with IceCube

R. Abbasi¹, M. Ackermann², J. Adams³, S. K. Agarwalla^{4,67}, J. A. Aguilar⁵, M. Ahlers⁶, J.M. Alameddine⁷, N. M. Amin⁸, K. Andeen⁹, C. Argüelles¹⁰, Y. Ashida¹¹, S. Athanasiadou², L. Ausborn¹², S. N. Axani⁸, X. Bai¹³, A. Balagopal V.⁴, M. Baricevic⁴, S. W. Barwick¹⁴, S. Bash¹⁵, V. Basu⁴, R. Bay¹⁶, J. J. Beatty^{17,18}, J. Becker Tjus^{19,68}, J. Beise²⁰, C. Bellenghi¹⁵, C. Benning¹², S. BenZvi²¹, D. Berley²², E. Bernardini²³, D. Z. Besson²⁴, E. Blaufuss²², L. Bloom²⁵, S. Blot², F. Bontempo²⁶, J. Y. Book Motzkin¹⁰, C. Boscolo Meneguolo²³, S. Böser²⁷, O. Botner²⁰, J. Böttcher¹², J. Braun⁴, B. Brinson²⁸, J. Brostean-Kaiser², L. Brusa¹², R. T. Burley²⁹, D. Butterfield⁴, M. A. Campana³⁰, I. Caracas²⁷, K. Carloni¹⁰, J. Carpio^{31,32}, S. Chattopadhyay^{4,67}, N. Chau⁵, Z. Chen³³, D. Chirkin⁴, S. Choi^{34,35}, B. A. Clark²², A. Coleman²⁰, G. H. Collin³⁶, A. Connolly^{17,18}, J. M. Conrad³⁶, P. Coppin³⁷, R. Corley¹¹, P. Correa³⁷, D. F. Cowen^{38,39}, P. Dave²⁸, C. De Clercq³⁷, J. J. DeLaunay²⁵, D. Delgado¹⁰, S. Deng¹², A. Desai⁴, P. Desiati⁴, K. D. de Vries³⁷, G. de Wasseige⁴⁰, T. DeYoung⁴¹, A. Diaz³⁶, J. C. Díaz-Vélez⁴, P. Dierichs¹², M. Dittmer⁴², A. Domi⁴³, L. Draper¹¹, H. Dujmovic⁴, K. Dutta²⁷, M. A. DuVernois⁴, T. Ehrhardt²⁷, L. Eidenschink¹⁵, A. Eimer⁴³, P. Eller¹⁵, E. Ellinger⁴⁴, S. El Mentawi¹², D. Elsässer⁷, R. Engel^{26,45}, H. Erpenbeck⁴, J. Evans²², P. A. Evenson⁸, K. L. Fan²², K. Fang⁴, K. Farrag⁴⁶, A. R. Fazely⁴⁷, A. Fedynitch⁴⁸, N. Feigl⁴⁹, S. Fiedlschuster⁴³, C. Finley⁵⁰, L. Fischer², D. Fox³⁸, A. Franckowiak¹⁹, S. Fukami², P. Fürst¹², J. Gallagher⁵¹, E. Ganster¹², A. Garcia¹⁰, M. Garcia⁸, G. Garg^{4,67}, E. Genton^{10,40}, L. Gerhardt⁵², A. Ghadimi²⁵, C. Girard-Carillo²⁷, C. Glaser²⁰, T. Glüsenskamp^{20,43}, J. G. Gonzalez⁸, S. Goswami^{31,32,70}, A. Granados⁴¹, D. Grant⁴¹, S. J. Gray²², O. Gries¹², S. Griffin⁴, S. Griswold²¹, K. M. Groth⁶, C. Günther¹², P. Gutjahr⁷, C. Ha⁵³, C. Haack⁴³, A. Hallgren²⁰, L. Halve¹², F. Halzen⁴, H. Hamdaoui³³, M. Ha Minh¹⁵, M. Handt¹², K. Hanson⁴, J. Hardin³⁶, A. A. Harnisch⁴¹, P. Hatch⁵⁴, A. Haungs²⁶, J. Häussler¹², K. Helbing⁴⁴, J. Hellrung¹⁹, J. Hermannsgabner¹², L. Heuermann¹², N. Heyer²⁰, S. Hickford⁴⁴, A. Hidvegi⁵⁰, C. Hill⁴⁶, G. C. Hill²⁹, K. D. Hoffman²², S. Hori⁴, K. Hoshina^{4,69}, M. Hostert¹⁰, W. Hou²⁶, T. Huber²⁶, K. Hultqvist⁵⁰, M. Hünnefeld⁷, R. Hussain⁴, K. Hyman⁷, A. Ishihara⁴⁶, W. Iwakiri⁴⁶, M. Jacquart⁴, O. Janik⁴³, M. Jansson⁵⁰, G. S. Japaridze⁵⁵, M. Jeong¹¹, M. Jin¹⁰, B. J. P. Jones⁵⁶, N. Kamp¹⁰, D. Kang²⁶, W. Kang³⁴, X. Kang³⁰, A. Kappes⁴², D. Kappesser²⁷, L. Kardum⁷, T. Karg², M. Karl¹⁵, A. Karle⁴, A. Katil⁵⁷, U. Katz⁴³, M. Kauer⁴, J. L. Kelley⁴, M. Khanal¹¹, A. Khattee Zathul⁴, A. Kheirandish^{31,32}, J. Kiryluk³³, S. R. Klein^{16,52}, A. Kochocki⁴¹, R. Koirala⁸, H. Kolanoski⁴⁹, T. Kontrimas¹⁵, L. Köpke²⁷, C. Kopper⁴³, D. J. Koskinen⁶, P. Koundal⁸, M. Kovacevich³⁰, M. Kowalski^{2,49}, T. Kozynets⁶, J. Krishnamoorthi^{4,67}, K. Kruiswijk⁴⁰, E. Krupczak⁴¹, A. Kumar², E. Kun¹⁹, N. Kurahashi³⁰, N. Lad², C. Lagunas Gualda², M. Lamoureux⁴⁰, M. J. Larson²², S. Latseva¹², F. Lauber⁴⁴, J. P. Lazar⁴⁰, J. W. Lee³⁴, K. Leonard DeHolton³⁹, A. Leszczyńska⁸, J. Liao²⁸, M. Lincetto¹⁹, Y. T. Liu³⁹, M. Liubarska⁵⁷, E. Lohfink²⁷, C. Love³⁰, C. J. Lozano Mariscal⁴², L. Lu⁴, F. Lucarelli⁵⁸, W. Luszczak^{17,18}, Y. Lyu^{16,52}, J. Madsen⁴, E. Magnus³⁷, K. B. M. Mahn⁴¹, Y. Makino⁴, E. Manao¹⁵, S. Mancina^{4,23}, W. Marie Sainte⁴, I. C. Mariş⁵, S. Marka⁵⁹, Z. Marka⁵⁹, M. Marsee²⁵, I. Martinez-Soler¹⁰, R. Maruyama⁶⁰, F. Mayhew⁴¹, F. McNally⁶¹, J. V. Mead⁶, K. Meagher⁴, S. Mechbal², A. Medina¹⁸, M. Meier⁴⁶, Y. Merckx³⁷, L. Merten¹⁹, J. Micallef⁴¹, J. Mitchell⁴⁷, T. Montaruli⁵⁸, R. W. Moore⁵⁷, Y. Morii⁴⁶, R. Morse⁴, M. Moulai⁴, T. Mukherjee²⁶, R. Naab², R. Nagai⁴⁶, M. Nakos⁴, U. Naumann⁴⁴, J. Necker², A. Negi⁵⁶, L. Neste⁵⁰, M. Neumann⁴², H. Niederhausen⁴¹, M. U. Nisa⁴¹, K. Noda⁴⁶, A. Noell¹², A. Novikov⁸, A. Obertacke Pollmann⁴⁶, V. O'Dell⁴, B. Oeyen⁶², A. Olivas²², R. Orsoe¹⁵, J. Osborn⁴, E. O'Sullivan²⁰, H. Pandya⁸, N. Park⁵⁴, G. K. Parker⁵⁶, E. N. Paudel⁸, L. Paul¹³, C. Pérez de los Heros²⁰, T. Pernice², J. Peterson⁴, S. Philippen¹², A. Pizzuto⁴, M. Plum¹³, A. Pontén²⁰, Y. Popovych²⁷, M. Prado Rodriguez⁴, B. Pries⁴¹, G. C. Privon⁶³, R. Procter-Murphy²², G. T. Przybylski⁵², C. Raab⁴⁰, J. Rack-Helleis²⁷, M. Ravn²⁰, K. Rawlins⁶⁴, Z. Rechav⁴, A. Rehman⁸, P. Reichherzer¹⁹, E. Resconi¹⁵, S. Reusch², W. Rhode⁷, B. Riedel⁴, A. Rifaie¹², E. J. Roberts²⁹, S. Robertson^{16,52}, S. Rodan^{34,35}, G. Roellinghoff³⁴, M. Rongen⁴³, A. Rosted⁴⁶, C. Rott^{11,34}, T. Ruhe⁷, L. Ruohan¹⁵, D. Ryckbosch⁶², I. Safa⁴, J. Saffer⁴⁵, D. Salazar-Gallegos⁴¹, P. Sampathkumar²⁶, A. Sandroock⁴⁴, M. Santander²⁵, S. Sarkar⁵⁷, S. Sarkar⁶⁵, J. Savelberg¹², P. Savina⁴, P. Schaile¹⁵, M. Schaufel¹², H. Schieler²⁶, S. Schindler⁴³, B. Schlüter⁴², F. Schlüter⁵, N. Schmeisser⁴⁴, T. Schmidt²², J. Schneider⁴³, F. G. Schröder^{8,26}, L. Schumacher⁴³, S. Sclafani²², D. Seckel⁸, M. Seikh²⁴, M. Seo³⁴, S. Seunarine⁶⁶, P. Sevlé Myhr⁴⁰, R. Shah³⁰, S. Shefali⁴⁵, N. Shimizu⁴⁶, M. Silva⁴, B. Skrzypek¹⁶, B. Smithers⁵⁶, R. Snihur⁴, J. Soedingrekso⁷, A. Sègaard⁶, D. Soldin¹¹, P. Soldin¹², G. Sommani¹⁹, C. Spannfellner¹⁵, G. M. Spiczak⁶⁶, C. Spiering², M. Stamatikos¹⁸, T. Stanev⁸, T. Stezelberger⁵², T. Stürwald⁴⁴, T. Stuttard⁶, G. W. Sullivan²², I. Taboada²⁸, S. Ter-Antonyan⁴⁷, A. Terliuk¹⁵, M. Thiesmeyer¹², W. G. Thompson¹⁰, J. Thwaites⁴, S. Tilav⁸, K. Tollefson⁴¹, C. Tönnis³⁴, S. Toscano⁵, D. Tosi⁴, A. Trettin², R. Turcotte²⁶, J. P. Twagirayezu⁴¹, M. A. Unland Elorrieta⁴², A. K. Upadhyay^{4,67}, K. Upshaw⁴⁷, A. Vaidyanathan⁹, N. Valtonen-Mattila²⁰, J. Vandenbroucke⁴, N. van Eijndhoven³⁷, D. Vannerom³⁶, J. van Santen², J. Vara⁴², F. Varsi⁴⁵, J. Veitch-Michaelis⁴, M. Venugopal²⁶, M. Vereecken⁴⁰, S. Verpoest⁸, D. Veske⁵⁹, A. Vijai²², C. Walck⁵⁰, A. Wang²⁸, C. Weaver⁴¹, P. Weigel³⁶, A. Weindl²⁶, J. Weldert³⁹, A. Y. Wen¹⁰, C. Wendt⁴, J. Werthebach⁷, M. Weyrauch²⁶, N. Whitehorn⁴¹, C. H. Wiebusch¹²

D. R. Williams²⁵, L. Witthaus⁷ , A. Wolf¹², M. Wolf¹⁵ , G. Wrede⁴³, X. W. Xu⁴⁷, J. P. Yanez⁵⁷, E. Yildizci⁴, S. Yoshida⁴⁶ ,
R. Young²⁴, S. Yu¹¹ , T. Yuan⁴ , Z. Zhang³³, P. Zhelnin¹⁰ , P. Zilberman⁴, and M. Zimmerman⁴

IceCube Collaboration

- ¹ Department of Physics, Loyola University Chicago, Chicago, IL 60660, USA
- ² Deutsches Elektronen-Synchrotron DESY, Platanenallee 6, D-15738 Zeuthen, Germany
- ³ Dept. of Physics and Astronomy, University of Canterbury, Private Bag 4800, Christchurch, New Zealand
- ⁴ Dept. of Physics and Wisconsin IceCube Particle Astrophysics Center, University of Wisconsin—Madison, Madison, WI 53706, USA
- ⁵ Université Libre de Bruxelles, Science Faculty CP230, B-1050 Brussels, Belgium
- ⁶ Niels Bohr Institute, University of Copenhagen, DK-2100 Copenhagen, Denmark
- ⁷ Dept. of Physics, TU Dortmund University, D-44221 Dortmund, Germany
- ⁸ Bartol Research Institute and Dept. of Physics and Astronomy, University of Delaware, Newark, DE 19716, USA
- ⁹ Department of Physics, Marquette University, Milwaukee, WI 53201, USA
- ¹⁰ Department of Physics and Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge, MA 02138, USA
- ¹¹ Department of Physics and Astronomy, University of Utah, Salt Lake City, UT 84112, USA
- ¹² III. Physikalisches Institut, RWTH Aachen University, D-52056 Aachen, Germany
- ¹³ Physics Department, South Dakota School of Mines and Technology, Rapid City, SD 57701, USA
- ¹⁴ Dept. of Physics and Astronomy, University of California, Irvine, CA 92697, USA
- ¹⁵ Physik-department, Technische Universität München, D-85748 Garching, Germany
- ¹⁶ Dept. of Physics, University of California, Berkeley, CA 94720, USA
- ¹⁷ Dept. of Astronomy, Ohio State University, Columbus, OH 43210, USA
- ¹⁸ Dept. of Physics and Center for Cosmology and Astro-Particle Physics, Ohio State University, Columbus, OH 43210, USA
- ¹⁹ Fakultät für Physik & Astronomie, Ruhr-Universität Bochum, D-44780 Bochum, Germany
- ²⁰ Dept. of Physics and Astronomy, Uppsala University, Box 516, SE-75120 Uppsala, Sweden
- ²¹ Dept. of Physics and Astronomy, University of Rochester, Rochester, NY 14627, USA
- ²² Dept. of Physics, University of Maryland, College Park, MD 20742, USA
- ²³ Dipartimento di Fisica e Astronomia Galileo Galilei, Università Degli Studi di Padova, I-35122 Padova PD, Italy
- ²⁴ Dept. of Physics and Astronomy, University of Kansas, Lawrence, KS 66045, USA
- ²⁵ Dept. of Physics and Astronomy, University of Alabama, Tuscaloosa, AL 35487, USA
- ²⁶ Karlsruhe Institute of Technology, Institute for Astroparticle Physics, D-76021 Karlsruhe, Germany
- ²⁷ Institute of Physics, University of Mainz, Staudinger Weg 7, D-55099 Mainz, Germany
- ²⁸ School of Physics and Center for Relativistic Astrophysics, Georgia Institute of Technology, Atlanta, GA 30332, USA
- ²⁹ Department of Physics, University of Adelaide, Adelaide, 5005, Australia
- ³⁰ Dept. of Physics, Drexel University, 3141 Chestnut Street, Philadelphia, PA 19104, USA
- ³¹ Department of Physics & Astronomy, University of Nevada, Las Vegas, NV 89154, USA; sreetama.goswami@unlv.edu
- ³² Nevada Center for Astrophysics, University of Nevada, Las Vegas, NV 89154, USA
- ³³ Dept. of Physics and Astronomy, Stony Brook University, Stony Brook, NY 11794-3800, USA
- ³⁴ Dept. of Physics, Sungkyunkwan University, Suwon 16419, Republic of Korea
- ³⁵ Institute of Basic Science, Sungkyunkwan University, Suwon 16419, Republic of Korea
- ³⁶ Dept. of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- ³⁷ Vrije Universiteit Brussel (VUB), Dienst ELEM, B-1050 Brussels, Belgium
- ³⁸ Dept. of Astronomy and Astrophysics, Pennsylvania State University, University Park, PA 16802, USA
- ³⁹ Dept. of Physics, Pennsylvania State University, University Park, PA 16802, USA
- ⁴⁰ Centre for Cosmology, Particle Physics and Phenomenology—CP3, Université catholique de Louvain, Louvain-la-Neuve, Belgium
- ⁴¹ Dept. of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA
- ⁴² Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, D-48149 Münster, Germany
- ⁴³ Erlangen Centre for Astroparticle Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg, D-91058 Erlangen, Germany
- ⁴⁴ Dept. of Physics, University of Wuppertal, D-42119 Wuppertal, Germany
- ⁴⁵ Karlsruhe Institute of Technology, Institute of Experimental Particle Physics, D-76021 Karlsruhe, Germany
- ⁴⁶ Dept. of Physics and The International Center for Hadron Astrophysics, Chiba University, Chiba 263-8522, Japan
- ⁴⁷ Dept. of Physics, Southern University, Baton Rouge, LA 70813, USA
- ⁴⁸ Institute of Physics, Academia Sinica, Taipei, 11529, Taiwan
- ⁴⁹ Institut für Physik, Humboldt-Universität zu Berlin, D-12489 Berlin, Germany
- ⁵⁰ Oskar Klein Centre and Dept. of Physics, Stockholm University, SE-10691 Stockholm, Sweden
- ⁵¹ Dept. of Astronomy, University of Wisconsin—Madison, Madison, WI 53706, USA
- ⁵² Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA
- ⁵³ Dept. of Physics, Chung-Ang University, Seoul 06974, Republic of Korea
- ⁵⁴ Dept. of Physics, Engineering Physics, and Astronomy, Queen's University, Kingston, ON K7L 3N6, Canada
- ⁵⁵ CTSPS, Clark-Atlanta University, Atlanta, GA 30314, USA
- ⁵⁶ Dept. of Physics, University of Texas at Arlington, 502 Yates St., Science Hall Rm 108, Box 19059, Arlington, TX 76019, USA
- ⁵⁷ Dept. of Physics, University of Alberta, Edmonton, Alberta, T6G 2E1, Canada
- ⁵⁸ Département de physique nucléaire et corpusculaire, Université de Genève, CH-1211 Genève, Switzerland
- ⁵⁹ Columbia Astrophysics and Nevis Laboratories, Columbia University, New York, NY 10027, USA
- ⁶⁰ Dept. of Physics, Yale University, New Haven, CT 06520, USA
- ⁶¹ Department of Physics, Mercer University, Macon, GA 31207-0001, USA
- ⁶² Dept. of Physics and Astronomy, University of Gent, B-9000 Gent, Belgium
- ⁶³ National Radio Astronomy Observatory, Charlottesville, VA 22903, USA
- ⁶⁴ Dept. of Physics and Astronomy, University of Alaska Anchorage, 3211 Providence Dr., Anchorage, AK 99508, USA

⁶⁵ Dept. of Physics, University of Oxford, Parks Road, Oxford OX1 3PU, UK⁶⁶ Dept. of Physics, University of Wisconsin, River Falls, WI 54022, USA

Received 2024 June 21; revised 2024 December 9; accepted 2025 January 9; published 2025 March 4

Abstract

Active galactic nuclei (AGN) are promising candidate sources of high-energy astrophysical neutrinos, since they provide environments rich in matter and photon targets where cosmic-ray interactions may lead to the production of gamma rays and neutrinos. We searched for high-energy neutrino emission from AGN using the Swift-BAT Spectroscopic Survey catalog of hard X-ray sources and 12 yr of IceCube muon track data. First, upon performing a stacked search, no significant emission was found. Second, we searched for neutrinos from a list of 43 candidate sources and found an excess from the direction of two sources, the Seyfert galaxies NGC 1068 and NGC 4151. We observed NGC 1068 at flux $\phi_{\nu_\mu+\bar{\nu}_\mu} = 4.02^{+1.58}_{-1.52} \times 10^{-11} \text{ TeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$ normalized at 1 TeV, with a power-law spectral index $\gamma = 3.10^{+0.26}_{-0.22}$, consistent with previous IceCube results. The observation of a neutrino excess from the direction of NGC 4151 is at a posttrial significance of 2.9σ . If interpreted as an astrophysical signal, the excess observed from NGC 4151 corresponds to a flux $\phi_{\nu_\mu+\bar{\nu}_\mu} = 1.51^{+0.99}_{-0.81} \times 10^{-11} \text{ TeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$ normalized at 1 TeV and $\gamma = 2.83^{+0.35}_{-0.28}$.

Unified Astronomy Thesaurus concepts: [Neutrino astronomy \(1100\)](#); [High energy astrophysics \(739\)](#); [Active galaxies \(17\)](#); [Seyfert galaxies \(1447\)](#)

1. Introduction

In 2013, IceCube reported the detection of high-energy neutrinos of astrophysical origin with a sky distribution consistent with isotropy (M. G. Aartsen et al. 2013). This steady flux of astrophysical neutrinos observed in the energy range TeV–PeV has so far been characterized by a single power-law model (M. G. Aartsen et al. 2020a; R. Abbasi et al. 2021a, 2022a, 2022b). To find the origin of the neutrino emission, IceCube has searched for neutrino excesses from various astronomical source classes, e.g., supernovae (R. Abbasi et al. 2023a), pulsar wind nebulae (M. G. Aartsen et al. 2020b), ultraluminous infrared galaxies (R. Abbasi et al. 2022c), X-ray binaries (R. Abbasi et al. 2022d), galaxy clusters (R. Abbasi et al. 2022e), GRBs (R. Abbasi et al. 2022f), and others. Evidence of neutrinos from the galactic plane (R. Abbasi et al. 2023b), at a significance level of 4.5σ , indicates a contribution of $\sim 6\%$ – 13% to the all-sky astrophysical flux at 30 TeV. So far, no significant evidence of a dominant source class has been identified.

Active galactic nuclei (AGN) are promising source candidates, owing to their electromagnetic emission (E. Lusso et al. 2012), which spans several orders of magnitude in luminosity, as well as their likely ability to accelerate charged particles to ultra-high energies, $E \sim 10^{20} \text{ eV}$ (R. Lovelace 1976; R. Mbarek & D. Caprioli 2019). AGN are the compact innermost regions of galaxies that, in some cases, are bright enough to outshine the host galaxy. Matter spirals into a central supermassive black hole (SMBH), forming an accretion disk releasing vast amounts of energy. The accretion disk is enveloped in a dusty torus that obstructs the view of the AGN core. AGN ubiquitously show evidence of thermal plasma being ejected with a range of velocities

and opening angles. Many AGN are observed with ejected winds over wide angles, while some outflows are relativistic. A subset of AGN constituting less than 10% are distinguishable by highly collimated and ultrarelativistic jets (A. Königl 2006). The orientation of the jet with respect to the observer and the degree of obscuration by the dusty torus result in differences in the observed spectral features (R. Antonucci 1993; C. M. Urry & P. Padovani 1995). Blazars are the brightest AGN, with their jets pointing toward the Earth.

The emission of AGN spans the entire electromagnetic spectrum, from radio waves to gamma rays (R. A. Edelson et al. 1996). The gases contained within the inner boundaries of the torus are under the gravitational effect of the SMBH and have a very high velocity. Observationally, they are identified by Doppler-broadened emission lines, and the region is referred to as the broad-line region (BLR). The other key region around AGN is the narrow-line region (NLR) ascribed to the gases around the torus with lower velocities and extending to a few kiloparsecs. For obscured AGN, only the NLR is observed, while the BLR remains hidden by the torus. The emission from the jets is nonthermal, with intense radiation of high apparent luminosities resulting from Doppler boosting. A prominent source of X-rays within the AGN is the corona, sometimes referred to as the hot corona. Though the exact geometry, size, and location of the corona remain unknown, the hot corona is believed to be situated above the inner region of the accretion disk and is responsible for reprocessing the UV photons from the accretion disk to X-ray energies by the inverse Compton process. There is a possible feedback mechanism between various outflows and the accretion disk, where the hot corona also plays an important role. The accretion disk is believed to power the jets in many AGN. The interplay of physical processes underlying these salient features of AGN gives rise to myriad observable emission features.

AGN have long been considered as production sites of astrophysical neutrinos (see, e.g., F. W. Stecker et al. 1991; K. Mannheim 1995; F. Halzen & E. Zas 1997). This hypothesis is supported by the detection of a high-energy neutrino in 2017 by IceCube in spatial and temporal coincidence with the blazar TXS 0506+056, in a flaring state (M. G. Aartsen et al. 2018a), and the evidence of neutrinos from the active galaxy NGC 1068 (M. G. Aartsen et al. 2018b; R. Abbasi et al. 2022g). Neutrino emission models suggest that high-energy neutrinos produced in AGN by the decay of charged pions are accompanied by a high-

⁶⁷ Also at Institute of Physics, Sachivalaya Marg, Sainik School Post, Bhubaneswar 751005, India.

⁶⁸ Also at Department of Space, Earth and Environment, Chalmers University of Technology, 412 96 Gothenburg, Sweden.

⁶⁹ Also at Earthquake Research Institute, University of Tokyo, Bunkyo, Tokyo 113-0032, Japan.

⁷⁰ Corresponding author.



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

energy gamma-ray flux arising from the decay of neutral pions (see, e.g., S. Gao et al. 2017; A. Keivani et al. 2018; K. Murase & F. W. Stecker 2022). These models propose that regions within AGN, such as the corona and jet, which are rich in photons and matter as indicated by electromagnetic signatures, are possible sites of neutrino production. Previous studies have searched for neutrino emission from gamma-ray-bright blazars (see, e.g., M. G. Aartsen et al. 2017a; M. Huber 2019) and concluded that this source class contributes only a small fraction toward the total flux of astrophysical neutrinos observed by IceCube. The lack of a simple association between gamma-ray AGN and neutrinos motivated reevaluations and new implementations of models (see, e.g., S. Gao et al. 2019; M. Petropoulou et al. 2020; P. Padovani 2023) where neutrinos escape the AGN environment but gamma rays interact with matter or photon fields surrounding the active core and cascade down to MeV gamma rays or hard X-rays. These predictions led to searches for neutrino emission from sources detected in lower-energy bands, which found background-compatible results (see, e.g., R. Abbasi et al. 2022h, 2022i).

In this study, we search for a possible correlation between high-energy neutrinos and AGN detected in hard X-rays. We use the Swift-BAT AGN Spectroscopic Survey (BASS; M. Koss et al. 2017), which is the most complete all-sky catalog of hard X-ray AGN detected in the 14–195 keV range, and we perform two different analyses as detailed in Sections 3.1 and 3.2.⁷¹

2. Data Set and Hard X-Ray Sources

2.1. The Neutrino Data Set

The IceCube Neutrino Observatory is situated at the Amundsen–Scott South Pole Station, encompassing a cubic kilometer of ice (M. G. Aartsen et al. 2017b) that acts as the detection medium. Neutrino interactions produce secondary charged particles that move at relativistic speeds, emitting Cherenkov radiation. IceCube detects this radiation using 5160 digital optical modules deployed on 86 cables known as “strings” at depths between 1450 m and 2450 m from the surface forming a three-dimensional array (M. G. Aartsen et al. 2017c). The events detected by IceCube are classified based on their optical signatures. The $\sim$ kilometer-long track-like events are produced by charged-current (CC) interactions of muon neutrinos (M. G. Aartsen et al. 2013). The roughly spherical cascade events are produced by all other CC interactions and neutral current (NC) interactions (M. G. Aartsen et al. 2020a). The background for the astrophysical neutrino flux is dominated by the atmospheric muons and neutrinos produced in cosmic-ray air showers (M. G. Aartsen et al. 2016).

We used IceCube data recorded from 2008 to 2020 for the analyses discussed in this paper. The data set has the same selection criteria as in M. G. Aartsen et al. (2020c) applied to an additional two years of data. The detector operated with partial configurations of 40, 59, and 79 strings from 2008 to 2011 and with the full 86 string configuration from 2011 onward (see Table 3). The sample covers the entire sky and is largely made of through-going muon tracks (M. G. Aartsen et al. 2020c). The data set was chosen due to the tracks having a better angular resolution compared to other event morphologies, with a median $<1^\circ$ above 1 TeV (M. G. Aartsen et al. 2017b; R. Abbasi et al. 2021b). See M. G. Aartsen et al.

(2020c) for more information about the selection criteria and the reconstruction methods for direction and energy.

2.2. Hard X-Ray Sources

The BASS catalog (M. Koss et al. 2017; C. Ricci et al. 2017) used in this study contains AGN selected from sources in the 70 month Swift-BAT hard X-ray survey (W. H. Baumgartner et al. 2013). As the most complete hard X-ray AGN catalog, this provides an opportunity to perform a sensitive study of the potential contribution of hard X-ray AGN to the neutrino flux. Only six of the 838 sources do not have redshift estimates. Most are nearby, with a median of 0.04. We used the updated source class and redshift values from the BASS DR-2 catalog (M. J. Koss et al. 2022a, 2022b).

Gas and dust surrounding AGN can act as targets for hadronic interactions. The amount of material, therefore, may impact neutrino production and is quantified by the column density (N_H) of neutral hydrogen along the line of sight. Due to their high penetrating power, hard X-rays are well suited to probe the obscured AGN core. Only two of the 838 sources do not have column density estimates. Since we have used column density, N_H , to classify the sources in the first analysis (see Section 3.1), the two sources without the parameter measured were excluded.

3. Analysis

We perform two analyses: a stacked search and an individual source search, described in Sections 3.1 and 3.2, respectively. The analyses in this paper are time-integrated searches performed using an unbinned likelihood ratio method that tests how well a hypothesis representing a combination of the signal and the diffuse background describes the data, as compared to a null hypothesis representing only the background (A. Achterberg et al. 2006; J. Braun et al. 2008). We assume a point source emits neutrinos following a power-law energy spectrum, with the neutrino flux per flavor given by $\phi_{\nu_\mu+\bar{\nu}_\mu}(E_\nu) = \phi_0 \cdot (E_\nu/E_0)^{-\gamma}$, where E_0 is the normalization energy, ϕ_0 is the normalization flux, E_ν is the neutrino energy, and γ is the spectral index. The likelihood function is defined as follows for a single point source (M. G. Aartsen et al. 2019a):

$$\mathcal{L}(n_s, \gamma) = \prod_i^N \left(\frac{n_s}{N} \mathcal{S}_i(\mathbf{x}_i, \sigma_i, E_i, \gamma) + \left(1 - \frac{n_s}{N} \right) \mathcal{B}_i(\delta_i, E_i) \right), \quad (1)$$

where n_s is the number of signal neutrinos. $\mathcal{S}_i$, the signal probability density functions (PDFs), have spatial and energy components where $\mathbf{x}_i$, E_i , and σ_i are the reconstructed direction, energy, and estimated directional uncertainty of each candidate neutrino event in the data set, respectively. The spatial part of $\mathcal{B}_i$, the background PDF, depends only on decl. because the distribution of events in R.A. is uniform over long timescales for IceCube. The signal PDF is constructed using Monte Carlo simulated events and the background PDF using data randomized in R.A. The number of signal events n_s and the power-law index γ are determined by maximizing the likelihood $\mathcal{L}$, which yields best-fit parameter values $\hat{n}_s$ and $\hat{\gamma}$.

3.1. The Stacked Search

A stacking analysis evaluates the cumulative neutrino signal from a given population of sources. In this method, the signal

⁷¹ Swift-BAT AGN Spectroscopic Survey

from all sources is summed to increase the sensitivity and possibility of detection. The likelihood function used in this analysis is similar to Equation (1) but with a stacked signal PDF where each source term contains two types of weights:

1. ω is a theoretical weight, which is a chosen observed physical property hypothesized to be proportional to the neutrino flux. In this analysis, we use the intrinsic X-ray flux (see Appendix A.2) in the 14–195 keV range for each source as a weight. The values are provided in the BASS catalog.
2. $R(\delta_k, \gamma)$ is a detector weight, which is the sensitivity of the IceCube detector for a source at a decl. δ with a spectral index γ .

The stacked signal PDF for the i th neutrino event, where $i \in \{1, 2, \dots, N\}$, $k \in \{1, 2, \dots, M\}$, $\mathbf{x}_{S_k}$ is the position of the k th source, and M is the number of sources, is

$$\mathcal{S}_{i,\text{Stacked}} = \frac{\sum_{k=1}^M \omega^k R^k(\delta_k, \gamma) \cdot \mathcal{S}_i^k(\mathbf{x}_i, \mathbf{x}_{S_k}, \sigma_i, E_i, \gamma)}{\sum_{k=1}^M \omega^k R^k(\delta_k, \gamma)}. \quad (2)$$

For this analysis, we divide our catalog using two different criteria: AGN type and column density. We divide the catalog into blazar and nonblazar AGN according to the first criterion. Subdividing the catalog by the column density (N_H) results in three samples: unobscured AGN ($N_H < 10^{22} \text{ cm}^{-2}$), obscured AGN ($10^{22} \text{ cm}^{-2} < N_H < 10^{24} \text{ cm}^{-2}$), and Compton-thick (CT) AGN ($N_H > 10^{24} \text{ cm}^{-2}$). Enough absorbing matter surrounds CT sources, resulting in a large optical depth for Compton scattering. We also test an unphysical hypothesis of equal neutrino emission from all AGN as an unbiased basis for comparison with other tests. In effect, we test a total of seven hypotheses. The first six use flux weights for each of the following AGN classes: (i) all 836 AGN, (ii) 104 blazars, (iii) 732 nonblazars, (iv) 457 unobscured AGN, (v) 323 obscured AGN, and (vi) 56 CT AGN, while the last (vii) uses equal weights for all 836 AGN.

Any flux measurement (or flux upper limit) derived from the stacking analysis applies solely to the selected AGN of a given class that forms a subset of the entire source population in the Universe. Many AGN fall below the instrument's detection threshold, due to large distance, low intrinsic luminosity, or high obscuration. To consider all the undetected sources, we scale the fluxes by a catalog completeness factor. In evaluating the contribution from the entire source population in the Universe, we integrate over luminosity functions for each source class (blazars from L. Marcotulli et al. 2022 and nonblazar AGN from Y. Ueda et al. 2014) and assume the neutrino flux is proportional to the intrinsic X-ray flux. See Appendix A.4.1 for the details of this method and the estimated catalog completeness factors for each AGN class.

3.2. The Individual Source Search

In this analysis, we study sources that are most likely to be individually detected by IceCube. We hypothesize that neutrino emission is proportional to the intrinsic hard X-ray flux in the 14–195 keV range. The strong decl. dependence of the IceCube sensitivity to neutrino point sources implies that some sources have a higher likelihood of being detected because of their sky location.

To select the sources to be probed individually, we define a “figure of merit” (FOM) as the ratio of the intrinsic hard X-ray flux in the 14–195 keV range to the neutrino flux sensitivity at the source decl. and rank all AGN in our catalog according to

this value. Since a power-law energy spectrum with spectral index $\gamma = 2.5$ approximately describes the astrophysical diffuse flux, we considered the same for the neutrino flux from each source. Starting from the source with the highest value of FOM, we select all the sources that have up to a factor of 10 lower FOM than the top source (see Appendix A.5). The 43 AGN thus selected from the BASS catalog used to perform the second analysis are listed in Table 5. Most of these selected sources are in the Northern Hemisphere, since IceCube has better sensitivity in the north as compared to the southern sky.

Using the method described in Section 3, we find the best-fit values of n_s and γ for each source. NGC 1068 is present in the list used for this analysis. Since a previous IceCube study (R. Abbasi et al. 2022g) has identified evidence for neutrino emission at a significance of 4.2σ , we do not report a posttrial significance for it.

4. Results

Stacking analysis. We tested seven hypotheses: first, by using the entire AGN sample and five subsamples with flux weights, and second, by all AGN using equal weights. For each hypothesis, we computed pretrial p -value (p_{local}). Since we test multiple hypotheses by performing several statistical tests and using samples having considerable overlaps, the p_{local} for the most significant test is corrected using a trial factor (see Appendix A.4.2). For the analysis performed using all 836 AGN with equal weights, the best-fit number of neutrinos is zero and p_{local} is 1.0. The hypothesis showing the highest statistical significance is an obscured AGN ($10^{22} \text{ cm}^{-2} < N_H < 10^{24} \text{ cm}^{-2}$) at a posttrial significance of 2.1σ .

Since there is no significant evidence of neutrino emission, we derive 90% confidence level (CL) flux upper limits per flavor for power-law spectra with three different spectral indices: $E^{-3.0}$, $E^{-2.5}$, and $E^{-2.0}$ (see Table 1). For spectral indices of $\gamma = 3.0, 2.5$, and 2.0 , the analyses with all AGN are most sensitive in the energy ranges 0.5 TeV to 0.1 PeV, 2.5 TeV to 1.1 PeV, and 6 TeV to 10 PeV, respectively (see Figures 1 and 7). We used these energy ranges for all AGN samples, since they are expected to be comparable (see Appendix A.4.3). The fluxes are scaled by the catalog completeness factor to take into account all unresolved sources (see Appendix A.4.1).

We constrain the maximum contribution from all hard X-ray blazars to the astrophysical diffuse flux in R. Abbasi et al. (2022b) to be less than 7%. This is evaluated using the flux upper limits obtained for blazars considering a power-law spectrum of $E^{-2.5}$ at 100 TeV and corrected for catalog completeness. Notably, the completeness factor for the blazars has large uncertainties as given in Table 4. Previously, another IceCube result (M. G. Aartsen et al. 2017a) constrained the contribution from Fermi-2LAC blazars to be 27%. There exists only a small overlap between the blazars in this analysis and the previous study. They independently conclude that blazars emit a small fraction of neutrinos. The flux upper limits obtained for all AGN or nonblazar AGN, considering a power-law spectrum with an index $\gamma = 3.0$ at an energy ~ 30 TeV, are comparable to the diffuse flux. Thus, we cannot exclude the possibility that a major fraction of the diffuse flux can be attributed to these source classes.

Individual source search. From the list of 43 sources, the two sources with pretrial significance $> 3\sigma$ are Seyfert galaxies NGC 1068 and NGC 4151, and their results are listed in Table 2. NGC 1068 is found at a flux and spectral index consistent with previous results (R. Abbasi et al. 2022g). Since it was observed




Figure 1. The 90% CL flux upper limit for (left) all AGN in the catalog, blazars, and nonblazars, and (right) for unobscured, obscured, and Compton-thick sources. Flux upper limits are shown here for a spectral index of 3.0, since most samples result in a best-fit value of ~ 3 . These flux upper limits are displayed within the most relevant energy range for the analysis and are scaled using the catalog completeness factor described in Appendix A.4.1. For reference, the plot shows the all-sky diffuse flux from muon tracks (light blue dotted line; R. Abbasi et al. 2022b) and from cascades (purple; M. G. Aartsen et al. 2019b).




Figure 2. Left: 68% CL contour lines for NGC 1068 (blue) and NGC 4151 (black), along with the best-fit values (crosses) of flux normalization and spectral index. Right: flux as a function of energy from NGC 1068 (black) and NGC 4151 (orange), obtained from the current analysis. We compare them to the all-sky total astrophysical neutrino flux showing 9.5 yr of northern tracks data (R. Abbasi et al. 2022b) and six years of cascade data (M. G. Aartsen et al. 2020a). The all-sky fluxes shown in the plot are calculated by integrating the measured quantities from R. Abbasi et al. (2022b) and from M. G. Aartsen et al. (2020a) over 4π . Lines are shown for the best-fit values, and the shaded region represents the 68% CL region, which represents the statistical uncertainties. The systematic uncertainties are subdominant for this analysis. The energy range corresponds to the majority of the contribution toward the neutrino excess: we found the central 68% of the contribution to the test statistic to be from neutrinos with energies of 0.7 TeV–15 TeV for NGC 1068 and 4.3 TeV–65.2 TeV for NGC 4151.

Table 1
Summary of Results from the Stacked Search

Sample	No. of Sources	$\hat{n}_s$	$\hat{\gamma}$	P_{local} (Significance)	$\phi_{90\%}$ ($\gamma = 2.0$)	$\phi_{90\%}$ ($\gamma = 2.5$)	$\phi_{90\%}$ ($\gamma = 3.0$)
All AGN	836	161	2.89	0.01 (2.2 σ)	0.68	8.97	39.25
Blazars	104	10	2.04	0.14 (1.1 σ)	0.16	1.83	7.38
Nonblazars	732	180	3.02	0.01 (2.4 σ)	0.75	9.89	41.94
Compton-thick	56	45	3.14	0.12 (1.2 σ)	0.30	3.63	14.40
Obscured	323	148	2.91	0.003 (2.7 σ)	0.57	6.85	27.06
Unobscured	457	0	0	1.00 (0.0 σ)	0.24	3.02	13.85

Note. Flux upper limit normalized at 1 TeV is in units of $10^{-11} \text{ TeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$. These upper limits are scaled by respective completeness factors and shown for different energies in Figure 1. The best-fit number of neutrinos, $\hat{n}_s$, is the sum over all sources in the sample.




Figure 3. Top row: high-resolution scan of the sky around the region of the two most significant source positions (a) NGC 1068 and (b) NGC 4151. For details of the method used to obtain the scans, see R. Abbasi et al. (2022g). The white cross shows the best-fit position, and the red star shows the source position cataloged in BASS. The solid and dashed contours show the 68% and 95% CL regions around the sources, respectively. Bottom row: number of events as a function of the square of the angular distances from the source for (c) NGC 1068 and (d) NGC 4151. The best-fit astrophysical neutrino signal, the background, and their sum are shown in purple, orange, and gray, respectively. They are obtained using Monte Carlo simulations. The neutrino event data points are shown in black with error bars.

Table 2
The Two Most Significant Sources from the Individual Source Search

Name	Type	$\hat{n}_s$	$\hat{\gamma}$	$\phi_{\nu_\mu + \bar{\nu}_\mu}^{1 \text{ TeV}}$ ($\text{TeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$)	p_{local} (Significance)	posttrial p -value (Significance)
NGC 1068	Compton-thick Seyfert	81.7	$3.10^{+0.26}_{-0.22}$	$4.02^{+1.58}_{-1.52} \times 10^{-11}$	1.27×10^{-6} (4.7σ)	...
NGC 4151	Obscured Seyfert	49.8	$2.83^{+0.35}_{-0.28}$	$1.51^{+0.99}_{-0.81} \times 10^{-11}$	3.99×10^{-5} (3.9σ)	1.67×10^{-3} (2.9σ)

with a high significance, we exclude it from the estimation of the posttrial p -value, evaluated as $1 - (1 - p_{\text{local}})^N$, where N is the number of sources. NGC 4151, the source with the highest FOM among the 43 AGN used for the individual source search, is found to have a posttrial p -value of 1.67×10^{-3} (2.9σ). An excess of neutrinos was previously reported from a direction ($\sim 0.18^\circ$ angular deviation) compatible with that of NGC 4151 in R. Abbasi et al. (2022g).

While we do not reject the background hypothesis at $>3\sigma$ significance, we can consider the excess to be astrophysical in origin and derive a best-fit flux normalization. Figure 2 (left) shows the best-fit values and 68% CL contours from a likelihood scan using Wilk's theorem. The contours are evaluated at 5 TeV, where the correlation between fit parameters is minimal. The power-law spectra and uncertainties are shown in Figure 2 (right).

The uncertainties are primarily statistical and the systematic uncertainties are subdominant, as evident from the estimates found in multiple searches of neutrino sources, e.g., M. G. Aartsen et al. (2014b, 2017b).

We scan the region around the most significant sources as shown in Figure 3 (top). It shows the offset of the best-fit positions as a result of the analysis from the cataloged coordinates of the sources. The known source position lies within the 68% contour line for both sources. See Figure 3 (bottom) for the distribution of the neutrino events with increasing angular distance from the sources.

5. Discussion

The most significant neutrino sources observed are two Seyfert galaxies, NGC 1068 and NGC 4151. However, the stacking

analysis of nonblazar AGN (731 of 732 nonblazar AGN are Seyfert galaxies) shows no significant emission. It should be noted that we have not tested the predictions of any specific model while performing the stacking analyses. The apparent disagreement indicates that the potential neutrino emission is not directly proportional to the hard X-ray flux. The stacking analysis considering only blazars, a class of jetted AGN, also did not show any significant emission. This is consistent with the previous neutrino searches that have not found a significant correlation between blazars and neutrinos detected by IceCube. Several models suggest that the neutrino fluxes from blazars possibly peak around 100 PeV, where their fluxes are too small to be detected by IceCube (X. Rodrigues et al. 2024).

NGC 1068 and NGC 4151 are two nearby AGN, with photometric distances 11.14 ± 0.54 Mpc, 14.20 ± 0.88 Mpc (N. A. Tikhonov & O. A. Galazutdinova 2021), that are obscured ($N_H = 10^{24.95} \text{ cm}^{-2}$, $10^{22.71} \text{ cm}^{-2}$) and bright in X-rays (flux = $2.06 \times 10^{-10} \text{ erg cm}^{-2} \text{ s}^{-1}$, $5.26 \times 10^{-10} \text{ erg cm}^{-2} \text{ s}^{-1}$). They have similar SMBH masses, $1.3 \times 10^7 M_\odot$ (J.-M. Wang et al. 2020) and $1.66 \times 10^7 M_\odot$ (M. C. Bentz et al. 2022). However, while NGC 1068 is a starburst galaxy, NGC 4151 shows little evidence of star formation (C. G. Mundell et al. 1999). Both sources show evidence of jet–disk interaction (D. R. A. Williams et al. 2020; G. Venturi et al. 2021). The relativistic jets and the accretion disk are the two most promising sites for efficient particle acceleration, which is essential for neutrino production.

Acceleration mechanisms describing neutrino production predict a comparable flux of gamma rays and neutrinos. However, NGC 1068 is “hidden” in gamma rays, as the neutrino flux exceeds the gamma-ray flux (see Figure 9). NGC 4151 is also an obscured source, with the core surrounded by heavy amounts of dust and gas, and it has not been detected in gamma rays. It has been suggested that the interaction sites of the cosmic rays within these AGN are close enough to the core that the photon density is sufficiently high and the gamma rays on the order of GeV-energy cannot escape. For details of the modeling of similar sources, see K. Murase (2022), C. Blanco et al. (2023), A. Das et al. (2024), and K. Murase et al. (2024).

Neutrino emission from these sources was also tested in a complementary analysis of Seyfert galaxies (R. Abbasi et al. 2024). The two analyses overlap in the sources examined. Nevertheless, the hypotheses tested, the neutrino data set, and the analysis techniques are different. The complementary analysis tests specific models of emission (A. Kheirandish et al. 2021) using a data set restricted to the northern sky. The conclusions from the two studies are consistent.

After the present work was unblinded, and while the manuscript was in preparation, the results of A. Neronov et al. (2024) were shared with us and subsequently published. In that work, the authors find a $\sim 3\sigma$ neutrino signal from a population of Seyfert galaxies. The population of Seyferts they considered includes NGC 4151 and NGC 3079, the second- and third-most significant neutrino sources observed in the second analysis of this work (see Table 5).

The best-fit spectra found for the two sources are $\sim E^{-3}$ and softer than the diffuse astrophysical flux obtained by IceCube studies (M. G. Aartsen et al. 2019b; R. Abbasi et al. 2022b). To resolve this discrepancy, we require either a dominant source population with a harder spectrum or a class of similar sources with some emitting at higher energies than the observed ones. Considering NGC 1068 and NGC 4151 as two sources representative of a class of neutrino emitters, we find that they

contribute $\sim 1\%$ of neutrinos to the diffuse flux but only $\sim 0.1\%$ to the total hard X-ray flux at a spectral index $\gamma = 3.0$. For the neutrino flux, we chose 15 TeV to evaluate the flux, since this is within the sensitive energy range of both sources and the diffuse flux. We also assumed a direct relation between the hard X-ray and neutrino flux and corrected for catalog completeness. This suggests underlying physical parameters besides hard X-ray flux may determine neutrino emission despite the limitations of considering a sample of only two nearby AGN. The intrinsic fluxes from obscured sources, especially CT AGN, have high uncertainties, as it is challenging to observe the emission from the core. In conclusion, growing evidence points to Seyfert galaxies as neutrino emitters. Better X-ray flux measurements of AGN especially for CT AGN may prove important to the discovery of high-energy neutrino sources.

Acknowledgments

The IceCube collaboration acknowledges significant contributions to this manuscript from J.D., S.G., and G.C.P.

The authors gratefully acknowledge the support from the following agencies and institutions: USA—U.S. National Science Foundation—Office of Polar Programs, U.S. National Science Foundation—Physics Division, U.S. National Science Foundation-EPSCoR, U.S. National Science Foundation—Office of Advanced Cyberinfrastructure, Wisconsin Alumni Research Foundation, Center for High Throughput Computing (CHTC) at the University of Wisconsin–Madison, Open Science Grid (OSG), Partnership to Advance Throughput Computing (PATH), Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS), Frontera computing project at the Texas Advanced Computing Center, U.S. Department of Energy—National Energy Research Scientific Computing Center, Particle Astrophysics Research Computing Center at the University of Maryland, Institute for Cyber-Enabled Research at Michigan State University, Astroparticle Physics Computational Facility at Marquette University, NVIDIA Corporation, and Google Cloud Platform; Belgium—Funds for Scientific Research (FRS-FNRS and FWO), FWO Odysseus and Big Science programmes, and Belgian Federal Science Policy Office (Belspo); Germany—Bundesministerium für Bildung und Forschung (BMBF), Deutsche Forschungsgemeinschaft (DFG), Helmholtz Alliance for Astroparticle Physics (HAP), Initiative and Networking Fund of the Helmholtz Association, Deutsches Elektronen Synchrotron (DESY), and High Performance Computing cluster of the RWTH Aachen; Sweden—Swedish Research Council, Swedish Polar Research Secretariat, Swedish National Infrastructure for Computing (SNIC), and Knut and Alice Wallenberg Foundation; European Union—EGI Advanced Computing for Research; Australia—Australian Research Council; Canada—Natural Sciences and Engineering Research Council of Canada, Calcul Québec, Compute Ontario, Canada Foundation for Innovation, WestGrid, and Digital Research Alliance of Canada; Denmark—Villum Fonden, Carlsberg Foundation, and European Commission; New Zealand—Marsden Fund; Japan—Japan Society for Promotion of Science (JSPS) and Institute for Global Prominent Research (IGPR) of Chiba University; Korea—National Research Foundation of Korea (NRF); Switzerland—Swiss National Science Foundation (SNSF).

Appendix

A.1. Neutrino Data Set

Table 3 shows the different configurations in which the IceCube detector operated, the livetime in days, and the number of events for each detector configuration. For the incomplete detector configurations with 40, 59, and 79 strings, the detector operated for about a year collecting data. With the complete configuration of 86 strings, there are various data-taking periods also about a year in length, each of which is called a “season.” The table includes the breakdown of the IC 86 configuration into different seasons with the livetime and the number of events in each season.

Table 3
Details of the Neutrino Data Set Used for the Analysis

Configuration	Livetime (days)	No. of Events
IC 40	2008 Apr 6–2009 May 20 (376.4)	36,900
IC 59	2009 May 20–2010 May 31 (353.6)	107,011
IC 79	2010 Jun 1–2011 May 13 (316.0)	93,133
IC 86, 2011	2011 May 13–2012 May 15 (340.1)	119,169
IC 86, 2012	2012 Apr 26–2013 May 2 (327.7)	116,715
IC 86, 2013	2013 Apr 18–2014 May 5 (355.6)	126,337
IC 86, 2014	2014 Apr 10–2015 May 18 (365.5)	129,823
IC 86, 2015	2015 Apr 24–2016 May 20 (365.3)	130,434
IC 86, 2016	2016 May 20–2017 May 18 (357.2)	126,438
IC 86, 2017	2017 May 18–2018 Jul 10 (405.6)	145,602
IC 86, 2018	2018 Jun 19–2019 Jul 17 (362.5)	129,230
IC 86, 2019	2019 Jun 28–2020 May 29 (304.7)	109,616
IC 86, all	2011 May 13–2020 May 29 (3184.2)	1,133,364

Note. The start date of some configurations is earlier than the end date of the previous season. This is because these dates denote the start date of test runs of the new processing, and the data-taking begins once the previous run ends.

A.2. Hard X-Ray Sources

The intrinsic X-ray flux is the observed flux after correcting for absorption and performing k -correction, i.e., shifting the flux such that it is in the 14–195 keV range in the source frame. The absorption arises mainly from photoelectric absorption and Compton scattering, which are both taken into account. The intrinsic X-ray flux and other parameters have been estimated in BASS using X-ray spectral analysis. Many models were tested by the authors, to find the one that is most appropriate for each source. Different models are used for the analysis of blazars and nonblazar AGN (P. Magdziarz & A. A. Zdziarski 1995). The authors computed the flux in the energy range 14–195 keV by extrapolating the fluxes in the 14–150 keV energy range, because the Swift-BAT detector response for the highest-energy range, 150–195 keV, is poor, with a signal-to-noise ratio ~ 100 –1000 times lower than those of other energy bands.

To evaluate the column density, the authors of the BASS catalog first used a simple power-law spectrum to fit the observations and then increased the number of free parameters as and when required. From the spectral analysis, if a source resulted in column density, $N_H > 10^{24} \text{ cm}^{-2}$, it is analyzed using a spheroidal torus model from M. Brightman & K. Nandra (2011) and classified as CT in the BASS catalog (C. Ricci et al. 2015).

In Figure 4 (left), we show the positions of the hard X-ray AGN from the BASS catalog. In Figure 4 (right), we can see from the y-axis that the sources in the catalog have a wide range of luminosity spanning around seven orders of magnitude. The most distant sources observed are around $z \sim 3$ and are also the most luminous, and as the distance increases, the AGN detected are more likely to be blazars with a higher luminosity and also detected in gamma rays. There are fewer nonblazar AGN observed at higher redshift in comparison, since the dust and gas in the torus prevent any light from the core from being visible at very large distances. However, for the blazars, their orientation makes the powerful jets point toward us, and due to relativistic




Figure 4. Left: sky map in equatorial coordinates that shows all the hard X-ray AGN in the catalog used to perform the analyses in this study. The color scheme of the sources shows the hard X-ray intrinsic flux (in the 14–195 keV energy range). The black dashed line represents the Galactic plane, and the star on the lower left represents the Galactic center. Right: the distribution of the hard X-ray sources in the parameter space of luminosity as a function of redshift. The color of each data point represents the column density of the sources. The sources are shown using different markers that correspond to the type of AGN, and the circled sources represent the AGN that have been detected in gamma rays and are found in the Fermi 4-FGL catalog (S. Abdollahi et al. 2020). The Galactic blazar is defined as a blazar with an additional component from the galaxy (M. J. Koss et al. 2022a).

beaming, the emission from the jet is boosted, making them visible at a far greater distance. The jet is also able to penetrate any gas or dust it encounters, even though the obscuration arises mostly from the torus whose axis aligns with the jet—and hence does not block any light from it.

A.3. Hypothesis Testing

The statistical method followed to perform the two analyses is a hypothesis test. We evaluate the test statistic (TS) using the likelihood function given in Equation (1) and defined as follows, with a larger TS indicating that the data favor the signal hypothesis more than a lower TS:

$$TS = -2 \log \left(\frac{\mathcal{L}(n_s = 0)}{\mathcal{L}(n_s = \hat{n}_s, \gamma = \hat{\gamma})} \right). \quad (\text{A1})$$

The pretrial p -value, p_{local} , is defined as the survival probability of the observed TS value compared to a distribution of TS values computed by generating background-only pseudo-experiments. The background PDF is obtained by scrambling only the R.A. of neutrino events to preserve the decl. distribution of the events. Each scrambling realization is a trial. We plot the TS values of these trials to obtain the background PDF. We then fit a χ^2_ν distribution plus a delta function at $TS = 0$ to our histogram. To compute a p -value associated with a certain TS, we find the background-only TS distribution by performing repeated searches on events with randomized R.A. values. p -values are then converted to one-sided statistical significance.

A.4. Stacked Search Methods and Additional Results

A.4.1. Catalog Completeness Correction

To calculate the catalog completeness factor, we compute the fraction of the neutrino flux expected from the entire source population due to the sources contained in the catalog.

In the analysis, we hypothesize

$$\frac{dN_\nu}{dE_\nu} \propto F_{\text{X-ray}}^{\text{int}}, \quad (\text{A2})$$

where $\frac{dN_\nu}{dE_\nu}$ is the neutrino differential energy spectrum and $F_{\text{X-ray}}^{\text{int}}$ is the intrinsic X-ray flux.

Most of the sources in the BASS catalog are at relatively low redshift ($z_{\text{mean}} = 0.16$), but for the most distant AGN, cosmological redshift plays a significant role in the observed neutrino flux. The flux is corrected for absorption and k -corrected. We measure the neutrino flux at a specific energy of 1 TeV and need to correct it to evaluate the intrinsic neutrino flux at the source. The observed neutrino differential energy spectrum is assumed to follow a power law given by

$$\frac{dN_\nu}{dE_\nu}^{\text{obs}} = \frac{dN_\nu}{dE_\nu}^{\text{source}} (1+z)^{-\gamma} \propto F_{\text{X-ray}}^{\text{int}} (1+z)^{-\gamma}. \quad (\text{A3})$$

The values for $F_{\text{X-ray}}^{\text{int}}$ and redshift z are available in the BASS catalog. Finally, by integrating the luminosity function of the source class (from L. Marcotulli et al. 2022 for blazars and from Y. Ueda et al. 2014 for nonblazar AGN), we obtain the contribution from the entire source population. The catalog completeness correction factor, or completeness factor for short, derived for each source class and spectral index are reported in Table 4. The neutrino flux upper limits obtained from the stacked search for various spectral indices are divided by these values.

Table 4
Fractions Showing the Completeness Factors of the Source Populations

Category	Completeness Fraction $\gamma = 2.0$ ($E^{-2.0}$)	Completeness Fraction $\gamma = 2.5$ ($E^{-2.5}$)	Completeness Fraction $\gamma = 3.0$ ($E^{-3.0}$)
All AGN	2.42 $^{+0.59}_{-0.59}\%$	3.17 $^{+0.72}_{-0.74}\%$	4.01 $^{+0.92}_{-0.89}\%$
Nonblazar AGN	2.31 $^{+0.62}_{-0.51}\%$	3.04 $^{+0.75}_{-0.64}\%$	3.86 $^{+0.95}_{-0.80}\%$
Blazars	11.52 $^{+24.15}_{-8.74}\%$	14.58 $^{+26.58}_{-10.86}\%$	17.97 $^{+28.74}_{-12.50}\%$
Unobscured	5.05 $^{+1.14}_{-1.57}\%$	6.49 $^{+1.32}_{-1.89}\%$	8.06 $^{+1.68}_{-2.23}\%$
Obscured	2.26 $^{+0.59}_{-0.52}\%$	2.98 $^{+0.71}_{-0.65}\%$	3.78 $^{+0.90}_{-0.81}\%$
Compton-thick	1.52 $^{+0.42}_{-0.34}\%$	2.03 $^{+0.50}_{-0.43}\%$	2.59 $^{+0.64}_{-0.54}\%$

We need the sum of the neutrino signal expectation from both the sources being tested and from the entire source class population in the observable Universe. To calculate the completeness factor, we define a quantity $S(\gamma)$ using Equation (A3), where $S(\gamma) = F_{\text{X-ray}}^{\text{int}} (1+z)^{-\gamma}$ and is proportional to the neutrino signal expectation. To find $S(\gamma)_{\text{total}}$, the sum of $S(\gamma)$ from the entire source class population, we use

$$S(\gamma)_{\text{total}} = \int_0^{4\pi} \int_0^{z_{\text{max}}} \int_{L_{\text{min}}}^{L_{\text{max}}} \frac{dV}{dzd\Omega} \times S(L, z, \gamma) \phi(L, z) dL dz d\Omega, \quad (\text{A4})$$

where $\phi(L, z)$ is the luminosity function and $\frac{dV}{dzd\Omega}$ is the comoving volume per unit redshift and solid angle. To compute this integral, we need an accurate luminosity function that captures the source density through all relevant luminosities and redshifts.

For the nonblazar AGN and their column density subclasses, we use the best-fit luminosity function from Y. Ueda et al. (2014). For their fit, those authors used X-ray surveys from multiple instruments that covered a combined energy range of 0.5–195 keV. The luminosity function is presented in the 2–10 keV range, so we use their spectral template of a cutoff power law to convert it to the 14–195 keV range. Since the luminosity function is in terms of intrinsic and k -corrected luminosity, no other adjustments need to be made.

For the blazar AGN, we use the best-fit luminosity function from L. Marcotulli et al. (2022). They tested multiple models, so we used the model that has the lowest Akaike information criterion. Those authors gave the luminosity function in terms of intrinsic and k -corrected luminosity, so no adjustments need to be made.

When integrating these luminosity functions we use the same minimum and maximum luminosities as those authors: $L_{\text{min}} = 10^{43} \text{ erg s}^{-1}$ and $L_{\text{max}} = 10^{50} \text{ erg s}^{-1}$ for the blazars and $L_{\text{min}} = 10^{41} \text{ erg s}^{-1}$ and $L_{\text{max}} = 10^{47} \text{ erg s}^{-1}$ for the nonblazars. We use the same maximum redshift of $z_{\text{max}} = 6$ for both classes.

To calculate $S(\gamma)_{\text{total}}$ for the different classes of column density, we use the nonblazar luminosity function separated into its contributions from each class. We also need the blazar contribution for each class, but the blazar luminosity function is not a function of column density. Most blazars have low column densities, as the jet is not traveling through the obscuring torus, so we add the blazar luminosity function to the unobscured nonblazar AGN luminosity function to calculate the unobscured $S(\gamma)_{\text{total}}$. Some BASS blazars fall into the obscured class (12% of total BASS blazars), either through




Figure 5. Plotted here are the inverse cumulative distributions of $S(\gamma)$ for $\gamma = 2$, for both the entire population and the BASS sources. The distributions for both all AGN and just blazars are plotted. The catalog completeness fraction is also listed along with an arrow illustrating the difference between the numerator and denominator in the fraction. At the highest fluxes, there is an overfluctuation of bright sources, which results in a higher value for sources in the catalog as compared to the total flux from all possible sources. Since the plot shows the cumulative distribution, this effect is not seen at the lower fluxes.

truly having that much obscuring material or through poor model fits. To account for this, we add 0.12 times the blazar luminosity function to the obscured nonblazar luminosity function to calculate the obscured $S(\gamma)_{\text{total}}$. The fractional blazar contributions to each class are rough estimates, but the blazar total flux is subdominant to the nonblazar AGN flux by ~ 2 orders of magnitude, so little difference is made by changing these fractional contributions.

In Figure 5, we have plotted the inverse cumulative $S(\gamma)$ for $\gamma = 2$, as a function of $S(\gamma)$. The arrows in the figure show the difference between S_{total} for just the sources in the catalog and the entire source class. The ratio of these two values gives our catalog completeness fraction. Figure 5 outlines this for all AGN, which is dominated by nonblazar AGN, and for blazars. The curves for the BASS sources closely match the curves for the entire population up until the sensitivity of the catalog is reached.

To find the error of $S(\gamma)_{\text{total}}$, we use a bootstrapping method along with the reported errors for each fit parameter from both luminosity functions. We randomly sample the parameters' values from their error PDFs, assuming they are Gaussian and uncorrelated. They are most likely correlated, but we do not have that information and the assumption of no correlation gives a larger and more conservative error. Then, we calculate $S(\gamma)_{\text{total}}$ for many samples and find the 16th and 84th percentiles of the distribution of $S(\gamma)_{\text{total}}$ to determine the 1σ range of the completeness factors. The errors are shown in Table 4 along with the completeness factors for three different γ 's.

A.4.2. Trial Factor Correction

We have tested seven different hypotheses in the stacking analysis (see Section 3.1) using different subclasses of AGN with a significant overlap in the source populations. To evaluate the correlation between the different AGN samples used to test each hypothesis, we performed a large number of trials, $\mathcal{O}(10^5)$, since a simple analytical function as was used




Figure 6. The cumulative distribution function of the minimum of the p -values from each AGN sample resulting from background trials, and the pretrial p -values of the samples in the unblinded analysis. “All-flux” and “all-equal” refer to the samples containing all 836 AGN tested using hard X-ray flux as weights and equal weights, respectively.

previously for the individual source search is not sufficient in this case. From each background trial, we computed the p -values for each of the seven AGN samples and found the minimum p -value. Figure 6 shows the distribution of the minimum p -values. The corresponding value of p_{local} gives the posttrial p -value. For the AGN samples tested, the pretrial p -values and pretrial significances are reported in Table 1.

A.4.3. Energy Range Evaluation

As the sensitivity of the IceCube detector is not constant for all neutrinos and varies with energy, we evaluate the energy range most relevant to the analyses. We follow two methods for the two analyses. This range reflects most of the neutrino events that contribute to the TS.

Since we find flux upper limits in the stacking analysis and the results do not show a high significance, we utilize pseudo-signal events from Monte Carlo instead of experimental data to determine the central 90% energy range. To evaluate the upper bound of the energy range, we performed many trials by injecting signals over the background, with a fixed value for the lower-energy bound and varying the upper energy bound. Next, we compute the sensitivity for each energy range and plot each one a function of the energy. The lowest value of the higher-bound energy for which the evaluated sensitivity is below 1.05 times the total sensitivity flux evaluated over an energy range of zero to infinity gives us the most sensitive higher-bound energy value. To evaluate the lower bound of the energy range, we performed many trials by injecting signals over the background with a fixed value of higher energy bound and varying the lower-energy bound to find the sensitivity for each energy range and plot them as a function of the energy. The lowest value of the varying lower-bound energy for which the evaluated sensitivity is above 1.05 times the total sensitivity flux evaluated over an energy range of zero to infinity gives us the most sensitive higher-bound energy value.

The most sensitive energy range is not expected to vary significantly with the different AGN samples, but it will change for different spectral indices. Therefore, we evaluate the energy range using all AGN and use the values for the other samples as well. The evaluated energy range where the analysis is most

sensitive for $\gamma = 2.0$ is 6 TeV–10 PeV, that for $\gamma = 2.5$ is 2.5 TeV–1.1 PeV, and that for $\gamma = 3.0$ is 0.5 TeV–0.1 PeV.

For the individual source search, the relevant energy ranges were determined using the experimental data collected from muon tracks over 12 yr. The resultant fluxes for the two sources, NGC 1068 and NGC 4151, were obtained assuming a power-law spectrum. Even though these fluxes extend across the entire energy range covered by the neutrino data set, the most relevant energy range corresponds to that which contributes 68% toward the TS obtained for each source.

In the case of NGC 1068, the energy range is 0.7–15 TeV. This is slightly different from the value reported for NGC 1068 in R. Abbasi et al. (2022g), where the energy range is

1.5–15 TeV. This discrepancy is due to the different data sets being used for the two analyses and their different effective areas. While in this analysis, we used the Point Source Tracks sample, which includes tracks from the whole sky (for details, see M. G. Aartsen et al. 2020c), the previous analyses used the Northern Tracks sample, which focuses on muon tracks from the northern sky (for details, see M. G. Aartsen et al. 2019a).

A.4.4. Stacked Search: Additional Results

In addition to the flux upper limits evaluated for index $\gamma = 3.0$, we computed upper limits for indices $\gamma = 2.5$ and $\gamma = 2.0$ as well. These flux upper limits are given in Table 1 and shown in Figure 7.




Figure 7. Panels (a) and (c) show the 90% CL flux upper limits for all AGN, AGN subclasses of blazars, and nonblazars, and panels (b) and (d) show the upper limits for the unobscured, obscured, and Compton-thick sources. The flux is for power-law spectra with a spectral index of 2.0 (top row) and 2.5 (bottom row). These flux values are derived using the completeness correction described in Appendix A.4.1. For reference, the plot shows the all-sky diffuse flux from muon tracks (R. Abbasi et al. 2022b) in light blue dotted lines and that from cascades (M. G. Aartsen et al. 2019b) in purple.

A.5. Individual Source Search: Source Selection

To evaluate the “figure of merit” that formed the basis of the selection of 43 sources to be analyzed more closely in the individual source search in Section 3.2, we first computed the neutrino flux sensitivity, defined as the required signal, in terms

of flux or number of neutrinos, that yields a p -value smaller than 0.5 in 90% of the trials, as a function of the decl. for spectral index $\gamma = 2.5$. The sensitivity allows the IceCube detector to observe a source at different declinations and is shown in Figure 8 (left). The list’s final selection of 43 sources and their distribution in the sky is shown in Figure 8 (right).





Figure 8. Left: neutrino sensitivity flux (in blue) and the flux upper limits (in orange) at normalization energy 1 TeV and power-law spectral index $\gamma = 2.5$ as a function of the sine of decl. Right: sky map showing the 43 sources from which we search for the emission of neutrinos, individually. Most sources are in the Northern Hemisphere, as seen in Figure 8 (right).

A.6. Individual Source Search: Additional Results

We calculated the p_{local} for each of the sources, which is reported in Table 5. For the two most significant sources, we show their electromagnetic spectra observed by different telescopes overlaid with the neutrino spectrum obtained from this analysis in Figure 9.

Other than the two sources found with the highest significance, we derived the flux upper limits for each source at three different power-law indices: $\gamma = 2.0, 2.5$, and 3.0 . See the flux upper limits in Table 5.

Table 5
Summary of Results from the Individual Source Search

Type	Source	α (deg)	δ (deg)	$\hat{n}_s$	$\hat{\gamma}$	p_{local}	X-Ray Flux	$\phi_{90\%}^{2.0}$	$\phi_{90\%}^{2.5}$	$\phi_{90\%}^{3.0}$
Seyfert	NGC 1068	40.67	−0.01	63.18	3.1	1.27×10^{-6}	206.0	...	...	...
	NGC 4151	182.64	39.41	43.67	2.83	3.99×10^{-5}	525.7	...	...	...
	NGC 3079	150.49	55.68	29.53	4.0	0.003	112.6	13.84	77.91	203.50
	MCG 4-48-2	307.15	25.73	21.74	3.75	0.060	63.9	6.44	49.50	150.52
	NGC 4992	197.27	11.63	17.41	2.86	0.086	66.8	4.89	46.79	166.27
	3C 111	64.59	38.03	18.22	3.45	0.108	118.7	7.43	45.91	127.20
	NGC 1275	49.95	41.51	16.13	3.55	0.117	79.2	7.42	45.01	123.17
	Q0241+622	41.24	62.47	18.4	3.84	0.129	89.6	10.17	49.92	120.86
	Cygnus A	299.87	40.73	1.84	1.38	0.141	137.9	5.98	30.95	114.64
	NGC 1194	45.95	−1.1	12.74	3.75	0.229	114.5	2.69	35.22	144.95
	NGC 5548	214.5	25.14	11.18	4.0	0.246	75.9	4.86	33.63	103.11
	Z164-19	221.4	27.03	8.43	4.0	0.252	250.1	4.74	33.90	102.57
	IRAS 05589+2828	90.54	28.47	7.58	4.0	0.268	62.4	4.89	31.53	90.33
	4C 50.55	321.16	50.97	6.41	3.41	0.329	217.3	5.56	29.49	79.88
	Mrk 348	12.2	31.96	5.27	4.0	0.356	159.2	4.38	25.92	81.12
	NGC 1142	43.8	−0.18	7.43	3.18	0.359	123.7	2.26	28.56	113.46
	Mrk 417	162.38	22.96	1.72	2.44	0.407	50.0	3.46	25.03	106.33
	NGC 4102	181.6	52.71	4.05	3.75	0.441	71.8	4.91	25.60	65.94
	Mrk 1040	37.06	31.31	3.51	3.49	0.460	61.1	3.53	21.35	67.19
	Mrk 110	141.3	52.29	0.41	1.43	0.466	57.1	4.79	24.10	64.50
	3C 120	68.3	5.35	4.69	3.82	0.472	93.4	2.19	21.49	78.00
	NGC 5252	204.57	4.54	0.73	2.24	0.493	106.3	2.01	22.24	84.06
	NGC 7469	345.82	8.87	0.17	2.58	0.534	64.3	2.16	20.42	73.51
	IRAS 05078+1626	77.69	16.5	0.0	0.0	1.0	92.3	2.90	22.94	71.02
	Mrk 1210	121.02	5.11	0.0	0.0	1.0	61.5	1.96	20.87	85.90
	NGC 2110	88.05	−7.46	0.0	0.0	1.0	317.5	3.51	75.21	608.59
	NGC 7682	352.27	3.53	0.0	0.0	1.0	55.8	1.98	22.55	85.59
	Ark 120	79.05	−0.15	0.0	0.0	1.0	68.0	1.80	22.15	91.29
	3C 382	278.76	32.7	0.0	0.0	1.0	75.0	3.52	21.65	64.13
	2MASX J20145928+2523010	303.75	25.38	0.0	0.0	1.0	73.6	3.12	21.15	67.60
	NGC 3516	166.7	72.57	0.0	0.0	1.0	113.9	8.01	36.84	93.62
	NGC 6240	253.25	2.4	0.0	0.0	1.0	348.2	1.89	21.86	85.01
	NGC 3227	155.88	19.87	0.0	0.0	1.0	106.6	2.90	21.62	69.19
	NGC 5506	213.31	−3.21	0.0	0.0	1.0	234.7	1.87	23.38	94.73
	UGC 3374	88.72	46.44	0.0	0.0	1.0	132.9	4.14	22.71	60.82
	UGC 11910	331.76	10.23	0.0	0.0	1.0	60.8	2.34	21.63	79.23
	NGC 4388	186.44	12.66	0.0	0.0	1.0	323.0	2.55	22.78	78.43
	Mrk 3	93.9	71.04	0.0	0.0	1.0	279.4	7.98	37.09	94.26
	LEDA 168563	73.02	49.55	0.0	0.0	1.0	58.7	4.28	22.35	62.87
	3C 445	335.96	−2.1	0.0	0.0	1.0	55.2	1.77	22.88	96.98
FSRQ	3C 454.3	343.49	16.15	3.71	2.01	0.150	124.9	4.54	35.05	114.93
	3C 273	187.28	2.05	3.51	2.20	0.230	434.7	2.88	33.35	133.67
BL Lac	Mrk 421	166.11	38.21	0.0	0.0	1.0	182.9	3.86	22.84	61.88

Note. The table contains the results from the individual source search. It shows the sources, their type, their positions in equatorial coordinates (J2000), the best-fit number of astrophysical neutrino events $\hat{n}_s$, the best-fit spectral index $\hat{\gamma}$, the pretrial p -value, and the 90% CL flux upper limit per flavor for $\gamma = 2.0, 2.5$, and 3.0 at a normalization energy of 1 TeV given by $\phi_{90\%}^{2.0}$, $\phi_{90\%}^{2.5}$, and $\phi_{90\%}^{3.0}$ in units of $10^{-13} \text{ TeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$. X-ray flux in the table is the intrinsic hard X-ray flux in the energy range 14–195 keV and in units of $10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$. The source type is obtained from the BASS DR-2 catalog and differs from the 105 month BAT survey classification for some sources; e.g., NGC 1275, Mrk 348, and 3C 120 are classified as blazars instead of Seyferts in the 105 month BAT survey.




Figure 9. The spectral energy distribution of the electromagnetic (EM) emissions from NGC 4151 and NGC 1068 and their neutrino spectra obtained from this analysis. Electromagnetic data and upper limits are taken from <https://tools.ssdc.asi.it/SED/>.

ORCID iDs

- R. Abbasi <https://orcid.org/0000-0001-6141-4205>
M. Ackermann <https://orcid.org/0000-0001-8952-588X>
S. K. Agarwalla <https://orcid.org/0000-0002-9714-8866>
J. A. Aguilar <https://orcid.org/0000-0003-2252-9514>
M. Ahlers <https://orcid.org/0000-0003-0709-5631>
J.M. Alameddine <https://orcid.org/0000-0002-9534-9189>
K. Andeen <https://orcid.org/0000-0001-9394-0007>
C. Argüelles <https://orcid.org/0000-0003-4186-4182>
S. N. Axani <https://orcid.org/0000-0001-8866-3826>
X. Bai <https://orcid.org/0000-0002-1827-9121>
A. Balagopal V. <https://orcid.org/0000-0001-5367-8876>
S. W. Barwick <https://orcid.org/0000-0003-2050-6714>
V. Basu <https://orcid.org/0000-0002-9528-2009>
J. J. Beatty <https://orcid.org/0000-0003-0481-4952>
J. Becker Tjus <https://orcid.org/0000-0002-1748-7367>
J. Beise <https://orcid.org/0000-0002-7448-4189>
C. Bellenghi <https://orcid.org/0000-0001-8525-7515>
S. BenZvi <https://orcid.org/0000-0001-5537-4710>
E. Bernardini <https://orcid.org/0000-0003-3108-1141>
E. Blaufuss <https://orcid.org/0000-0001-5450-1757>
L. Bloom <https://orcid.org/0009-0005-9938-3164>
S. Blot <https://orcid.org/0000-0003-1089-3001>
J. Y. Book Motzkin <https://orcid.org/0000-0001-6687-5959>
C. Boscolo Meneguolo <https://orcid.org/0000-0001-8325-4329>
S. Böser <https://orcid.org/0000-0002-5918-4890>
O. Botner <https://orcid.org/0000-0001-8588-7306>
J. Böttcher <https://orcid.org/0000-0002-3387-4236>
B. Brinson <https://orcid.org/0000-0001-9128-1159>
M. A. Campana <https://orcid.org/0000-0003-4162-5739>
J. Carpio <https://orcid.org/0000-0003-0667-6557>
S. Chattopadhyay <https://orcid.org/0009-0006-1352-2248>
D. Chirkin <https://orcid.org/0000-0003-4911-1345>
B. A. Clark <https://orcid.org/0000-0003-4089-2245>
A. Coleman <https://orcid.org/0000-0003-1510-1712>
J. M. Conrad <https://orcid.org/0000-0002-6393-0438>
P. Coppin <https://orcid.org/0000-0001-6869-1280>
P. Correa <https://orcid.org/0000-0002-1158-6735>
D. F. Cowen <https://orcid.org/0000-0003-4738-0787>
P. Dave <https://orcid.org/0000-0002-3879-5115>
C. De Clercq <https://orcid.org/0000-0001-5266-7059>
J. J. DeLaunay <https://orcid.org/0000-0001-5229-1995>
D. Delgado <https://orcid.org/0000-0002-4306-8828>
A. Desai <https://orcid.org/0000-0001-7405-9994>
P. Desiati <https://orcid.org/0000-0001-9768-1858>
K. D. de Vries <https://orcid.org/0000-0002-9842-4068>
G. de Wasseige <https://orcid.org/0000-0002-1010-5100>
T. DeYoung <https://orcid.org/0000-0003-4873-3783>
A. Diaz <https://orcid.org/0000-0001-7206-8336>
J. C. Díaz-Vélez <https://orcid.org/0000-0002-0087-0693>
H. Dujmovic <https://orcid.org/0000-0003-1891-0718>
M. A. DuVernois <https://orcid.org/0000-0002-2987-9691>
P. Eller <https://orcid.org/0000-0001-6354-5209>
D. Elsässer <https://orcid.org/0000-0001-6796-3205>
H. Erpenbeck <https://orcid.org/0000-0001-6319-2108>
A. R. Fazely <https://orcid.org/0000-0002-6907-8020>
A. Fedynitch <https://orcid.org/0000-0003-2837-3477>
C. Finley <https://orcid.org/0000-0003-3350-390X>
L. Fischer <https://orcid.org/0000-0002-7645-8048>
D. Fox <https://orcid.org/0000-0002-3714-672X>
A. Franckowiak <https://orcid.org/0000-0002-5605-2219>
P. Fürst <https://orcid.org/0000-0002-7951-8042>
J. Gallagher <https://orcid.org/0000-0001-8608-0408>
E. Ganster <https://orcid.org/0000-0003-4393-6944>
A. Garcia <https://orcid.org/0000-0002-8186-2459>
A. Ghadimi <https://orcid.org/0000-0002-6350-6485>
T. Glüsenskamp <https://orcid.org/0000-0002-2268-9297>
S. J. Gray <https://orcid.org/0000-0003-2907-8306>
S. Griffin <https://orcid.org/0000-0002-0779-9623>
S. Griswold <https://orcid.org/0000-0002-7321-7513>
K. M. Groth <https://orcid.org/0000-0002-1581-9049>
P. Gutjahr <https://orcid.org/0000-0001-7980-7285>
C. Haack <https://orcid.org/0000-0003-3932-2448>
A. Hallgren <https://orcid.org/0000-0001-7751-4489>
L. Halve <https://orcid.org/0000-0003-2237-6714>
F. Halzen <https://orcid.org/0000-0001-6224-2417>
H. Hamdaoui <https://orcid.org/0000-0001-5709-2100>
A. Haungs <https://orcid.org/0000-0002-9638-7574>
K. Helbing <https://orcid.org/0000-0003-2072-4172>
J. Hellrung <https://orcid.org/0009-0006-7300-8961>
N. Heyer <https://orcid.org/0000-0001-9036-8623>
C. Hill <https://orcid.org/0000-0003-0647-9174>
S. Hori <https://orcid.org/0009-0007-2644-5955>
M. Hostert <https://orcid.org/0000-0002-9584-8877>
W. Hou <https://orcid.org/0000-0003-3422-7185>
T. Huber <https://orcid.org/0000-0002-6515-1673>
K. Hultqvist <https://orcid.org/0000-0003-0602-9472>
M. Hünnefeld <https://orcid.org/0000-0002-2827-6522>
W. Iwakiri <https://orcid.org/0000-0002-0207-9010>
G. S. Japaridze <https://orcid.org/0000-0002-7000-5291>
M. Jeong <https://orcid.org/0000-0003-2420-6639>
M. Jin <https://orcid.org/0000-0003-0487-5595>
B. J. P. Jones <https://orcid.org/0000-0003-3400-8986>
D. Kang <https://orcid.org/0000-0002-5149-9767>
W. Kang <https://orcid.org/0000-0003-3980-3778>
A. Kappes <https://orcid.org/0000-0003-1315-3711>
T. Karg <https://orcid.org/0000-0003-3251-2126>
M. Karl <https://orcid.org/0000-0003-2475-8951>
A. Karle <https://orcid.org/0000-0001-9889-5161>
U. Katz <https://orcid.org/0000-0002-7063-4418>
M. Kauer <https://orcid.org/0000-0003-1830-9076>
J. L. Kelley <https://orcid.org/0000-0002-0846-4542>
A. Khatee Zathul <https://orcid.org/0000-0002-8735-8579>

- A. Kheirandish  <https://orcid.org/0000-0001-7074-0539>
J. Koryluk  <https://orcid.org/0000-0003-0264-3133>
S. R. Klein  <https://orcid.org/0000-0003-2841-6553>
A. Kochocki  <https://orcid.org/0000-0003-3782-0128>
R. Koirala  <https://orcid.org/0000-0002-7735-7169>
H. Kolanoski  <https://orcid.org/0000-0003-0435-2524>
T. Kontrimas  <https://orcid.org/0000-0001-8585-0933>
C. Kopper  <https://orcid.org/0000-0001-6288-7637>
D. J. Koskinen  <https://orcid.org/0000-0002-0514-5917>
P. Koundal  <https://orcid.org/0000-0002-5917-5230>
M. Kovacevich  <https://orcid.org/0000-0002-5019-5745>
M. Kowalski  <https://orcid.org/0000-0001-8594-8666>
K. Kruiswijk  <https://orcid.org/0009-0002-9261-0537>
A. Kumar  <https://orcid.org/0000-0002-8367-8401>
N. Kurahashi  <https://orcid.org/0000-0003-1047-8094>
N. Lad  <https://orcid.org/0000-0001-9302-5140>
C. Lagunas Gualda  <https://orcid.org/0000-0002-9040-7191>
M. Lamoureux  <https://orcid.org/0000-0002-8860-5826>
M. J. Larson  <https://orcid.org/0000-0002-6996-1155>
F. Lauber  <https://orcid.org/0000-0001-5648-5930>
J. P. Lazar  <https://orcid.org/0000-0003-0928-5025>
J. W. Lee  <https://orcid.org/0000-0001-5681-4941>
K. Leonard DeHolton  <https://orcid.org/0000-0002-8795-0601>
A. Leszczyńska  <https://orcid.org/0000-0003-0935-6313>
J. Liao  <https://orcid.org/0009-0008-8086-586X>
M. Lincetto  <https://orcid.org/0000-0002-1460-3369>
L. Lu  <https://orcid.org/0000-0003-3175-7770>
F. Lucarelli  <https://orcid.org/0000-0002-9558-8788>
W. Luszczak  <https://orcid.org/0000-0003-3085-0674>
Y. Lyu  <https://orcid.org/0000-0002-2333-4383>
J. Madsen  <https://orcid.org/0000-0003-2415-9959>
E. Magnus  <https://orcid.org/0009-0008-8111-1154>
E. Manao  <https://orcid.org/0009-0002-6197-8574>
S. Mancina  <https://orcid.org/0009-0003-9879-3896>
I. C. Mariş  <https://orcid.org/0000-0002-5771-1124>
S. Marka  <https://orcid.org/0000-0002-3957-1324>
Z. Marka  <https://orcid.org/0000-0003-1306-5260>
R. Maruyama  <https://orcid.org/0000-0003-2794-512X>
F. Mayhew  <https://orcid.org/0000-0001-7609-403X>
F. McNally  <https://orcid.org/0000-0002-0785-2244>
K. Meagher  <https://orcid.org/0000-0003-3967-1533>
M. Meier  <https://orcid.org/0000-0002-9483-9450>
L. Merten  <https://orcid.org/0000-0003-1332-9895>
T. Montaruli  <https://orcid.org/0000-0001-5014-2152>
R. W. Moore  <https://orcid.org/0000-0003-4160-4700>
M. Moulai  <https://orcid.org/0000-0001-7909-5812>
T. Mukherjee  <https://orcid.org/0000-0002-0962-4878>
R. Naab  <https://orcid.org/0000-0003-2512-466X>
R. Nagai  <https://orcid.org/0000-0001-7503-2777>
J. Necker  <https://orcid.org/0000-0003-0280-7484>
L. Neste  <https://orcid.org/0000-0002-4829-3469>
H. Niederhausen  <https://orcid.org/0000-0002-9566-4904>
M. U. Nisa  <https://orcid.org/0000-0002-6859-3944>
K. Noda  <https://orcid.org/0000-0003-1397-6478>
A. Obertacke Pollmann  <https://orcid.org/0000-0002-2492-043X>
V. O'Dell  <https://orcid.org/0000-0003-0903-543X>
B. Oeyen  <https://orcid.org/0000-0003-2940-3164>
E. O'Sullivan  <https://orcid.org/0000-0003-1882-8802>
H. Pandya  <https://orcid.org/0000-0002-6138-4808>
N. Park  <https://orcid.org/0000-0002-4282-736X>
E. N. Paudel  <https://orcid.org/0000-0001-9276-7994>
L. Paul  <https://orcid.org/0000-0003-4007-2829>
C. Pérez de los Heros  <https://orcid.org/0000-0002-2084-5866>
S. Philippen  <https://orcid.org/0000-0002-0276-0092>
A. Pizzuto  <https://orcid.org/0000-0002-8466-8168>
M. Plum  <https://orcid.org/0000-0001-8691-242X>
B. Pries  <https://orcid.org/0000-0003-4811-9863>
G. C. Privon  <https://orcid.org/0000-0003-3474-1125>
C. Raab  <https://orcid.org/0000-0001-9921-2668>
A. Rehman  <https://orcid.org/0000-0001-7616-5790>
E. Resconi  <https://orcid.org/0000-0003-0705-2770>
W. Rhode  <https://orcid.org/0000-0003-2636-5000>
B. Riedel <https://orcid.org/0000-0002-9524-8943>
M. Rongen <https://orcid.org/0000-0002-7057-1007>
A. Rosted <https://orcid.org/0000-0003-2410-400X>
C. Rott <https://orcid.org/0000-0002-6958-6033>
T. Ruhe <https://orcid.org/0000-0002-4080-9563>
I. Safa <https://orcid.org/0000-0001-8737-6825>
D. Salazar-Gallegos <https://orcid.org/0000-0002-9312-9684>
A. Sandrock <https://orcid.org/0000-0002-6779-1172>
M. Santander <https://orcid.org/0000-0001-7297-8217>
S. Sarkar <https://orcid.org/0000-0002-1206-4330>
S. Sarkar <https://orcid.org/0000-0002-3542-858X>
H. Schieler <https://orcid.org/0000-0002-2637-4778>
S. Schindler <https://orcid.org/0000-0001-5507-8890>
F. Schlüter <https://orcid.org/0000-0002-5545-4363>
J. Schneider <https://orcid.org/0000-0001-7752-5700>
F. G. Schröder <https://orcid.org/0000-0001-8495-7210>
L. Schumacher <https://orcid.org/0000-0001-8945-6722>
S. Sclafani <https://orcid.org/0000-0001-9446-1219>
M. Seikh <https://orcid.org/0000-0002-4464-7354>
S. Seunarine <https://orcid.org/0000-0003-3272-6896>
P. Sevlé Myhr <https://orcid.org/0009-0005-9103-4410>
N. Shimizu <https://orcid.org/0000-0001-6857-1772>
M. Silva <https://orcid.org/0000-0001-6940-8184>
B. Skrzypek <https://orcid.org/0000-0002-0910-1057>
B. Smithers <https://orcid.org/0000-0003-1273-985X>
D. Soldin <https://orcid.org/0000-0003-3005-7879>
P. Soldin <https://orcid.org/0000-0003-1761-2495>
G. Sommani <https://orcid.org/0000-0002-0094-826X>
G. M. Spiczak <https://orcid.org/0000-0002-0030-0519>
C. Spiering <https://orcid.org/0000-0001-7372-0074>
T. Stezelberger <https://orcid.org/0000-0003-2676-9574>
T. Stuttard <https://orcid.org/0000-0001-7944-279X>
G. W. Sullivan <https://orcid.org/0000-0002-2585-2352>
I. Taboada <https://orcid.org/0000-0003-3509-3457>
S. Ter-Antonyan <https://orcid.org/0000-0002-5788-1369>
W. G. Thompson <https://orcid.org/0000-0003-2988-7998>
J. Thwaites <https://orcid.org/0000-0001-9179-3760>
K. Tollefson <https://orcid.org/0000-0001-9725-1479>
S. Toscano <https://orcid.org/0000-0002-1860-2240>
M. A. Unland Elorrieta <https://orcid.org/0000-0002-6124-3255>
A. K. Upadhyay <https://orcid.org/0000-0003-1957-2626>
N. Valtonen-Mattila <https://orcid.org/0000-0002-1830-098X>
J. Vandenbroucke <https://orcid.org/0000-0002-9867-6548>
N. van Eijndhoven <https://orcid.org/0000-0001-5558-3328>
J. van Santen <https://orcid.org/0000-0002-2412-9728>
S. Verpoest <https://orcid.org/0000-0002-3031-3206>
A. Wang <https://orcid.org/0009-0006-9420-2667>

C. Weaver  <https://orcid.org/0000-0003-2385-2559>
 C. Wendt  <https://orcid.org/0000-0001-8076-8877>
 N. Whitehorn  <https://orcid.org/0000-0002-3157-0407>
 C. H. Wiebusch  <https://orcid.org/0000-0002-6418-3008>
 L. Witthaus  <https://orcid.org/0009-0000-0666-3671>
 M. Wolf  <https://orcid.org/0000-0001-9991-3923>
 S. Yoshida  <https://orcid.org/0000-0003-2480-5105>
 S. Yu  <https://orcid.org/0000-0003-4811-9863>
 T. Yuan  <https://orcid.org/0000-0002-7041-5872>
 P. Zhelnin  <https://orcid.org/0000-0003-1019-8375>

References

- Aartsen, M. G., Abbasi, R., Abdou, Y., et al. 2013, *Sci*, **342**, 1242856
 Aartsen, M. G., Abbasi, R., Ackermann, M., et al. 2014a, *JInst*, **9**, P03009
 Aartsen, M. G., Abraham, K., Ackermann, M., et al. 2016, *Aph*, **78**, 1
 Aartsen, M. G., Abraham, K., Ackermann, M., et al. 2017a, *ApJ*, **835**, 45
 Aartsen, M. G., Abraham, K., Ackermann, M., et al. 2017b, *ApJ*, **835**, 151
 Aartsen, M. G., Ackermann, M., Adams, J., et al. 2014b, *PhRvL*, **113**, 101101
 Aartsen, M. G., Ackermann, M., Adams, J., et al. 2017c, *JINST*, **12**, P03012
 Aartsen, M. G., Ackermann, M., Adams, J., et al. 2018a, *Sci*, **361**, 147
 Aartsen, M. G., Ackermann, M., Adams, J., et al. 2018b, *Sci*, **361**, eaat1378
 Aartsen, M. G., Ackermann, M., Adams, J., et al. 2019a, *EPJC*, **79**, 234
 Aartsen, M. G., Ackermann, M., Adams, J., et al. 2019b, *ApJ*, **886**, 12
 Aartsen, M. G., Ackermann, M., Adams, J., et al. 2020a, *PhRvL*, **125**, 121104
 Aartsen, M. G., Ackermann, M., Adams, J., et al. 2020b, *ApJ*, **898**, 117
 Aartsen, M. G., Ackermann, M., Adams, J., et al. 2020c, *PhRvL*, **124**, 051103
 Aartsen, M. G., Ackermann, M., Adams, J., et al. 2021a, *PhRvD*, **104**, 022002
 Abbasi, R., Ackermann, M., Adams, J., et al. 2021b, *JInst*, **16**, P08034
 Abbasi, R., Ackermann, M., Adams, J., et al. 2022a, *EPJC*, **82**, 1031
 Abbasi, R., Ackermann, M., Adams, J., et al. 2022b, *ApJ*, **928**, 50
 Abbasi, R., Ackermann, M., Adams, J., et al. 2022c, *ApJ*, **926**, 59
 Abbasi, R., Ackermann, M., Adams, J., et al. 2022d, *ApJL*, **930**, L24
 Abbasi, R., Ackermann, M., Adams, J., et al. 2022e, *ApJL*, **938**, L11
 Abbasi, R., Ackermann, M., Adams, J., et al. 2022f, *ApJ*, **939**, 116
 Abbasi, R., Ackermann, M., Adams, J., et al. 2022g, *Sci*, **378**, 538
 Abbasi, R., Ackermann, M., Adams, J., et al. 2022h, *ApJ*, **938**, 38
 Abbasi, R., Ackermann, M., Adams, J., et al. 2022i, *PhRvD*, **106**, 022005
 Abbasi, R., Ackermann, M., Adams, J., et al. 2023a, *ApJL*, **949**, L12
 Abbasi, R., Ackermann, M., Adams, J., et al. 2023b, *Sci*, **380**, 1338
 Abbasi, R., Ackermann, M., Adams, J., et al. 2024, arXiv:2406.07601
 Abdollahi, S., Acero, F., Ackermann, M., et al. 2020, *ApJS*, **247**, 33
 Achterberg, A., Ackermann, M., Adams, J., et al. 2006, *Aph*, **26**, 282
 Antonucci, R. 1993, *ARA&A*, **31**, 473
 Baumgartner, W. H., Tueller, J., Markwardt, C. B., et al. 2013, *ApJS*, **207**, 19
 Bentz, M. C., Williams, P. R., & Treu, T. 2022, *ApJ*, **934**, 168
 Blanco, C., Hooper, D., Linden, T., & Pinetti, E. 2023, arXiv:2307.03259
 Braun, J., Dumm, J., De Palma, F., et al. 2008, *Aph*, **29**, 299
 Brightman, M., & Nandra, K. 2011, *MNRAS*, **413**, 1206
 Das, A., Zhang, B. T., & Murase, K. 2024, *ApJ*, **972**, 44
 Edelson, R. A., Alexander, T., Crenshaw, D. M., et al. 1996, *ApJ*, **470**, 364
 Gao, S., Fedynitch, A., Winter, W., & Pohl, M. 2019, *NatAs*, **3**, 88
 Gao, S., Pohl, M., & Winter, W. 2017, *ApJ*, **843**, 109
 Halzen, F., & Zas, E. 1997, *ApJ*, **488**, 669
 Huber, M. 2019, *ICRC (Madison, WI)*, **36**, 916
 Keivani, A., Murase, K., Petropoulou, M., et al. 2018, *ApJ*, **864**, 84
 Kheirandish, A., Murase, K., & Kimura, S. S. 2021, *ApJ*, **922**, 45
 Königl, A. 2006, *MmSAI*, **77**, 598
 Koss, M., Trakhtenbrot, B., Ricci, C., et al. 2017, *ApJ*, **850**, 74
 Koss, M. J., Ricci, C., Trakhtenbrot, B., et al. 2022a, *ApJS*, **261**, 2
 Koss, M. J., Trakhtenbrot, B., Ricci, C., et al. 2022b, *ApJS*, **261**, 1
 Lovelace, R. 1976, *Natur*, **262**, 649
 Lusso, E., Comastri, A., Simmons, B. D., et al. 2012, *MNRAS*, **425**, 623
 Magdziarz, P., & Zdziarski, A. A. 1995, *MNRAS*, **273**, 837
 Mannheim, K. 1995, *Aph*, **3**, 295
 Marcotulli, L., Ajello, M., Urry, C. M., et al. 2022, *ApJ*, **940**, 77
 Mbarek, R., & Caprioli, D. 2019, *ApJ*, **886**, 8
 Mundell, C. G., Pedlar, A., Shone, D. L., & Robinson, A. 1999, *MNRAS*, **304**, 481
 Murase, K. 2022, *ApJL*, **941**, L17
 Murase, K., Karwin, C. M., Kimura, S. S., Ajello, M., & Buson, S. 2024, *ApJL*, **961**, L34
 Murase, K., & Stecker, F. W. 2023, in *The Encyclopedia of Cosmology*, ed. F. W. Stecker, Vol. 2 (Singapore: World Scientific)
 Neronov, A., Savchenko, D., & Semikoz, D. V. 2024, *PhRvL*, **132**, 101002
 Padovani, P. 2023, arXiv:2302.12540
 Petropoulou, M., Murase, K., Santander, M., et al. 2020, *ApJ*, **891**, 115
 Ricci, C., Trakhtenbrot, B., Koss, M. J., et al. 2017, *ApJS*, **233**, 17
 Ricci, C., Ueda, Y., Koss, M. J., et al. 2015, *ApJL*, **815**, L13
 Rodrigues, X., Karl, M., Padovani, P., et al. 2024, *A&A*, **689**, A147
 Stecker, F. W., Done, C., Salamon, M. H., & Sommers, P. 1991, *PhRvL*, **66**, 2697
 Tikhonov, N. A., & Galazutdinova, O. A. 2021, *AstBu*, **76**, 255
 Ueda, Y., Akiyama, M., Hasinger, G., Miyaji, T., & Watson, M. G. 2014, *ApJ*, **786**, 104
 Urry, C. M., & Padovani, P. 1995, *PASP*, **107**, 803
 Venturi, G., Cresci, G., Marconi, A., et al. 2021, *A&A*, **648**, A17
 Wang, J.-M., Songsheng, Y.-Y., Li, Y.-R., Du, P., & Yu, Z. 2020, *MNRAS*, **497**, 1020
 Williams, D. R. A., Baldi, R. D., McHardy, I. M., et al. 2020, *MNRAS*, **495**, 3079