



## BRIEF COMMUNICATION

# Trends in gastrointestinal infections before and during non-pharmaceutical interventions in Korea in comparison with the United States

Soyeoun Kim<sup>1</sup>, Jinhyun Kim<sup>2</sup>, Bo Youl Choi<sup>3</sup>, Boyoung Park<sup>3</sup><sup>1</sup>Department of Health Sciences, Hanyang University College of Medicine, Seoul, Korea; <sup>2</sup>Economics & Business Economics, University of Amsterdam, Amsterdam, The Netherlands; <sup>3</sup>Department of Preventive Medicine, Hanyang University College of Medicine, Seoul, Korea

**OBJECTIVES:** This study examined how trends in the weekly frequencies of gastrointestinal infectious diseases changed before and during the coronavirus disease 2019 (COVID-19) pandemic in Korea, and compared them with the trends in the United States.

**METHODS:** We compared the weekly frequencies of gastrointestinal infectious diseases (16 bacterial and 6 viral diseases) in Korea during weeks 5-52 before and after COVID-19. In addition, the weekly frequencies of 5 gastrointestinal infectious diseases in the United States (data from the Centers for Disease Control and Prevention) that overlapped with those in Korea were compared.

**RESULTS:** The mean weekly number of total cases of gastrointestinal infectious diseases in Korea showed a significant decrease (from 522 before COVID-19 to 245 after COVID-19,  $p < 0.01$ ). Only bacterial gastrointestinal infectious diseases caused by *Campylobacter* increased significantly; other bacterial gastrointestinal infectious diseases showed either a decrease or no change. The incidence of all other viral diseases decreased. In the United States, the weekly numbers of *Salmonella*, *Campylobacter*, typhoid, shigellosis, and hepatitis A virus cases sharply decreased after the COVID-19 outbreak. The weekly case numbers of all viral diseases markedly decreased in both countries; however, bacterial gastrointestinal infectious diseases showed a different pattern.

**CONCLUSIONS:** The incidence of gastrointestinal infectious diseases decreased after the COVID-19 outbreak. In contrast, *Campylobacter* infections showed an increasing trend in Korea, but a decreasing trend in the United States. Further studies are needed to elucidate the different trends in bacterial and viral infectious diseases before and after non-pharmaceutical interventions and between different countries.

**KEY WORDS:** COVID-19, Gastrointestinal diseases, Personal protective equipment

**Correspondence:** Boyoung ParkDepartment of Preventive Medicine, Hanyang University College of Medicine, 222 Wangsimni-ro, Seongdong-gu, Seoul 04763, Korea  
E-mail: hayejine@hanmail.net

Received: Jul 10, 2021 / Accepted: Jan 2, 2022 / Published: Jan 3, 2022

This article is available from: <https://e-epih.org/>

© This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

© 2022, Korean Society of Epidemiology

## INTRODUCTION

On March 11, 2020, the World Health Organization (WHO) declared coronavirus disease 2019 (COVID-19) as a pandemic. After the WHO declaration, non-pharmacological interventions (NPIs) against COVID-19 were implemented in several countries [1]. NPIs such as movement restrictions and isolation measures began in March 2020 worldwide [2], and Korea also started implementing nationwide NPIs in March 2020 [3,4]. As a result of NPIs, the weekly positivity rates of viral respiratory infections significantly decreased in 2020 compared with those in the same

weeks in 2010–2019 [5]. Interventions, including personal hygiene and social distancing measures, might affect infectious diseases other than respiratory infectious diseases. However, whether NPIs could influence the development of other infectious diseases is unknown owing to the limited number of studies. Therefore, we examined how trends in the weekly frequencies of gastrointestinal infectious diseases changed before and during the COVID-19 pandemic in Korea, considering trends in COVID-19 incidence, and compared them with the trends in the United States.

## MATERIALS AND METHODS

We analyzed the weekly number of cases of gastrointestinal infectious diseases from the Infectious Disease Surveillance System established by the Korea Disease Control and Prevention Agency (KDCA; formerly the Korea Centers for Disease Control and Prevention). Physicians reported cases to this surveillance system based on the Infectious Disease Control and Prevention Act [6]. After collecting data from the local health authorities, these data were transferred to the KDCA. A report of national statistics was then published using confirmed data from the KDCA [7]. According to reporting guidelines, reported infectious diseases are classified into 3 groups; pathogen identified, suspected infection, or a pathogen carrier [8]. The infectious disease surveillance system in Korea is composed of a mandatory surveillance system and a sentinel surveillance system [6,9]. The mandatory surveillance system requires obligatory reporting from all health centers immediately or within 24 hours of identifying 64 types of infectious diseases. The sentinel surveillance system receives weekly reports on 23 types of infectious diseases from designated health institutions. In this study, we considered gastrointestinal infectious diseases, including cholera, shigellosis, typhoid, paratyphoid, enterohemorrhagic *Escherichia coli* (EHEC), and hepatitis A virus, which are collected through the mandatory surveillance system: *Salmonella*, enteroinvasive *E. coli* (EIEC), *Yersinia enterocolitica*, *Clostridium perfringens*, *Vibrio parahaemolyticus*, enteropathogenic *E. coli* (EPEC), *Bacillus cereus*, *Campylobacter*, enterotoxigenic *E. coli* (ETEC), *Listeria monocytogenes*, *Staphylococcus aureus*, group A rotavirus, astrovirus, adenovirus, norovirus, and sapovirus infections, which are collected through the sentinel surveillance system [9]. For a cross-country comparison with the United States, we used weekly reports from the Nationally Notifiable Diseases Surveillance System (NNDSS) administered by the United States Centers for Disease Control and Prevention (CDC) [10], to which infectious and contagious diseases are immediately reported by healthcare providers [11,12]. The NNDSS is collected from all 50 states and the District of Columbia and monitors about 120 diseases classified by the main transmission: sexually transmitted, foodborne or waterborne, vector borne, injection drug use-associated, and respiratory [13]. As a result, the types of infectious agents reported in the KDCA in Korea and the CDC in the United States differed. Thus, we selected infectious diseases from the CDC that overlapped with those in the KDCA for comparison:

salmonellosis, *Campylobacter* infection, typhoid, shigellosis, and hepatitis A virus infection. For the period before COVID-19, the mandatory surveillance system used the mean weekly number of cases from 2015 to 2019, which was calculated as the sum of all weekly case numbers divided by 5 (the number of years). The sentinel surveillance system used data from July 30, 2017 to 2019, considering the expansion of medical centers for sentinel surveillance in July 2017. For the period after COVID-19, we used the weekly number of cases of each type of gastrointestinal infectious disease from weeks 5–52 of 2020. We then compared the mean weekly case number for the same weeks before and after COVID-19 in the mandatory surveillance system and the sentinel surveillance system using the paired t-test.

We excluded 1–4 weeks from each year to reflect the incidence of COVID-19 because the first COVID-19 patient was identified in Korea on January 20, 2020. Data on the number of weekly COVID-19 cases from the CDC in the United States [14] and from the KDCA in Korea [3] were used. For visualization, we plotted the trends in the weekly number of new COVID-19 cases together with the mean weekly number of each gastrointestinal infectious disease for 2 periods (before and after COVID-19). All statistical analyses were conducted using Microsoft Excel (version 2016; Microsoft, Redmond, WA, USA).

## Ethics statement

This study used data from official websites which are opened to the public. In accordance with Article 13 of the Enforcement Ordinance, this study is subject to institutional review board exception.

## RESULTS

Table 1 shows the mean weekly number of cases of gastrointestinal infectious diseases during weeks 5–52 before COVID-19 compared with that after COVID-19 in Korea and the United States. In Korea, the weekly case number of total gastrointestinal infectious diseases showed a significant decrease (from 522 before COVID-19 to 245 after COVID-19,  $p < 0.01$ ). Statistically significant decreases were observed for all types of viral gastrointestinal infectious diseases (range of decrease: -55.2 to -5.8,  $p < 0.01$ ). The decreases for total bacterial gastrointestinal infectious diseases were not statistically significant ( $p = 0.12$ ). In contrast, the number of total viral gastrointestinal infectious diseases markedly decreased after COVID-19 compared with the period before COVID-19. Meanwhile, in the United States, the number of cases significantly decreased for both total bacterial gastrointestinal diseases and total viral gastrointestinal diseases. In particular, in Korea, statistically significant decreases were observed for *Salmonella*, *V. parahaemolyticus*, ETEC, EIEC, *S. aureus*, *B. cereus*, typhoid, and shigellosis. A statistically significant increase was observed for *Campylobacter* infections. There were no statistically significant differences in *C. perfringens*, *Y. enterocolitica*, *Listeria monocytogenes*, EHEC, cholera, paratyphoid, and EPEC infections. In the United States, significant decreases were found for salmonel-

**Table 1.** Mean weekly numbers of cases gastrointestinal infectious diseases during weeks 5-52 before COVID-19 compared with the case numbers after COVID-19 in Korea and the United States

Variables	Korea				United States			
	Before COVID-19	After COVID-19	Difference	p-value	Before COVID-19	After COVID-19	Difference	p-value
Total no. of gastrointestinal infectious diseases	522.1±83.3	245.0±98.2	-277.1	<0.01	1,021.0±257.0	616.0±205.0	-405	<0.01
<b>Bacterial</b>								
Total	190.7±70.8	182.0±91.6	-8.7	0.12	965.5±252.5	576.2±193.6	-389.3	<0.01
Sentinel surveillance <sup>1</sup>								
<i>Salmonella</i>	52.4±32.2	38.0±27.2	-14.4	<0.01	350.8±132.9	199.0±94.1	-151.8	<0.01
<i>Vibrio parahaemolyticus</i>	2.6±2.6	1.0±1.4	-1.6	<0.01	-	-	-	-
Enterotoxigenic <i>Escherichia coli</i>	1.8±1.6	1.0±1.4	-0.8	<0.01	-	-	-	-
Enteroinvasive <i>Escherichia coli</i>	0.3±0.6	0.1±0.3	-0.2	0.02	-	-	-	-
<i>Campylobacter</i>	57.1±38.3	67.0±55.1	9.9	<0.01	497.3±126.4	333.4±104.5	-163.9	<0.01
<i>Clostridium perfringens</i>	57.9±9.9	59.0±13.9	1.1	0.35	-	-	-	-
<i>Staphylococcus aureus</i>	3.7±2.2	1.0±1.2	-2.7	<0.01	-	-	-	-
<i>Bacillus cereus</i>	0.6±0.6	0.2±0.5	-0.4	<0.01	-	-	-	-
<i>Yersinia enterocolitica</i>	2.2±1.0	2.0±1.6	-0.2	0.73	-	-	-	-
<i>Listeria monocytogenes</i>	0.2±0.3	0.1±0.5	-0.1	0.84	-	-	-	-
Enteropathogenic <i>Escherichia coli</i>	2.3±1.9	2.0±2.5	-0.3	0.60	-	-	-	-
Mandatory surveillance <sup>2</sup>								
Cholera	0.1±0.1	0.1±0.1	-0.1	0.15	-	-	-	-
Typhoid	2.6±1.1	0.7±0.9	-1.8	<0.01	2.1±1.0	0.6±0.9	-1.5	<0.01
Paratyphoid	1.1±0.6	1.1±1.3	0.0	0.88	-	-	-	-
Shigellosis	2.4±1.1	0.4±0.7	-2.0	<0.01	115.2±13.7	43.3±29.0	-71.9	<0.01
Enterohemorrhagic <i>Escherichia coli</i>	2.7±1.9	5.5±10.8	2.8	0.06	-	-	-	-
<b>Viral</b>								
Total	371.3±87.9	147.7±80.1	-223.6	<0.01	54.6±10.9	39.1±18.4	-15.5	<0.01
Sentinel surveillance <sup>1</sup>								
Group A rotaviruses	73.0±48.3	23.0±16.8	-50.0	<0.01	-	-	-	-
Astrovirus	15.4±6.0	2.0±5.1	-13.4	<0.01	-	-	-	-
Adenovirus	18.5±6.0	3.0±2.6	-15.5	<0.01	-	-	-	-
Norovirus	95.2±66.1	40.0±59.5	-55.2	<0.01	-	-	-	-
Sapovirus	6.5±3.0	0.7±1.0	-5.8	<0.01	-	-	-	-
Mandatory surveillance <sup>2</sup>								
Hepatitis A	124.0±42.4	77.6±17.0	-46.3	<0.01	54.6±10.9	39.1±18.4	-15.5	<0.01

Values are presented as mean weekly number±standard deviation.

<sup>1</sup>The weekly number of each type of gastrointestinal infectious disease in weeks 5-52 of 2020 with the mean of the same weeks during July 30, 2017-2019 in Korea or during 2015-2019 in United States.


<sup>2</sup>The weekly number of each type of gastrointestinal infectious disease in weeks 5-52 of 2020 with the mean of the same weeks during 2015-2019 in Korea and United States.

losis (-151.8), *Campylobacter* infection (-163.9), typhoid (-1.5), shigellosis (-71.9), and hepatitis A virus infection (-15.5) after COVID-19 compared before COVID-19 ( $p < 0.01$  for all).

Figures 1 and 2 present the epidemic curves of each type of gastrointestinal infectious disease determined by sentinel surveillance and mandatory surveillance during weeks 5–52 before and after COVID-19 in Korea with the weekly number of new COVID-19 cases. After the start of the COVID-19 outbreak, followed by the establishment of NPIs in Korea, the frequency of the 6 types of viral diseases decreased, especially group A rotavirus, astrovirus,


adenovirus, norovirus, and sapovirus infections. In particular, the frequency of astrovirus and sapovirus infections reached nearly 0 from week 41. The epidemic patterns of bacterial diseases did not show a dramatic decrease after COVID-19.

Figure 3 shows that the weekly case numbers of *Salmonella* infection, *Campylobacter* infection, typhoid, shigellosis, and hepatitis A virus infection sharply decreased after the COVID-19 outbreak in the United States. The decrease in the number of shigellosis cases was most pronounced, from 124 on week 9 to 9 on week 18.




**Figure 1.** Mean weekly numbers of gastrointestinal infections identified by sentinel surveillance in Korea before coronavirus disease 2019 (COVID-19) compared with after COVID-19. The red dotted lines indicate the mean weekly incidence in Korea before COVID-19. The black lines show the weekly incidence after COVID-19. The blue zones indicate the COVID-19 incidence in Korea. (A) Total gastrointestinal infectious diseases. (B) *Bacillus cereus*. (C) *Campylobacter*. (D) *Clostridium perfringens*. (E) Enteroinvasive *Escherichia coli*. (F) Enteropathogenic *E. coli*. (G) Enterotoxigenic *E. coli*. (H) *Listeria monocytogenes*. (I) *Salmonella*. (J) *Staphylococcus aureus*. (K) *Vibrio parahaemolyticus*. (L) *Yersinia enterocolitica*. (M) Adenovirus. (N) Astrovirus. (O) Group A rotaviruses. (P) Norovirus. (Q) Sapovirus.





**Figure 2.** Mean weekly numbers of gastrointestinal infections by mandatory surveillance in Korea before coronavirus disease 2019 (COVID-19) compared with after COVID-19. The red dotted lines indicate the mean weekly incidence in Korea before COVID-19. The black lines show the weekly incidence after COVID-19. The blue zones indicate the COVID-19 incidence in Korea. (A) Cholera. (B) Enterohemorrhagic *Escherichia coli*. (C) Paratyphoid. (D) Shigellosis. (E) Typhoid. (F) Hepatitis A.



**Figure 3.** Mean weekly numbers of gastrointestinal infections in the United States during 2015-2019 compared with those in 2020. The red dotted lines indicate the mean weekly incidence in the United States before coronavirus disease 2019 (COVID-19). The black lines show the weekly incidence after COVID-19. The blue zones indicate the COVID-19 incidence in the United States. (A) *Campylobacter*. (B) Hepatitis A. (C) *Salmonella*. (D) Shigellosis. (E) Typhoid.

## DISCUSSION

With respect to the 5 gastrointestinal infections that were collected in both countries, not only the weekly case numbers of each bacterial and viral disease, but also the total case number of bacterial and viral gastrointestinal diseases markedly decreased in the United States. However, Korea showed a different pattern. Specifically, the total number of viral gastrointestinal diseases decreased significantly, while the total number of bacterial gastrointestinal

diseases did not significantly decrease. In addition, each bacterial gastrointestinal disease had distinct trends.

Infectious diseases are expected to decrease after the COVID-19 pandemic due to worldwide NPIs. In Korea, the total number of viral gastrointestinal infections showed a considerable decrease, but the decreases in the total bacterial gastrointestinal infections did not reach a statistically significant level. Unlike the viral gastrointestinal infectious diseases, there were fluctuations in the trends of bacterial gastrointestinal infections (downward, stable, and then

upward) in 2020 compared with previous years. Significant decreases were observed for *Salmonella*, *V. parahaemolyticus*, ETEC, EIEC, *S. aureus*, *B. cereus*, typhoid, and shigellosis. However, only the increase in *Campylobacter* infections was statistically significant. NPIs, such as hand hygiene and wearing a mask, can reduce the spread of viral diseases [15], and these viral gastrointestinal infectious diseases also showed a significant decrease like that observed for respiratory viruses, suggesting the existence of a food-borne or waterborne infection route for gastrointestinal viruses [16,17]. However, *C. perfringens* and EHEC did not show significant increases. This finding was probably due to an unstable small number. In the case of EHEC, the outbreak in 2019 may have affected the increase in EHEC [18]. Bacterial gastrointestinal infections were difficult to identify both before and after the COVID-19 outbreak. Bacterial pathogens propagate through rotten food or water, especially during the summer [19,20].

Since the types of gastrointestinal infections collected in Korea and the United States differed, the possibility of a direct comparison would be limited; however, there are several potential reasons for the observed divergence. First, the United States issued stay-at-home orders between March and April 2020. In Korea, personal hygiene measures, such as mask-wearing, were used. Second, compared to the United States, Korea implemented a less strict social distancing policy, and therefore, accessibility to medical care may have been lower in the United States than in Korea [7]. Consequently, people who had gastrointestinal infectious diseases with less severe symptoms might have had a limited ability to access medical centers during the COVID-19 pandemic than during the period before it, which could have had a greater impact in the United States than in Korea. Thus, despite the possibility of under-reporting followed by underestimation of actual infection numbers in both countries, this would be more impactful in the United States [21]. Despite these potential reasons, *Campylobacter* infections showed clearly different patterns between Korea and the United States. Gastrointestinal infectious diseases have a wide range of infection pathways. The agents of *Campylobacter* infections are animals, including dogs, cats, pigs, sheep, rodents, birds, and poultry [9]. According to a Korean study, retail raw chickens were confirmed in about 68% of cases for *Campylobacter*. After the COVID-19 outbreak, the annual consumption of chicken per adult increased by 1.2 kg compared to COVID-19 [22]. Thus, *Campylobacter* infections are a zoonotic contagious disease that can infect humans through food based on these assumptions. Additionally, in Korea, cluster outbreaks in restaurants accounted for the highest number of cases before COVID-19; however, after COVID-19, the most common place of cluster outbreaks shifted to pre-primary schools [9]. In Korea, kindergartens and pre-primary schools have different purposes. Kindergartens focus on education, and pre-primary schools focus on childcare. Therefore, this could explain the poor control of *Campylobacter* in the context of COVID-19. Our data, including the mandatory surveillance and sentinel surveillance, and, in the case of sentinel surveillance, the number of hospitals (including public hospitals) collecting data, in-

creased from July 30, 2017 [23]. Thus, we compared the periods before and after COVID-19 separately for each surveillance system.

In conclusion, while the incidence of viral gastrointestinal diseases, including hepatitis A, prominently decreased after COVID-19, bacterial diseases showed different trends in Korea and a decreasing trend in the United States. Further studies are needed to elucidate the different trends in bacterial and viral infectious diseases before and after NPIs and between different countries, as well as considering the characteristics of each pathogen.

## CONFLICT OF INTEREST

The authors have no conflicts of interest to declare for this study.

## FUNDING

This research was supported by the Government-wide R&D Fund Project for Infectious Disease Research (GFID), Republic of Korea (grant No. HG18C00880).

This study was conducted under the research project “Research and Development on Integrated Surveillance System for Early Warning of Infectious Diseases” (RISEWIDs).

## ACKNOWLEDGEMENTS

None.

## AUTHOR CONTRIBUTIONS

Conceptualization: Park B, Kim S. Data curation: Kim S, Kim J. Formal analysis: Kim S. Funding acquisition: Choi BY. Methodology: Park B, Kim S. Project administration: Park B, Choi BY. Visualization: Kim S. Writing – original draft: Kim S, Kim J, Park B. Writing – review & editing: Kim S, Park B, Choi BY.

## ORCID

Soyeoun Kim: <https://orcid.org/0000-0001-9158-5590>; Jinhyun Kim: <https://orcid.org/0000-0002-2740-7535>; Bo Youl Choi: <https://orcid.org/0000-0003-0115-5736>; Boyoung Park: <https://orcid.org/0000-0003-1902-3184>

## REFERENCES

1. World Health Organization. WHO Director-General's opening remarks at the media briefing on COVID-19 - 30 March 2020 [cited 2021 Jun 10]. Available from: <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19--30-march-2020>.
2. Askatas N, Tatsiramos K, Verheyden B. Estimating worldwide effects of non-pharmaceutical interventions on COVID-19 incidence and population mobility patterns using a multiple-event study.

- Sci Rep 2021;11:1972.
3. Korea Disease Control and Prevention Agency. Coronavirus (COVID-19), Republic of Korea [cited 2020 Jun 11]. Available from: <http://ncov.mohw.go.kr/en>.
  4. Dighe A, Cattarino L, Cuomo-Dannenburg G, Skarp J, Imai N, Bhatia S, et al. Response to COVID-19 in South Korea and implications for lifting stringent interventions. *BMC Med* 2020;18:321.
  5. Yum S, Hong K, Sohn S, Kim J, Chun BC. Trends in viral respiratory infections during COVID-19 pandemic, South Korea. *Emerg Infect Dis* 2021;27:1685-1688.
  6. Korea Ministry of Government Legislation. Infectious Disease Control and Prevention Act [cited 2021 Nov 22]. Available from: <http://law.go.kr/> (Korean).
  7. Park S, Cho E. National infectious diseases surveillance data of South Korea. *Epidemiol Health* 2014;36:e2014030.
  8. Park M. Infectious disease-related laws: prevention and control measures. *Epidemiol Health* 2017;39:e2017033.
  9. Korea Disease Control and Prevention Agency. Infectious disease portal [cited 2021 Jun 10]. Available from: <https://www.kdca.go.kr/npt/biz/npp/nppMain.do> (Korean).
  10. Centers for Disease Control and Prevention. Notifiable infectious disease data tables; 2021 [cited 2021 Jun 10]. Available from: <https://www.cdc.gov/nndss/data-statistics/infectious-tables/index.html>.
  11. Troppy S, Haney G, Cocoros N, Cranston K, DeMaria A Jr. Infectious disease surveillance in the 21st century: an integrated web-based surveillance and case management system. *Public Health Rep* 2014;129:132-138.
  12. Adams DA, Jajosky RA, Ajani U, Kriseman J, Sharp P, Onwen DH, et al. Summary of notifiable diseases--United States, 2012. *MMWR Morb Mortal Wkly Rep* 2014;61:1-121.
  13. Crane MA, Popovic A, Panaparambil R, Stolbach AI, Romley JA, Ghanem KG. Reporting of infectious diseases in the United States during the COVID-19 pandemic. *Clin Infect Dis* 2021:ciab529.
  14. CDC COVID-19 Response Team. Coronavirus disease 2019 in children - United States, February 12-April 2, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:422-426.
  15. Laszkowska M, Kim J, Faye AS, Joelson AM, Ingram M, Truong H, et al. Prevalence of *Clostridioides difficile* and other gastrointestinal pathogens in patients with COVID-19. *Dig Dis Sci* 2021; 66:4398-4405.
  16. Matsuyama R, Miura F, Nishiura H. The transmissibility of noroviruses: statistical modeling of outbreak events with known route of transmission in Japan. *PLoS One* 2017;12:e0173996.
  17. Matsuyama R, Miura F, Tsuzuki S, Nishiura H. Household transmission of acute gastroenteritis during the winter season in Japan. *J Int Med Res* 2018;46:2866-2874.
  18. Lee SH, Yun JW, Lee JH, Jung YH, Lee DH. Trends in recent waterborne and foodborne disease outbreaks in South Korea, 2015-2019. *Osong Public Health Res Perspect* 2021;12:73-79.
  19. Wen SC, Best E, Nourse C. Non-typhoidal Salmonella infections in children: review of literature and recommendations for management. *J Paediatr Child Health* 2017;53:936-941.
  20. Kim JS, Lee MS, Kim JH. Recent updates on outbreaks of Shiga toxin-producing *Escherichia coli* and its potential reservoirs. *Front Cell Infect Microbiol* 2020;10:273.
  21. Kang HY, Shin E, Kim YS, Kim JK. Comparing physicians' reporting propensity with active and passive surveillance systems in South Korea. *J Korean Med Assoc* 2014;57:167-175.
  22. Han K, Jang SS, Choo E, Heu S, Ryu S. Prevalence, genetic diversity, and antibiotic resistance patterns of *Campylobacter jejuni* from retail raw chickens in Korea. *Int J Food Microbiol* 2007;114: 50-59.
  23. Korea Disease Control and Prevention Agency. Guideline for water & foodborne diseases prevention and control 2022 [cited 2021 Jun 10]. Available from: <https://www.kdca.go.kr/npt/biz/npp/portal/nppPblctDtaView.do?pblctDtaSeAt=8&pblctDtaSn=2567> (Korean).