

CORRECTION

Open Access



# Correction to: Influence of mechanical and TGF- $\beta$ 3 stimulation on the tenogenic differentiation of tonsil-derived mesenchymal stem cells

Jaeyeon Wee<sup>1†</sup>, Hyang Kim<sup>2†</sup>, Sang-Jin Shin<sup>3</sup>, Taeyong Lee<sup>4</sup> and Seung Yeol Lee<sup>2,5\*</sup>

## Correction to: *BMC Mol Cell Biol* 23, 3 (2022)

<https://doi.org/10.1186/s12860-021-00400-7>

Following publication of the original article [1], a typesetting error was noticed in the reference list. References 17 to 31 were published in the incorrect order. The correct order of references 17 to 31 are given below and the original article [1] has been corrected. The publishers apologize for the error.

17. Ryu KH, Cho KA, Park HS, Kim JY, Woo SY, Jo I, et al. Tonsil-derived mesenchymal stromal cells: evaluation of biologic, immunologic and genetic factors for successful banking. *Cytotherapy*. 2012;14(10):1193–202. <https://doi.org/10.3109/14653249.2012.706708>.

18. Choi JS, Lee BJ, Park HY, Song JS, Shin SC, Lee JC, et al. Effects of donor age, long-term passage culture, and cryopreservation on tonsil-derived mesenchymal stem cells. *Cell Physiol Biochem*. 2015;36(1):85–99. <https://doi.org/10.1159/000374055>.

19. Koh RH, Jin Y, Kang BJ, Hwang NS. Chondrogenically primed tonsil-derived mesenchymal stem cells encapsulated in riboflavin-induced photocrosslinking

collagen-hyaluronic acid hydrogel for meniscus tissue repairs. *Acta Biomater*. 2017;53:318–28. <https://doi.org/10.1016/j.actbio.2017.01.081>.

20. Park J, Kim IY, Patel M, Moon HJ, Hwang SJ, Jeong B. 2D and 3D hybrid Systems for Enhancement of chondrogenic differentiation of tonsil-derived mesenchymal stem cells. *Adv Funct Mater*. 2015;25(17):2573–82. <https://doi.org/10.1002/adfm.201500299>.

21. Patel M, Moon HJ, Ko DY, Jeong B. Composite system of graphene oxide and polypeptide thermogel as an injectable 3D scaffold for adipogenic differentiation of tonsil-derived mesenchymal stem cells. *ACS Appl Mater Interfaces*. 2016;8(8):5160–9. <https://doi.org/10.1021/acsami.5b12324>.

22. Li YH, Ramcharan M, Zhou ZP, Leong DJ, Akinbiyi T, Majeska RJ, et al. The role of Scleraxis in fate determination of mesenchymal stem cells for tenocyte differentiation. *Sci Rep-Uk*. 2015;5(1). <https://doi.org/10.1038/srep13149>.

23. Zhang JY, Wang JHC. The effects of mechanical loading on tendons - an in vivo and in vitro model study. *Plos One*. 2013;8(8).

24. Jager M, Feser T, Denck H, Krauspe R. Proliferation and osteogenic differentiation of mesenchymal stem cells cultured onto three different polymers in vitro. *Ann Biomed Eng*. 2005;33(10):1319–32. <https://doi.org/10.1007/s10439-005-5889-2>.

25. Fan HB, Liu HE, Toh SL, Goh JCH. Enhanced differentiation of mesenchymal stem cells co-cultured with ligament fibroblasts on gelatin/silk fibroin hybrid scaffold.

The original article can be found online at <https://doi.org/10.1186/s12860-021-00400-7>.

\*Correspondence: [kernels00@naver.com](mailto:kernels00@naver.com)

<sup>†</sup>Jaeyeon Wee and Hyang Kim contributed equally to this work.

<sup>2</sup>New Horizon Biomedical Engineering Institute, Myongji Hospital, Goyang, South Korea

<sup>5</sup>Department of Orthopaedic Surgery, Myongji Hospital, Hanyang University College of Medicine, 55, Hwasu-ro 14beon-gil, Deogyang-gu, Goyang, Gyeonggi 10475, South Korea

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Biomaterials. 2008;29(8):1017–27. <https://doi.org/10.1016/j.biomaterials.2007.10.048>.

26. Agathocleous M, Harris WA. Metabolism in physiological cell proliferation and differentiation. *Trends Cell Biol.* 2013;23(10):484–92. <https://doi.org/10.1016/j.tcb.2013.05.004>.

27. Lima EG, Bian L, Ng KW, Mauck RL, Byers BA, Tuan RS, et al. The beneficial effect of delayed compressive loading on tissue-engineered cartilage constructs cultured with TGF-beta 3. *Osteoarthr Cartil.* 2007;15(9):1025–33. <https://doi.org/10.1016/j.joca.2007.03.008>.

28. Thorpe SD, Buckley CT, Vinardell T, O'Brien FJ, Campbell VA, Kelly DJ. The response of bone marrow-derived mesenchymal stem cells to dynamic compression following TGF-beta 3 induced Chondrogenic differentiation. *Ann Biomed Eng.* 2010;38(9):2896–909. <https://doi.org/10.1007/s10439-010-0059-6>

29. Nam HY, Pingguan-Murphy B, Abbas AA, Merican AM, Kamarul T. Uniaxial cyclic tensile stretching at 8% strain exclusively promotes Tenogenic differentiation of human bone marrow-derived mesenchymal stromal cells. *Stem Cells Int.* 2019;2019:1–16. <https://doi.org/10.1155/2019/9723025>.

30. Bottagisio M, Lopa S, Granata V, Talo G, Bazzocchi C, Moretti M, et al. Different combinations of growth factors for the tenogenic differentiation of bone marrow mesenchymal stem cells in monolayer culture and in fibrin-based three-dimensional constructs. *Differentiation.* 2017;95:44–53. <https://doi.org/10.1016/j.diff.2017.03.001>.

31. Kraus A, Woon C, Raghavan S, Megerle K, Pham H, Chang J. Co-culture of human adipose-derived stem cells with tenocytes increases proliferation and induces differentiation into a tenogenic lineage. *Plast Reconstr Surg.* 2013;132(5):754e–66e

#### Author details

<sup>1</sup>School of Medicine, CHA University, Pocheon, South Korea. <sup>2</sup>New Horizon Biomedical Engineering Institute, Myongji Hospital, Goyang, South Korea.

<sup>3</sup>Department of Orthopaedic Surgery, Ewha Womans University Seoul Hospital, Seoul, South Korea. <sup>4</sup>Division of Mechanical and Biomedical Engineering, Ewha Womans University, Seoul, South Korea. <sup>5</sup>Department of Orthopaedic Surgery, Myongji Hospital, Hanyang University College of Medicine, 55, Hwasu-ro 14beon-gil, Deogyang-gu, Goyang, Gyeonggi 10475, South Korea.

Published online: 28 January 2022

#### Reference

1. Wee J, Kim H, Shin SJ, et al. Influence of mechanical and TGF- $\beta$ 3 stimulation on the tenogenic differentiation of tonsil-derived mesenchymal stem cells. *BMC Mol Cell Biol.* 2022;23:3. <https://doi.org/10.1186/s12860-021-00400-7>.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

