

# World Journal of *Clinical Cases*

*World J Clin Cases* 2021 February 16; 9(5): 999-1246



**MINIREVIEWS**

- 999 Remote nursing training model combined with proceduralization in the intensive care unit dealing with patients with COVID-19  
*Wang H, Kang K, Gao Y, Yang B, Li J, Wang L, Bi Y, Yu KJ, Dai QQ, Zhao MY*

**ORIGINAL ARTICLE****Case Control Study**

- 1005 Metabolic syndrome, ApoE genotype, and cognitive dysfunction in an elderly population: A single-center, case-control study  
*Wang JY, Zhang L, Liu J, Yang W, Ma LN*

- 1016 Serum neuron-specific enolase: A promising biomarker of silicosis  
*Huang HB, Huang JL, Xu XT, Huang KB, Lin YJ, Lin JB, Zhuang XB*

**Retrospective Study**

- 1026 Biochemical recurrence of pathological T2+ localized prostate cancer after robotic-assisted radical prostatectomy: A 10-year surveillance  
*Yang CH, Lin YS, Ou YC, Weng WC, Huang LH, Lu CH, Hsu CY, Tung MC*

**Observational Study**

- 1037 Clinical characteristics of perineal endometriosis: A case series  
*Liang Y, Zhang D, Jiang L, Liu Y, Zhang J*
- 1048 Safety of gastrointestinal endoscopy in patients with acute coronary syndrome and concomitant gastrointestinal bleeding  
*Elkafrawy AA, Ahmed M, Alomari M, Elkaryoni A, Kennedy KF, Clarkston WK, Campbell DR*

**SYSTEMATIC REVIEWS**

- 1058 Clinical features of SARS-CoV-2-associated encephalitis and meningitis amid COVID-19 pandemic  
*Huo L, Xu KL, Wang H*

**CASE REPORT**

- 1079 Neuropathy and chloracne induced by 3,5,6-trichloropyridin-2-ol sodium: Report of three cases  
*Ma Y, Cao X, Zhang L, Zhang JY, Qiao ZS, Feng WL*
- 1087 Effect of rifampicin on anticoagulation of warfarin: A case report  
*Hu YN, Zhou BT, Yang HR, Peng QL, Gu XR, Sun SS*
- 1096 Severe lumbar spinal stenosis combined with Guillain-Barré syndrome: A case report  
*Xu DF, Wu B, Wang JX, Yu J, Xie JX*

- 1103** Treatment of pediatric intracranial dissecting aneurysm with clipping and angioplasty, and next-generation sequencing analysis: A case report and literature review  
*Sun N, Yang XY, Zhao Y, Zhang QJ, Ma X, Wei ZN, Li MQ*
- 1111** Imaging characteristics of a rare case of monostotic fibrous dysplasia of the sacrum: A case report  
*Liu XX, Xin X, Yan YH, Ma XW*
- 1119** Primary aldosteronism due to bilateral micronodular hyperplasia and concomitant subclinical Cushing's syndrome: A case report  
*Teragawa H, Oshita C, Orita Y, Hashimoto K, Nakayama H, Yamazaki Y, Sasano H*
- 1127** Management of corneal ulceration with a moisture chamber due to temporary lagophthalmos in a brain injury patient: A case report  
*Yu XY, Xue LY, Zhou Y, Shen J, Yin L*
- 1132** Bronchoscopy for diagnosis of COVID-19 with respiratory failure: A case report  
*Chen QY, He YS, Liu K, Cao J, Chen YX*
- 1139** Pembrolizumab as a novel therapeutic option for patients with refractory thymic epithelial tumor: A case report  
*Wong-Chong J, Bernadach M, Ginzac A, Veyssi ere H, Durando X*
- 1148** Successful bailout stenting strategy against rare spontaneous retrograde dissection of partially absorbed magnesium-based resorbable scaffold: A case report  
*Liao ZY, Liou JY, Lin SC, Hung HF, Chang CM, Chen LC, Chua SK, Lo HM, Hung CF*
- 1156** Chronic myelomonocytic leukemia-associated pulmonary alveolar proteinosis: A case report and review of literature  
*Chen C, Huang XL, Gao DQ, Li YW, Qian SX*
- 1168** Obturator nerve impingement caused by an osteophyte in the sacroiliac joint: A case report  
*Cai MD, Zhang HF, Fan YG, Su XJ, Xia L*
- 1175** Venetoclax in combination with chidamide and dexamethasone in relapsed/refractory primary plasma cell leukemia without t(11;14): A case report  
*Yang Y, Fu LJ, Chen CM, Hu MW*
- 1184** Heterochronic triple primary malignancies with Epstein-Barr virus infection and tumor protein 53 gene mutation: A case report and review of literature  
*Peng WX, Liu X, Wang QF, Zhou XY, Luo ZG, Hu XC*
- 1196** Negative conversion of autoantibody profile in chronic hepatitis B: A case report  
*Zhang X, Xie QX, Zhao DM*
- 1204** Dumbbell-shaped solitary fibrous tumor in the parapharyngeal space: A case report  
*Li YN, Li CL, Liu ZH*
- 1210** Spontaneous small bowel perforation secondary to *Vibrio parahaemolyticus* infection: A case report  
*Chien SC, Chang CC, Chien SC*

- 1215** Management protocol for Fournier’s gangrene in sanitary regime caused by SARS-CoV-2 pandemic: A case report  
*Grabińska A, Michalczyk Ł, Banaczyk B, Szyrło T, Ząbkowski T*
- 1221** Infective bicuspid aortic valve endocarditis causing acute severe regurgitation and heart failure: A case report  
*Hou C, Wang WC, Chen H, Zhang YY, Wang WM*
- 1228** Endoscopic repair of delayed stomach perforation caused by penetrating trauma: A case report  
*Yoon JH, Jun CH, Han JP, Yeom JW, Kang SK, Kook HY, Choi SK*
- 1237** Bilateral musculocutaneous neuropathy: A case report  
*Jung JW, Park YC, Lee JY, Park JH, Jang SH*

**ABOUT COVER**

Editorial Board Member of *World Journal of Clinical Cases*, Dr. Antonio Corvino is a PhD in the Motor Science and Wellness Department at University of Naples "Parthenope". In 2008, he obtained his MD degree from the School of Medicine, Second University of Naples. Then, he completed a residency in Radiology in 2014 at University Federico II of Naples. In 2015, he undertook post-graduate training at Catholic University of Rome, obtaining the 2<sup>nd</sup> level Master's degree in "Internal Ultrasound Diagnostic and Echo-Guided Therapies". In 2016-2018, he served on the directive board of Young Directive of Italian Society of Ultrasound in Medicine and Biology. His ongoing research interests involve ultrasound and ultrasound contrast media in abdominal and non-abdominal applications, etc. (L-Editor: Filipodia)

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, Scopus, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for *WJCC* as 1.013; IF without journal self cites: 0.991; Ranking: 120 among 165 journals in medicine, general and internal; and Quartile category: Q3. The *WJCC*'s CiteScore for 2019 is 0.3 and Scopus CiteScore rank 2019: General Medicine is 394/529.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Jia-Hui Li; Production Department Director: Yu-Jie Ma; Editorial Office Director: Jin-Lai Wang.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Thrice Monthly

**EDITORS-IN-CHIEF**

Dennis A Bloomfield, Sandro Vento, Bao-gan Peng

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

February 16, 2021

**COPYRIGHT**

© 2021 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Bilateral musculocutaneous neuropathy: A case report

Ji Won Jung, Yu Chan Park, Jae Young Lee, Jae Hyeon Park, Seong Ho Jang

**ORCID number:** Ji Won Jung 0000-0001-6063-6244; Yu Chan Park 0000-0002-3395-8528; Jae Young Lee 0000-0002-8881-154X; Jae Hyeon Park 0000-0001-5619-4818; Seong Ho Jang 0000-0001-6267-0980.

**Author contributions:** Jung JW reviewed the literature and contributed to manuscript drafting; Park YC reviewed the literature and interviewed the patient; Lee JY and Park JH performed the electrodiagnostic study and contributed to interpretation of imaging and electrodiagnostic studies; Jang SH was responsible for manuscript revision and intellectual content; All authors issued approval for submission of the final version.

**Supported by** The Data Construction Learning Program for artificial intelligence (20200803-FS4T, Sector 20 Medical and Health AI Data) funded by the National Information Society Agency, Ministry of Science and ICT.

**Informed consent statement:** Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** The authors have no conflicts of interest to declare.

**CARE Checklist (2016) statement:** The authors have read the CARE

**Ji Won Jung, Yu Chan Park, Jae Young Lee,** Department of Rehabilitation Medicine, Hanyang University Hospital, Seoul 04763, South Korea

**Jae Hyeon Park, Seong Ho Jang,** Department of Rehabilitation Medicine, Hanyang University Guri Hospital, Guri 11923, South Korea

**Corresponding author:** Seong Ho Jang, MD, PhD, Professor, Department of Rehabilitation Medicine, Hanyang University Guri Hospital, No. 153 Gyeongchun-ro, Guri 11923, South Korea. [systole77@hanmail.net](mailto:systole77@hanmail.net)

### Abstract

#### BACKGROUND

Isolated musculocutaneous nerve injury is a rare condition. Herein, we report the first case of bilateral musculocutaneous neuropathy after vigorous stretching of both upper extremities with normal results of sensory nerve action potential. Clinicians should be aware of this rare condition that can appear bilaterally. In addition, the interpretation of the aberrant electrodiagnostic study results of this case was discussed.

#### CASE SUMMARY

A 29-year-old male complaining of bilateral forearm tingling and upper extremity weakness visited the outpatient clinic. The symptoms began 6 mo prior, and he visited another hospital before visiting our department. The diagnosis was not made even after cervical spine magnetic resonance imaging, electrodiagnostic study, brain magnetic resonance imaging, and arteriography were conducted. The patient performed unique exercises that stretched the pectoralis minor and coracobrachialis muscles. On the follow-up electrodiagnostic study, abnormal spontaneous activities in the bilateral biceps and brachialis muscles were observed. The patient was diagnosed with bilateral musculocutaneous neuropathy. Steroid pulse therapy was administered for approximately 6 wk. After treatment, his muscle strength returned to the predisease condition.

#### CONCLUSION

Clinicians should be aware of this condition, have adequate understanding of anatomy, and advise to correct inappropriate exercises.

**Key Words:** Musculocutaneous nerve; Peripheral neuropathy; Electrodiagnosis; Exercise; Diagnosis; Case report

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Checklist (2016) and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Unsolicited manuscript

**Specialty type:** Clinical neurology

**Country/Territory of origin:** South Korea

**Peer-review report's scientific quality classification**

Grade A (Excellent): A, A  
 Grade B (Very good): 0  
 Grade C (Good): C  
 Grade D (Fair): 0  
 Grade E (Poor): 0

**Received:** December 3, 2020

**Peer-review started:** December 3, 2020

**First decision:** December 13, 2020

**Revised:** December 14, 2020

**Accepted:** December 22, 2020

**Article in press:** December 22, 2020

**Published online:** February 16, 2021

**P-Reviewer:** de Melo FF, Fu T

**S-Editor:** Zhang H

**L-Editor:** Filipodia

**P-Editor:** Li JH



**Core Tip:** We report the first case of bilateral musculoskeletal neuropathy after vigorous stretching of both upper extremities with normal results of sensory nerve action potential. We recommend clinicians be aware of this rare condition that can occur bilaterally. In addition, the interpretation of aberrant electrodiagnostic study results of this case is discussed.

**Citation:** Jung JW, Park YC, Lee JY, Park JH, Jang SH. Bilateral musculoskeletal neuropathy: A case report. *World J Clin Cases* 2021; 9(5): 1237-1246

**URL:** <https://www.wjgnet.com/2307-8960/full/v9/i5/1237.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v9.i5.1237>

## INTRODUCTION

Isolated musculoskeletal nerve injury is a rare condition and generally associated with brachial plexopathy<sup>[1]</sup>. In previous reports, isolated musculoskeletal injury occurred after intense physical activity, repeated strenuous upper extremity activity<sup>[2,3]</sup>, after quick and strong movements<sup>[4]</sup>, trauma<sup>[5]</sup>, surgery<sup>[6]</sup>, wrestling, or incorrect pitching by a pitcher<sup>[7,8]</sup>. Musculoskeletal lesions usually occur proximal to the biceps and brachialis muscles<sup>[9]</sup>. The most common lesion site is the coracobrachialis muscle<sup>[9]</sup>. Due to hypertrophy or powerful contraction of the muscle, mechanical and ischemic nerve injury can occur<sup>[9]</sup>. In previous reports, patients presented with unilateral arm symptoms.

Electrodiagnostic evaluation is important for diagnosing peripheral neuropathy. Appropriate electrodiagnostic evaluation is needed for differential diagnosis<sup>[10]</sup>. Peripheral nerves respond to injury in various ways, and test findings are different depending on the timing of the test<sup>[11]</sup>. Furthermore, peripheral neuropathy has neurapraxic nature and is often self-resolving<sup>[5]</sup>. Therefore, performing and interpreting electrodiagnostic studies properly can be difficult during diagnosing neuropathy.

Herein, we report a case of bilateral musculoskeletal neuropathy after vigorous stretching of both upper extremities with normal results of sensory nerve action potential (SNAP). Because the importance of various stretching and exercise is emphasized, we recommend clinicians be aware of this rare condition that can occur bilaterally. In addition, the interpretation of the aberrant electrodiagnostic study results in this case is discussed.

## CASE PRESENTATION

### Chief complaints

A 29-year-old male complaining of bilateral forearm tingling and bilateral upper extremity weakness visited the outpatient clinic of our department.

### History of present illness

His primary symptoms were bilateral forearm tingling and pain. These symptoms began 6 mo prior to the visit. Within a few days, bilateral upper extremity weakness also began. There was no improvement in symptoms for several weeks. Thus, he visited the neurology department of another hospital. Under suspicion of radiculopathy, cervical spine magnetic resonance imaging (MRI) with enhancement and electrodiagnostic study were performed. However, abnormalities were not found. During the follow-up period at that hospital, brain MRI, computed tomography, and arteriography of the left upper extremity were performed, but a diagnosis could not be made.

Later, when the patient visited the hospital, he complained of bilateral forearm tingling, weakness, and bilateral arm muscle spasms.

### History of past illness

The patient was not taking any medications and had no previous diagnoses. The patient did not have any history of hospitalization or surgery.

**Personal and family history**

He had no familial history of congenital, allergic, or systemic disease. The patient smoked approximately half a pack of cigarettes a day for 5 years and did not drink alcohol.

**Physical examination**

On physical examination, no external wound was observed. The circumference of both arms was not different and distinct atrophy of biceps or deltoid muscles was not present. The range of motion for both shoulders and elbow joints was not limited. The pain was not aggravated by movement.

Paresthesia was present in both forearms, but the symptom site did not match with peripheral nerve distribution or cervical root dermatome. Manual muscle testing was grade 5 throughout the right and left upper extremities except in bilateral elbow and shoulder flexion. Manual muscle test of bilateral elbow and shoulder flexion showed grade 4. The patient could flex both his elbows with a 2 kg dumbbell but not with a 5 kg. The deep tendon reflex of both bicep muscles was symmetrically decreased.

Before the symptoms began, he started working out at a fitness center. He did weight training, which is commonly practiced. However, he extensively stretched his pectoralis minor muscles. [Figure 1](#) shows the stretching exercise of the pectoralis minor that the patient described. Both shoulder joints were mildly extended, 90° externally rotated, and 45° abducted. Both scapulae were retracted, and the elbow joints were bent approximately 90°. The patient placed his elbows on the wall right next to the door and pushed his trunk forward.

**Laboratory examinations**

There were no abnormalities in laboratory tests including complete blood count, electro profile, liver function test, kidney function test, routine urine analysis, blood coagulation test, and thyroid function test. Acetylcholine receptor antibody, erythrocyte sedimentation rate, C-reactive protein, vitamin B12, and folate were within normal limits. In addition, blood calcium, ionized calcium, creatine kinase, and phosphate were within normal limits.

Tables 1 and 2 show the electrodiagnostic study of the patient. Bilateral axillary, musculocutaneous, median, and ulnar compound motor action potential (CMAP) were within normal limits. Bilateral median, ulnar, and bilateral lateral antebrachial cutaneous SNAP were within normal limits. Bilateral median F-waves were within normal limits. Electromyography of the bilateral upper extremities showed abnormal spontaneous activities in the bilateral biceps and brachialis muscles with reduced recruitment and interference pattern. These electrophysiologic findings were indicative of bilateral musculocutaneous neuropathy.

Tables 3 and 4 show the patient's electrodiagnostic study performed at a previous hospital approximately 4 mo before the follow-up study. Compared with [Table 3](#), [Table 1](#) shows that amplitude of each musculocutaneous CMAP was decreased.

**Imaging examinations**

As mentioned above, electrophysiologic findings were indicative of bilateral musculocutaneous neuropathy. To confirm the diagnosis, MRI of both arms was performed.

Figures 2 and 3 showed the MRI of both arms and brachial plexus of the patient. Significant abnormality was not observed in either brachial plexus and distal peripheral nerves.

[Figure 4](#) showed the patient's cervical spine MRI performed at a previous hospital approximately 6 mo before the visit to our clinic. Specific abnormalities were not observed on cervical spine MRI.

---

**FINAL DIAGNOSIS**

Final diagnosis of the presented case was bilateral musculocutaneous neuropathy with active denervation.

---

**TREATMENT**

Steroid pulse therapy was administered for treatment. For the first 2 d, the patient was



**Table 1** Electrodiagnostic study: Nerve conduction study

Nerve conduction (sensory)	Recording site	Stimulation site	Peak latency (ms)	Amplitude (µV)	Distance (cm)	Conduction velocity (m/s)	Nerve conduction (motor)	Recording site	Stimulation site	Onset latency (ms)	Amplitude (mV)	Distance (cm)	Conduction velocity (m/s)
Right median	Index finger	Palm	1.65	49.9	7	60.9	Right median	APB	Wrist	2.80	8.4		
		Wrist	2.90	54.8	7	63.6			Antecubital	6.25	8.3	22	63.8
Left median	Index finger	Palm	1.60	59.8	7	63.6	Left median	APB	Wrist	2.95	8.3		
		Wrist	2.80	59.7	7	66.7			Antecubital	6.50	8.3	22	62.0
Right ulnar	Small finger	Wrist	2.70	44.0	13	63.4	Right ulnar	ADM	Wrist	2.05	8.3		
Left ulnar	Small finger	Wrist	2.90	37.0	14	59.6			Below elbow	5.85	8.8	23	60.5
Right lateral antebrachial	Lateral forearm	12 cm proximal	2.40	13.5	11	57.9	Left ulnar	ADM	Wrist	2.15	5.8		
Left lateral antebrachial	Lateral forearm	12 cm proximal	2.60	18.8	13	63.4			Below elbow	6.00	7.4	22	57.1
Right medial antebrachial	Medial forearm	12 cm proximal	2.30	12.6	11	66.7	Right ulnar	ADM	Wrist	2.25	8.2		
Left medial antebrachial	Medial forearm	12 cm proximal	2.40	19.9	12	64.9	(Post-exercise)		Below elbow	5.95	8.4	23	62.2
							Left ulnar	ADM	Wrist	2.45	8.1		
							(Post-exercise)		Below elbow	6.15	9.6	23	62.2
							Right axillary	Deltoid	Erb's point	3.65	8.6		
							Left axillary	Deltoid	Erb's point	4.05	7.2		
							Right musculocutaneous	Biceps	Erb's point	4.25	<b>2.4</b>		
							Left musculocutaneous	Biceps	Erb's point	4.05	<b>1.3</b>		

ADM: Abductor digiti minimi; APB: Abductor pollicis brevis. Bold text indicates abnormal findings.

prescribed 64 mg of triamcinolone (Ledercort®). The following 2 d, 48 mg of triamcinolone was prescribed and 32 mg for the following 3 d. During the treatment period, 30 mg of lansoprazole (Lanston LFTD®), a proton-pump inhibitor, was prescribed daily.

A week after diagnosis, the patient visited the outpatient clinic. His strength

Table 2 Electrodiagnostic study: Electromyography

Muscle	Spontaneous	MUAP	Recruitment pattern	Interference pattern	Muscle	Spontaneous	MUAP	Recruitment pattern	Interference pattern
<b>(Right)</b>	<b>Positive sharp wave</b>				<b>(Left)</b>	<b>Positive sharp wave</b>			
APB	None	Normal	Normal	Full	APB	None	Normal	Normal	Full
First DI	None	Normal	Normal	Full	First DI	None	Normal	Normal	Full
ECRL	None	Normal	Normal	Full	ECRL	None	Normal	Normal	Full
FCR	None	Normal	Normal	Full	FCR	None	Normal	Normal	Full
Biceps	<b>2+</b>	Normal	<b>Reduced</b>	<b>Reduced</b>	Biceps	<b>2+</b>	Normal	<b>Reduced</b>	<b>Reduced</b>
Brachialis	<b>1+</b>	Normal	<b>Reduced</b>	Full	Brachialis	<b>1+</b>	Normal	<b>Reduced</b>	Full
Infraspinatus	None	Normal	Normal	Full	Infraspinatus	None	Normal	Normal	Full
Deltoid	None	Normal	Normal	Full	Deltoid	None	Normal	Normal	Full
Triceps	None	Normal	Normal	Full	Triceps	None	Normal	Normal	Full
Cervical Paraspinal	None	NA	NA	NA	Cervical Paraspinal	None	NA	NA	NA

APB: Abductor pollicis brevis; DI: Dorsal interosseous; ECRL: Extensor carpi radialis longus; FCR: Flexor carpi radialis; MUAP: Motor unit action potential. Bold texts indicate abnormal findings.

improved sufficiently to perform elbow flexion with a 5 kg dumbbell. Because his strength did not recover to the predisease level, he was prescribed an additional 24 mg of triamcinolone for 2 d, 16 mg for 2 d, 8 mg for 3 d, and 4 mg for 4 d.

## OUTCOME AND FOLLOW-UP

On the follow-up visit after 11 d, the patient stated he was slightly improving and had started stretching exercise again. He was instructed not to perform the stretching, and 5 mg intramuscular injection of hydroxocobalamin (Lanobin®) was administered. Another 4 mg of triamcinolone and 30 mg of duloxetine hydrochloride (Cymbalta®) was prescribed for an additional 4 wk.

On the last follow-up visit after 4 wk, his muscle strength returned to the predisease condition. Informed written consent was obtained from the patient for publication of this report and any accompanying images.

## DISCUSSION

In the present case report, the diagnosis of a patient with bilateral musculocutaneous neuropathy was described. We would like to discuss the anatomy, etiology, diagnostic consideration, and differential diagnosis of this rare case.

The musculocutaneous nerve originates from the brachial plexus lateral cord and consists of fibers from the fifth, sixth, and seventh roots of the cervical spinal nerve. Furthermore, the components of the fifth and sixth cervical spinal nerve roots contribute primarily to the musculocutaneous nerve<sup>[12,13]</sup>. It passes through the coracobrachialis muscle and travels between the biceps brachii and brachialis muscles that it innervates<sup>[12]</sup>. The nerve becomes the lateral antebrachial cutaneous nerve between these muscles near the lateral margin of the bicipital aponeurosis<sup>[14]</sup>. The nerve is usually injured at the level of the coracobrachialis muscle and generally patients with this type of injury present with pain and weakness of the biceps brachii and paresthesia of the forearm<sup>[13,15]</sup>.

Various injury mechanisms for musculocutaneous neuropathy have been described. Prolonged positioning of the arm during surgery, direct injury to the nerve during surgery, repetitive vigorous upper extremity activity such as lifting, throwing, or carrying may be the etiology of musculocutaneous neuropathy<sup>[9]</sup>. In addition, a single event of forceful extension of the arm could also be a causative factor<sup>[4]</sup>.

**Table 3 Electrodiagnostic study from previous hospital: Nerve conduction studies**

Nerve conduction (sensory)	Recording site	Recording site	Stimulation site	Onset latency (ms) <sup>1</sup>	Amplitude (µV)	Nerve conduction (motor)	Recording site	Stimulation site	Onset latency (ms)	Amplitude (mV)	Distance (cm)	Conduction velocity (m/s)
Left median	Index finger	Index finger	Wrist	2.29	71.8	Left median	APB	Wrist	3.18	12.1		
Left ulnar	Small finger	Small finger	Wrist	2.34	49.8			Antecubital	6.77	12.4	19	52.9
Left radial	Thumb	Thumb	12 cm proximal	1.82	36.3	Left ulnar	ADM	Wrist	2.24	11.0		
Right medial antebrachial	Medial forearm	Medial forearm	12 cm proximal	1.93	15.9			Below elbow	5.73	11.6	19	54.4
Left medial antebrachial	Medial forearm	Medial forearm	12 cm proximal	1.61	15.8	Right axillary	Deltoid	Erb's point	3.70	3.6		
						Left axillary	Deltoid	Erb's point	4.38	3.2		
						Right musculocutaneous	Biceps	Erb's point	5.31	3.8		
						Left musculocutaneous	Biceps	Erb's point	4.95	5.1		

<sup>1</sup>This test was performed at another hospital and latency of sensory nerve action potential was evaluated with onset latency.  
ADM: Abductor digiti minimi; APB: Abductor pollicis brevis.

A case of musculocutaneous nerve injury after repeated sessions of skydiving simulation in a wind-tunnel was also reported<sup>[1]</sup>. Similar to our patient, the subject placed her arms in an abducted, extended, and externally rotated position while experiencing freefall. Our patient repetitively stretched his pectoralis minor muscles with both shoulder joints mildly extended, 90° externally rotated, and 45° abducted. In addition, during the follow-up period when the patient performed the stretching exercise again, his muscle strength recovery was slow. After he was advised to not perform the stretching exercise, the muscle strength recovery was noticeably faster. Therefore, we postulated his stretching exercise was the etiology of the bilateral musculocutaneous neuropathy. The coracobrachialis muscle originates from the coracoid process of the scapula and inserts into the anteromedial surface of the humerus<sup>[16]</sup>. The stretching exercise shown in **Figure 1** may have lengthened his coracobrachialis. Consequently, the musculocutaneous nerve passing through the coracobrachialis muscle might be squeezed. Furthermore, repetitive stretching may have caused the pathological insult and the resulting neuropathy.

Electrodiagnostic studies play an important role in diagnosing peripheral neuropathy<sup>[17]</sup>. The electrodiagnostic findings in the present case report showed that amplitude of each musculocutaneous CMAP was decreased compared with a previous


**Table 4** Electrodiagnostic study from previous hospital: Electromyography

Muscle (Left)	Spontaneous Positive sharp wave	MUAP analysis	Recruitment pattern	Interference pattern
First DI	None	Normal	Normal	Full
ADM	None	Normal	Normal	Full
FCU	None	Normal	Normal	Full

ADM: Abductor digiti minimi; DI: Dorsal interosseous; FCU: Flexor carpi ulnaris; MUAP: Motor unit action potential.




**Figure 1** Stretching exercise of the pectoralis minor muscle. Both shoulder joints were mildly extended, 90° externally rotated, and 45° abducted. Both scapulae were retracted and the elbow joints were bent approximately 90°. The patient placed his elbows on the wall right next to the door and pushed his trunk forward.




**Figure 2** Right arm and brachial plexus magnetic resonance imaging. A: Transverse view; B: Coronal view. Significant abnormality was not observed in the right brachial plexus and distal peripheral nerves.

study. In addition, we performed a nerve conduction study of the bilateral lateral antebrachial cutaneous nerve and needle electromyography of the biceps brachii, brachialis, and deltoid muscles. Proper evaluation of muscles and nerves based on presenting symptoms is important.

In general, electrodiagnostic findings of musculocutaneous neuropathy include abnormalities of lateral antebrachial cutaneous SNAP, musculocutaneous CMAP, and electromyography abnormalities of the biceps brachii and brachialis muscles<sup>[9]</sup>. However, in our case, lateral antebrachial cutaneous SNAP was within normal limits. There are two probable explanations for the test results.



**Figure 3 Left arm and brachial plexus magnetic resonance imaging.** A: Transverse view; B: Coronal view. Significant abnormality was not observed in the left brachial plexus and distal peripheral nerves.



**Figure 4 Cervical spine magnetic resonance imaging taken at a previous hospital.** A: Transverse view; B: Sagittal view. Specific abnormalities were not observed on cervical spine magnetic resonance imaging.

First, because SNAP is detectable later than CMAP after peripheral nerve injury<sup>[11]</sup>, the SNAP amplitude possibly decreased but might have not yet reached the upper normal limit at that time point. In a previous case report, Kissel *et al*<sup>[9]</sup> reported an electrodiagnostic study should be performed 10-21 d after peripheral nerve injury. At this time, the specific lesion extent and amount of axonal loss can be assessed<sup>[9]</sup>. However, the mechanism of injury in the present case was not due to single insult. Because the nerve damage occurred due to numerous repetitive stretching, the injury might have acute-on-chronic trait. Therefore, conducting an electrodiagnostic study at the appropriate time was difficult.

Second, the lateral antebrachial cutaneous SNAP values can be interpreted as abnormal. In general, electrodiagnostic studies are conducted with left-right comparison. However, in some cases when left-right comparison cannot be performed, the results are interpreted by comparison with other intact nerves<sup>[18]</sup>. The reference values of lateral and medial antebrachial cutaneous SNAP are shown in Table 5<sup>[9]</sup>. The upper normal limit of lateral antebrachial cutaneous SNAP is approximately 60% greater than of medial antebrachial cutaneous SNAP. In the current case, the amplitude of right and left lateral antebrachial cutaneous SNAP (13.5  $\mu$ V and 12.6  $\mu$ V, respectively) was proportionally smaller than of medial antebrachial cutaneous SNAP (12.6  $\mu$ V and 19.9  $\mu$ V, respectively). However, lateral antebrachial cutaneous SNAP has never been compared with medial antebrachial cutaneous SNAP, and appropriate comparison criteria for diagnosis are needed. In addition, the examiner should always remember to correlate the test results in a clinical aspect.

When confirming the diagnosis, excluding other possible differential diagnoses is also necessary. In the present case report, biceps tendon injury or rupture and strain or

Table 5 Reference values of nerve conduction studies<sup>[20]</sup>

Nerve conduction (sensory)	Onset latency (ms)	Peak latency (ms)	Amplitude ( $\mu$ V)
Lateral antebrachial cutaneous	1.6-2.1	2.2-2.6	12-50
Medial antebrachial cutaneous		1.7-2.6	10-30

tear of the biceps or brachialis muscle should have also been considered as differential diagnoses<sup>[9]</sup>. In case of biceps tendon injury or rupture or strain or tear of the biceps or brachialis muscles, sensory changes would not be observed. In the present case, the patient had sensory symptoms in both forearms. Furthermore, hematoma or muscle injury was not observed on the MRI of both arms<sup>[9]</sup>. In addition, cervical radiculopathy involving the fifth or sixth spinal nerve root, should have been considered as a differential diagnosis<sup>[9]</sup>. If the patient had cervical radiculopathy, sensory change would have matched the dermatome of the fifth or sixth spinal nerve root. Furthermore, other muscles supplied by the fifth or sixth spinal nerve root, such as deltoids and supraspinatus muscles, would also have been weak<sup>[20]</sup>. In addition, the patient did not show specific abnormalities on cervical spine MRI. Brachial plexus injury should also have been excluded for definite diagnosis. In general, patients with brachial plexus injury present with broader distributions of sensory change and muscle weakness in upper extremities. Therefore, the differential diagnoses described above were less likely to be applicable to our patient.

The present case report had several limitations. First, the radiologist reported no abnormalities on MRI of the bilateral arm and brachial plexus. MRI is a static diagnostic tool and performing proper nerve tracing based on the patient's movement is difficult. For a more accurate diagnosis, the nerve route could have been traced to detect any entrapment. Second, there is a previous report of a recreational parachutist presenting with bilateral arm weakness, who was later found to have hereditary neuropathy with predisposition to pressure palsies<sup>[21]</sup>. The patient in the present case report was also likely to have hereditary neuropathy with predisposition to pressure palsies. Because the pressure applied during recreational parachuting is greater than the compression during the stretching exercise, we could have recommended testing the *PMP22* gene.

## CONCLUSION

To the best of our knowledge, this is the first reported case of bilateral musculocutaneous neuropathy. Although rare, clinicians should take this disease into consideration when conducting electrodiagnostic studies and be aware the condition may appear bilaterally. Furthermore, people can usually access various exercises and stretching methods through the internet. However, some exercises can cause complications. Clinicians should have adequate understanding of anatomy and give appropriate advice.

## REFERENCES

- 1 Mautner K, Keel JC. Musculocutaneous nerve injury after simulated freefall in a vertical wind-tunnel: a case report. *Arch Phys Med Rehabil* 2007; **88**: 391-393 [PMID: 17321836 DOI: 10.1016/j.apmr.2006.12.018]
- 2 Pećina M, Bojanić I. Musculocutaneous nerve entrapment in the upper arm. *Int Orthop* 1993; **17**: 232-234 [PMID: 8407039 DOI: 10.1007/BF00194185]
- 3 Papanikolaou A, Maris J, Tsampazis K. Isolated musculocutaneous nerve palsy after heavy physical activity. *Injury Extra* 2005; **36**: 486-488 [DOI: 10.1016/j.injury.2005.04.010]
- 4 Yilmaz C, Eskandari MM, Colak M. Traumatic musculocutaneous neuropathy: a case report. *Arch Orthop Trauma Surg* 2005; **125**: 414-416 [PMID: 15843942 DOI: 10.1007/s00402-005-0818-2]
- 5 Merrell CA, Merrell KL. A variation of musculocutaneous neuropathy: implications for electromyographers. *PM R* 2010; **2**: 780-782 [PMID: 20709307 DOI: 10.1016/j.pmrj.2010.04.005]
- 6 Besleaga D, Castellano V, Lutz C, Feinberg JH. Musculocutaneous neuropathy: case report and discussion. *HSS J* 2010; **6**: 112-116 [PMID: 20013159 DOI: 10.1007/s11420-009-9143-6]
- 7 Henry D, Bonthius DJ. Isolated musculocutaneous neuropathy in an adolescent baseball pitcher. *J Child Neurol* 2011; **26**: 1567-1570 [PMID: 21693649 DOI: 10.1177/0883073811411189]
- 8 Chabra T. Case report on isolated musculocutaneous nerve injury following a wrestling match.

- Journal of Dental and Medical Sciences 2015; 14: 97-99 [DOI: [10.9790/0853-14329799](https://doi.org/10.9790/0853-14329799)]
- 9 **Kissel JA**, Leonardelli C. Isolated musculoskeletal neuropathy: a case report. *J Can Chiropr Assoc* 2019; **63**: 162-170 [PMID: [31988537](https://pubmed.ncbi.nlm.nih.gov/31988537/)]
  - 10 **Daniel D**, Machiel JZ. Focal Peripheral Neuropathies. In: Daneil D, Anthony AA, Machiel Z. *Electrodiagnostic Medicine*. 2nd. Philadelphia: Hanley & Belfus, Cop, 2002: 1043
  - 11 **Daniel D**, Machiel JZ, Anthony AA. Peripheral Nervous System's Reaction to Injury. In: Daneil D, Anthony AA, Machiel Z. *Electrodiagnostic Medicine*. 2nd. Philadelphia: Hanley & Belfus, Cop, 2002: 126-129
  - 12 **Mastaglia FL**. Musculoskeletal neuropathy after strenuous physical activity. *Med J Aust* 1986; **145**: 153-154 [PMID: [3016490](https://pubmed.ncbi.nlm.nih.gov/3016490/)]
  - 13 **Lorei MP**, Hershman EB. Peripheral nerve injuries in athletes. Treatment and prevention. *Sports Med* 1993; **16**: 130-147 [PMID: [8378668](https://pubmed.ncbi.nlm.nih.gov/8378668/) DOI: [10.2165/00007256-199316020-00005](https://doi.org/10.2165/00007256-199316020-00005)]
  - 14 **McCarty EC**, Tsairis P, Warren RF. Brachial neuritis. *Clin Orthop Relat Res* 1999; 37-43 [PMID: [10613151](https://pubmed.ncbi.nlm.nih.gov/10613151/)]
  - 15 **Bassett FH 3rd**, Nunley JA. Compression of the musculoskeletal nerve at the elbow. *J Bone Joint Surg Am* 1982; **64**: 1050-1052 [PMID: [7118970](https://pubmed.ncbi.nlm.nih.gov/7118970/)]
  - 16 **Ilayperuma I**, Nanayakkara BG, Hasan R, Uluwitiya SM, Palahepitiya KN. Coracobrachialis muscle: morphology, morphometry and gender differences. *Surg Radiol Anat* 2016; **38**: 335-340 [PMID: [26464302](https://pubmed.ncbi.nlm.nih.gov/26464302/) DOI: [10.1007/s00276-015-1564-y](https://doi.org/10.1007/s00276-015-1564-y)]
  - 17 **Quan D**, Bird S. Nerve Conduction Studies and Electromyography in the Evaluation of Peripheral Nerve Injuries. *The University of Pennsylvania Orthopaedic Journal* 1999; **12**: 45-51
  - 18 **Werner RA**, Albers JW. Relation between needle electromyography and nerve conduction studies in patients with carpal tunnel syndrome. *Arch Phys Med Rehabil* 1995; **76**: 246-249 [PMID: [7717817](https://pubmed.ncbi.nlm.nih.gov/7717817/) DOI: [10.1016/s0003-9993\(95\)80610-5](https://doi.org/10.1016/s0003-9993(95)80610-5)]
  - 19 **Daniel D**, Anthony AA, Macheil JZ. Nerve conduction studies. In: Daneil D, Anthony AA, Machiel Z. *Electrodiagnostic Medicine*. 2nd. Philadelphia: Hanley & Belfus, Cop, 2002: 206-207
  - 20 **Abbed KM**, Coumans JV. Cervical radiculopathy: pathophysiology, presentation, and clinical evaluation. *Neurosurgery* 2007; **60**: S28-S34 [PMID: [17204882](https://pubmed.ncbi.nlm.nih.gov/17204882/) DOI: [10.1227/01.NEU.0000249223.51871.C2](https://doi.org/10.1227/01.NEU.0000249223.51871.C2)]
  - 21 **Marriott M**, Macdonell R, McCrory P. Flail arms in a parachutist: an unusual presentation of hereditary neuropathy with liability to pressure palsies. *Br J Sports Med* 2002; **36**: 465-466 [PMID: [12453844](https://pubmed.ncbi.nlm.nih.gov/12453844/) DOI: [10.1136/bjism.36.6.465](https://doi.org/10.1136/bjism.36.6.465)]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

