



Anti-Rods and Rings Autoantibodies in a Patient With Hepatitis C Virus Infection

Kyung Ho Choi, M.D.¹, Young Ae Lim, M.D.¹, Think-You Kim, M.D.², La-He Jearn, M.D.², Sae Yun Baik, M.D.³,
Sung Won Cho, M.D.⁴, and Eunju Jeong, M.D.⁴

Departments of Laboratory Medicine¹ and Internal Medicine⁴, Ajou University School of Medicine, Suwon; Department of Laboratory Medicine², Hanyang University School of Medicine, Seoul; Green Cross Laboratories³, Yongin, Korea

Dear Editor

The fluorescent antinuclear antibody (FANA) test is widely used for the detection of systemic autoantibodies. Moreover, the fluorescent patterns seen with the FANA test have specific clinical significance [1, 2]. We recently observed an unfamiliar FANA pattern with several small cytoplasmic rod and ring immunofluorescence known as anti-rods and rings (RR) autoantibodies—the first instance in Korea—in a patient with hepatitis C virus (HCV) infection who received combined interferon (IFN) and ribavirin treatment [3-8].

A 69-yr-old man was admitted to Ajou university hospital in Korea in July 2014 for jaundice due to liver cirrhosis. On admission, his condition worsened owing to hepatic failure. He had been diagnosed as having HCV infection, hypertension, and type 2 diabetes in 2001. He refused medical care until 2009, when he received treatment with IFN and ribavirin for one year. The treatment was discontinued because of ineffectiveness. There was no other history of supportive treatment, including herbal medication.

On admission, the patient developed sudden jaundice with the following laboratory results: total bilirubin, 21 mg/dL; direct bilirubin, 15.0 mg/dL; ALT, 78 U/L; and AST, 265 U/L. As there were no previous FANA test data available, the physician re-

quested a FANA test to rule out autoimmune hepatitis as the cause of hepatic failure [7]. The FANA test was performed by using the HEPANA test with HEp-2 cells (Medical & Biological Laboratories Co., Ltd., Nagoya, Japan) and was read with a Zeiss Axio Imager A1 (Carl Zeiss Inc., Germany).

A novel finding on the FANA test included several small cytoplasmic RR immunofluorescences (Fig. 1A). The result was reproducible with a titer of 1:640. Experts at another laboratory confirmed the pattern as anti-RR autoantibodies by using an autoimmune target (AIT) test (ImmunoThink Co., Seoul, Korea). Other reagents, such as HEp-2 ANA slides (INOVA Diagnostics, San Diego, CA, USA) also showed the RR pattern [3-8]. However, Kallestad HEp-2 cell line substrate (Bio-Rad Laboratories, Hercules, CA, USA) showed other cytoplasmic patterns, such as autoantibodies recognizing mitotic spindle apparatus-related antigens, instead of the RR pattern (Fig. 1B). This discrepancy may be due to differences in the conditions used for particular tissue cultures and the fixation of HEp-2 cells. Therefore, all commercial HEp-2 slides may not be reactive to anti-RR autoantibodies [6, 8].

The patient developed acute renal failure likely owing to hepatorenal syndrome at admission and died on day 17 post-admission despite supportive treatment.

Received: January 29, 2015

Revision received: May 8, 2015

Accepted: July 22, 2015

Corresponding author: Young Ae Lim

Department of Laboratory Medicine, Ajou University School of Medicine,
164 World Cup-ro, Yeongtong-gu, Suwon 16499, Korea
Tel: +82-31-219-5786, Fax: +82-31-219-5778
E-mail: limyoung@ajou.ac.kr

© The Korean Society for Laboratory Medicine.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1. (A) Cytoplasmic rods and rings of HEP-2 cells with FITC-conjugated goat anti-human immunoglobulins (Medical & Biological Laboratories Co., Ltd., Nagoya, Japan) in sera from the patient with hepatitis C virus infection ($\times 200$). (B) Cytoplasmic pattern (autoantibodies recognizing mitotic spindle apparatus-related antigens) observed with Kallestad HEP-2 cell line substrate (Bio-Rad Laboratories, Hercules, CA, USA) in sera from the patient with hepatitis C virus infection ($\times 200$).

There have been several reports about the incidence of the anti-RR pattern with hepatitis C and other illnesses. It is present in approximately 30% of the patients with chronic hepatitis C treated with IFN-ribavirin [7], 14.1% with hepatitis C, and 3.4% with hepatitis B but has not been observed in patients without any form of viral hepatitis [8]. High anti-RR autoantibody titers occur in patients with no response or relapsers [3], although some studies found no association between anti-RR titer and treatment response [7, 8]. The patient's anti-RR titer was 1:640, which was high according to earlier reports [7, 8]. The patient did not respond to combination IFN and ribavirin therapy.

Anti-RR autoantibodies are reportedly detected more often in relapsers and non-responders than in sustained virological responders defined as having undetectable HCV RNA 24 weeks post-therapy [3, 4]. The viral load of the present patient, 196,254 copies/mL (72,687 IU/mL), exceeded the category limits of high viral load [7]. We detected no hepatitis B virus co-infection or other autoantibodies, including anti-mitochondrial antibody or anti-smooth muscle antibody. This patient's treatment with IFN and ribavirin was discontinued after a year owing to lack of effectiveness; hence, the patient appeared to be a non-responder [7].

We did not analyze the target antigens of anti-RR autoantibodies; several studies have reported that the target antigens of anti-RR autoantibodies are inosine monophosphate dehydrogenase 2 (IMPDH2), cytidine triphosphate synthase 1 (CTPS1), or both [3-5, 8]. These enzymes, which normally mask epitopes

that might participate in RR formation, are inhibited by IFN and ribavirin treatment, which induces structural modifications that make the epitopes accessible to the immune system [5, 8]. Other potential target antigens, including Myc-associated zinc finger protein (MAZI), have also been reported and must be confirmed [6].

Anti-RR autoantibodies can easily be overlooked if clinical pathologists ignore specific FANA patterns; therefore, FANA patterns require cautious interpretation. The anti-RR autoantibody pattern should also be distinguished from other cytoplasmic patterns and reported as the cytoplasmic RR pattern. Clinical information on patient history and present illness is helpful in the interpretation of unusual or unfamiliar FANA patterns.

Authors' Disclosures of Potential Conflicts of Interest

No potential conflicts of interest relevant to this article were reported.

REFERENCES

1. Vermeersch P and Bossuyt X. Prevalence and clinical significance of rare antinuclear antibody patterns. *Autoimmun Rev* 2013;12:998-1003.
2. Liberal R, Mieli-Vergani G, Vergani D. Clinical significance of autoantibodies in autoimmune hepatitis. *J Autoimmun* 2013;46:17-24.
3. Carcamo WC, Ceribelli A, Calise SJ, Krueger C, Liu C, Daves M, et al.

- Differential reactivity to IMPDH2 by anti-rods/rings autoantibodies and unresponsiveness to pegylated interferon-alpha/ribavirin therapy in US and Italian HCV patients. *J Clin Immunol* 2013;33:420-6.
4. Covini G, Carcamo WC, Bredi E, von Mühlen CA, Colombo M, Chan EK. Cytoplasmic rods and rings autoantibodies developed during pegylated interferon and ribavirin therapy in patients with chronic hepatitis C. *Antivir Ther* 2012;17:805-11.
 5. Carcamo WC, Satoh M, Kasahara H, Terada N, Hamazaki T, Chan JY, et al. Induction of cytoplasmic rods and rings structures by inhibition of the CTP and GTP synthetic pathway in mammalian cells. *PLoS One* 2011;6:e29690.
 6. Stinton LM, Myers RP, Coffin CS, Fritzier MJ. Clinical associations and potential novel antigenic targets of autoantibodies directed against rods and rings in chronic hepatitis C infection. *BMC Gastroenterol* 2013; 13:50.
 7. Novembrino C, Aghemo A, Ferraris Fusarini C, Maiavacca R, Martinato C, Lunghi G, et al. Interferon-ribavirin therapy induces serum antibodies determining 'rods and rings' pattern in hepatitis C patients. *J Viral Hepat* 2014;21:944-9.
 8. Keppeke GD, Nunes E, Ferraz ML, Silva EA, Granato C, Chan EK, Andrade LE. Longitudinal study of a human drug-induced model of autoantibody to cytoplasmic rods/rings following HCV therapy with ribavirin and interferon- α . *PLoS One* 2012;7:e45392.