

# Association of genetic variation in *FTO* with risk of obesity and type 2 diabetes with data from 96,551 East and South Asians

H. Li · T. O. Kilpeläinen · C. Liu · J. Zhu · Y. Liu · C. Hu · Z. Yang · W. Zhang · W. Bao · S. Cha · Y. Wu · T. Yang · A. Sekine · B. Y. Choi · C. S. Yajnik · D. Zhou · F. Takeuchi · K. Yamamoto · J. C. Chan · K. R. Mani · L. F. Been · M. Imamura · E. Nakashima · N. Lee · T. Fujisawa · S. Karasawa · W. Wen · C. V. Joglekar · W. Lu · Y. Chang · Y. Xiang · Y. Gao · S. Liu · Y. Song · S. H. Kwak · H. D. Shin · K. S. Park · C. H. D. Fall · J. Y. Kim · P. C. Sham · K. S. L. Lam · W. Zheng · X. Shu · H. Deng · H. Ikegami · G. V. Krishnaveni · D. K. Sanghera · L. Chuang · L. Liu · R. Hu · Y. Kim · M. Daimon · K. Hotta · W. Jia · J. S. Kooner · J. C. Chambers · G. R. Chandak · R. C. Ma · S. Maeda · R. Dorajoo · M. Yokota · R. Takayanagi · N. Kato · X. Lin · R. J. F. Loos

Received: 24 August 2011 / Accepted: 10 October 2011 / Published online: 23 November 2011  
© The Author(s) 2011. This article is published with open access at Springerlink.com

## Abstract

**Aims/hypothesis** *FTO* harbours the strongest known obesity-susceptibility locus in Europeans. While there is growing evidence for a role for *FTO* in obesity risk in Asians, its association with type 2 diabetes, independently of BMI, remains inconsistent. To test whether there is an association of the *FTO* locus with obesity and type 2 diabetes, we conducted a meta-analysis of 32 populations including 96,551 East and South Asians.

**Methods** All studies published on the association between *FTO*-rs9939609 (or proxy [ $r^2 > 0.98$ ]) and BMI, obesity or type 2 diabetes in East or South Asians were invited. Each study group analysed their data according to a standardised analysis plan. Association with type 2 diabetes was also adjusted for BMI. Random-effects meta-analyses were performed to pool all effect sizes.

**Results** The *FTO*-rs9939609 minor allele increased risk of obesity by 1.25-fold/allele ( $p = 9.0 \times 10^{-19}$ ), overweight by

**Electronic supplementary material** The online version of this article (doi:10.1007/s00125-011-2370-7) contains peer-reviewed but unedited supplementary material, which is available to authorised users.

H. Li · C. Liu · J. Zhu · D. Zhou · X. Lin (✉)  
Institute for Nutritional Sciences,  
Shanghai Institutes for Biological Sciences,  
Chinese Academy of Sciences,  
294 Tai-Yuan Road,  
Shanghai 200031, People's Republic of China  
e-mail: xlin@sibs.ac.cn

T. O. Kilpeläinen · R. J. F. Loos (✉)  
MRC Epidemiology Unit, Institute of Metabolic Science Box 285,  
Addenbrooke's Hospital,  
Hills Road,  
Cambridge CB2 0QQ, UK  
e-mail: ruth.loos@mrc-epid.cam.ac.uk

Y. Liu  
Institutes of Biomedical Sciences, Fudan University,  
Shanghai, People's Republic of China

C. Hu · W. Jia  
Shanghai Diabetes Institute,  
Department of Endocrinology and Metabolism,  
Shanghai Clinical Center of Diabetes,  
Shanghai Jiao Tong University Affiliated Sixth People's Hospital,  
Shanghai, People's Republic of China

Z. Yang · R. Hu  
Department of Endocrinology and Metabolism, Huashan Hospital,  
Institute of Endocrinology and Diabetology at Fudan University,  
Shanghai Medical School, Fudan University,  
Shanghai, People's Republic of China

W. Zhang · J. C. Chambers  
Department Epidemiology and Biostatistics,  
School of Public Health, Imperial College London,  
London, UK

1.13-fold/allele ( $p=1.0\times 10^{-11}$ ) and type 2 diabetes by 1.15-fold/allele ( $p=5.5\times 10^{-8}$ ). The association with type 2 diabetes was attenuated after adjustment for BMI (OR 1.10-fold/allele,  $p=6.6\times 10^{-5}$ ). The *FTO*-rs9939609 minor allele increased BMI by 0.26 kg/m<sup>2</sup> per allele ( $p=2.8\times 10^{-17}$ ), WHR by 0.003/allele ( $p=1.2\times 10^{-6}$ ), and body fat percentage by 0.31%/allele ( $p=0.0005$ ). Associations were similar using dominant models. While the minor allele is less common in East Asians (12–20%) than South Asians (30–33%), the effect of *FTO* variation on obesity-related traits and type 2 diabetes was similar in the two populations.

**Conclusions/interpretation** *FTO* is associated with increased risk of obesity and type 2 diabetes, with effect sizes similar in East and South Asians and similar to those observed in Europeans. Furthermore, *FTO* is also associated with type 2 diabetes independently of BMI.

W. Bao · L. Liu

Department of Nutrition and Food Hygiene and MOE Key Lab of Environment and Health, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, People's Republic of China

S. Cha · J. Y. Kim

Division of Constitutional Medicine Research, Korea Institute of Oriental Medicine, Daejeon, South Korea

Y. Wu

Department of Genetics, University of North Carolina, Chapel Hill, NC, USA

T. Yang

Key Laboratory of Biomedical Information Engineering of Ministry of Education, and Institute of Molecular Genetics, School of Life Science and Technology, Xi'an Jiaotong University, Xi'an, People's Republic of China

A. Sekine · K. Hotta

EBM Research Center, Kyoto University Graduate School of Medicine, Kyoto, Japan

B. Y. Choi

Department of Preventive Medicine, HanYang University College of Medicine, Seoul, South Korea

C. S. Yajnik · C. V. Joglekar

Diabetology Research Centre, KEM Hospital and Research Centre, Pune, India

F. Takeuchi · N. Kato

National Center for Global Health and Medicine, Tokyo, Japan

K. Yamamoto

Division of Genome Analysis, Medical Institute of Bioregulation, Kyushu University, Fukuoka, Japan

**Keywords** Asians · *FTO* · Meta-analysis · Obesity · Type 2 diabetes

### Abbreviations

GWAS Genome-wide association study  
MAF Minor allele frequency  
PAR Population-attributable risk  
SNP Single-nucleotide polymorphism

### Introduction

Large-scale genome-wide association studies (GWAS) in mainly white Europeans have identified at least 50 genetic loci to be robustly associated with obesity-related traits [1–

J. C. Chan · R. C. Ma

Department of Medicine and Therapeutics, Chinese University of Hong Kong, Prince of Wales Hospital, Shatin, Hong Kong Special Administrative Region, People's Republic of China

K. R. Mani · G. R. Chandak

Centre for Cellular and Molecular Biology (CCMB-CSIR), Hyderabad, India

L. F. Been · D. K. Sanghera

University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA

M. Imamura · S. Maeda

Laboratory for Endocrinology and Metabolism, RIKEN Center for Genomic Medicine, Yokohama, Japan

E. Nakashima

Department of Diabetes and Endocrinology, Chubu Rosai Hospital, Nagoya, Japan

N. Lee

USC Office of Population Studies Foundation, University of San Carlos, Cebu, Philippines

T. Fujisawa

Department of Geriatric Medicine and Nephrology, Osaka University Graduate School of Medicine, Suita, Japan

S. Karasawa · M. Daimon

Third Department of Internal Medicine, and Global Center of Excellence Program Study Group, Yamagata University School of Medicine, Yamagata, Japan

W. Wen · W. Zheng · X. Shu

Division of Epidemiology, Department of Medicine, Vanderbilt School of Medicine, Nashville, TN, USA

12]. A cluster of common variants in the first intron of the fat mass and obesity-associated gene (*FTO*) was the first obesity-susceptibility locus to be identified by two independent GWAS in 2007 [1, 2] and has since been consistently replicated by many others and for a variety of obesity-related traits [7, 9, 13–15]. Of all currently identified obesity-susceptibility loci, the *FTO* locus has the most pronounced effect on BMI and obesity risk, at least in individuals of European descent. Each minor allele of any commonly investigated variant in *FTO* increases BMI by 0.30–0.40 kg/m<sup>2</sup> (equivalent to 870–1,150 g for a person 1.7 m tall) and risk of obesity by ~20% [7, 15]. The

minor allele of the *FTO* variant is common (minor allele frequency (MAF)~42%) in white Europeans, such that 66% of Europeans carry at least one risk allele and 18% carry two risk alleles. Because of the high prevalence of the risk allele and its relatively strong effect on BMI, the *FTO* locus explains most (0.34%), yet little, of the variation in BMI in Europeans [7].

*FTO* has also been examined as an obesity-susceptibility locus in populations of non-white European origin. While the initial replication efforts in East Asian populations were inconsistent [16, 17], a growing number of studies have provided evidence that genetic variation in *FTO* influences

W. Lu

Shanghai Institute of Preventive Medicine,  
Shanghai, People's Republic of China

Y. Chang

National Taiwan University Hospital Bei-Hu branch,  
Taipei, Taiwan

Y. Xiang · Y. Gao

Department of Epidemiology, Shanghai Cancer Institute,  
Shanghai, People's Republic of China

S. Liu

Center for Metabolic Disease Prevention, School of Public Health  
and David Geffen School of Medicine, UCLA,  
Los Angeles, CA, USA

Y. Song

Division of Preventive Medicine, Brigham & Women's Hospital,  
Harvard Medical School,  
Boston, MA, USA

S. H. Kwak · K. S. Park

Department of Internal Medicine, Seoul National University  
College of Medicine,  
Seoul, South Korea

H. D. Shin

Department of Life Science, Sogang University,  
Seoul, South Korea

C. H. D. Fall

MRC Lifecourse Epidemiology Unit, University of Southampton,  
Southampton General Hospital,  
Southampton, Hampshire, UK

P. C. Sham · K. S. L. Lam

Li Ka Shing Faculty of Medicine, University of Hong Kong,  
Hong Kong, Hong Kong Special Administrative Region,  
People's Republic of China

H. Deng

School of Medicine, University of Missouri,  
Kansas City, MO, USA

H. Deng

Center of Systematic Biomedical Research, University of  
Shanghai for Science and Technology,  
Shanghai, People's Republic of China

H. Deng

Institute of Bioscience and Biotechnology, School of Science,  
Beijing Jiaotong University,  
Beijing, People's Republic of China

H. Ikegami

Department of Endocrinology, Metabolism and Diabetes,  
Kinki University School of Medicine,  
Osaka, Japan

G. V. Krishnaveni

Epidemiology Research Unit, Holdsworth Memorial Hospital,  
Mysore, India

L. Chuang

Department of Internal Medicine,  
National Taiwan University Hospital,  
Taipei, Taiwan

Y. Kim

Department of Preventive Medicine,  
Dong-A University College of Medicine,  
Busan, South Korea

J. S. Kooner

National Heart & Lung Institute, Hammersmith Hospital,  
Hammersmith Campus, Faculty of Medicine,  
Imperial College London,  
London, UK

R. Dorajoo

Genome Institute of Singapore, Agency for Science,  
Technology and Research,  
Singapore, Republic of Singapore

R. Dorajoo

Department of Genomics of Common Disease, School of Public  
Health, Hammersmith Hospital, Imperial College London,  
London, UK

M. Yokota

Department of Genome Science, School of Dentistry,  
Aichi-Gakuin University,  
Nagoya, Japan

R. Takayanagi

Department of Medicine and Bioregulatory Science,  
Graduate School of Medical Sciences, Kyushu University,  
Fukuoka, Japan

BMI and obesity risk also in Chinese, Japanese, Korean and Filipino populations [18–27]. A GWAS for BMI in 7,861 Koreans identified variation in *FTO* (rs9939609) as the most significantly associated locus, nearly reaching genome-wide significance ( $p=1.5\times 10^{-7}$ ) [28]. Furthermore, literature-based meta-analyses in Asians reported that the minor allele for the rs9939609 *FTO* single-nucleotide polymorphism (SNP) significantly ( $p=9\times 10^{-9}$ ) increased the risk of obesity, but no other obesity-related traits were examined [18, 29, 30]. Fewer studies in South Asians have been reported, two of which confirmed the association between the *FTO* locus and obesity susceptibility [31, 32], whereas one did not [33]. The prevalence of the risk allele in East Asians (~20%) and South Asians (~30%) is substantially lower than in Europeans, and the reported effect sizes in both East and South Asians vary widely for BMI (OR 0.13–0.83 kg/m<sup>2</sup> per minor allele) and obesity risk (OR 1.02–1.48 per minor allele) [16, 18, 20–25, 27, 34–39].

*FTO* was first identified as a type 2 diabetes-susceptibility gene, but, as further adjustment for BMI abolished the association with type 2 diabetes [1], it was suggested that *FTO* is primarily an obesity-susceptibility locus. However, the BMI-independent role of *FTO* in type 2 diabetes remains a matter of debate, particularly in Asians but also in white Europeans. While several studies have reported that the association between the *FTO* locus and risk of type 2 diabetes remained significant after adjustment for BMI [15, 18, 33, 35, 40, 41], others could not confirm this [21, 30, 32, 37, 42].

To firmly establish the association between the *FTO* locus and obesity susceptibility in East and South Asians and to assess its effect size and potential heterogeneity across Asian populations, we performed a systematic meta-analysis of data from 32 populations, including 96,551 men and women, using standardised study-specific association analyses. Furthermore, we examined whether the *FTO* locus is associated with type 2 diabetes independently of its association with BMI.

## Methods

**Literature search and study identification** We designed a meta-analysis based on de novo analyses of data according to a standardised plan to achieve the greatest consistency possible across studies. We identified all published studies (before September 2010) that had examined the association of genetic variation in *FTO* with risk of obesity and type 2 diabetes and with obesity-related continuous traits in East and South Asian adults (age  $\geq 18$  years) by a PubMed literature search using the key words ‘*FTO*’, ‘fat mass and obesity associated gene’ and ‘genome-wide association study’. References from the identified papers were subse-

quently screened to identify additional studies and to ensure that the list of eligible studies was complete. The literature search was carried out by two investigators independently, who cross-checked their search results for completeness.

Our literature search identified 38 publications, one of which was excluded because it was a subsample of another identified study. We invited the corresponding authors of the remaining 37 publications to join our meta-analysis, of which 26 agreed to participate and eventually 22 submitted raw data or summary statistics. We also included a Korean population with previously unpublished data (Y. M. Kim, J. Shin, C.B. Lee, M.K. Kim, Y. Tabara, T. Miki and B.Y. Choi), which was presented by a contributing author.

Taken together, our meta-analysis included data for 31 populations from 22 publications and one unpublished study, with 96,551 individuals altogether. The study identification and selection process is illustrated in Fig. 1.


All studies were conducted according to the Declaration of Helsinki. Informed consent was obtained from all participants, and the studies were approved by the ethics committees of the participating institutions.

**Genotyping** The rs9939609 *FTO* SNP was examined in 18 studies, whereas proxy SNPs were used in 14 studies. More specifically, the rs8050135 SNP was genotyped in 11 studies of East Asians and one of South Asians, and the rs3751812 and rs17817449 SNP were each genotyped once in studies of East Asians (electronic supplementary material [ESM] Table 1). The linkage disequilibrium between rs9939609 and the three proxies (rs8050135, 3751812, rs17817449) is perfect ( $r^2=1$ ) in populations of East Asian origin, based on CHB+JPT data from the HapMap (Rel 24/Phase II). The linkage disequilibrium between rs9939609 and rs8050135 in Indian Asians is very high ( $r^2>0.98$ ), based on a subsample ( $n=305$ ) of the participating Lolipop study.

The genotyping success rate and concordance rate were  $>95\%$ , and genotype distributions were in Hardy–Weinberg equilibrium ( $p>0.01$ ) in all participating studies (ESM Table 1).

**Statistical analysis** As case–control definitions and statistical analyses used in the published papers were inconsistent, we asked analysts of each of the participating cohorts to re-analyse their data according to a standardised analysis plan. Summary statistics of each study were subsequently meta-analysed.

**Obesity-susceptibility traits and type 2 diabetes** Overweight was defined as a BMI  $\geq 24$  kg/m<sup>2</sup>, and obesity as a BMI  $\geq 28$  kg/m<sup>2</sup> according to the definition proposed by the Working Group on Obesity in China [43]. Anthropometric data, including weight, height, waist circumference, hip

**Fig. 1** Study identification and inclusion in the meta-analyses

circumference and body fat percentage, were collected in each study as described previously (ESM Table 1), BMI was calculated as weight (kg) divided by height squared ( $m^2$ ), and WHR as waist circumference (cm) divided by hip circumference (cm). Raw data were used for analyses.

Type 2 diabetes was defined as meeting one or more of the following criteria: (1) fasting glucose  $\geq 7.0$  mmol/l; (2) 2-h glucose  $\geq 11.1$  mmol/l; (3) previous diagnosis of type 2 diabetes; (4)  $HbA_{1c} \geq 6.5\%$  (48 mmol/mol); (5) self-reported type 2 diabetes (ESM Table 1).

**Study-specific de novo data analyses** Association analyses within each study were performed for the total population and for men and women separately using additive and dominant genetic models. The associations of *FTO*-rs9939609 (or proxy) with risk of obesity and type 2 diabetes were assessed with multiple logistic regression models. Generalised linear models were used to assess the associations of *FTO*-rs9939609 (or proxy) with obesity-related continuous traits. In studies with a case–control design, analyses for continuous traits were conducted in control samples only. All analyses were adjusted for age and sex (sex-stratified analyses were only adjusted for age). The association with type 2 diabetes was also analysed with adjustment for BMI. Adjustments were performed by

including the covariates (age, sex and/or BMI) as a linear term in the association model.

Summary statistics from the study-specific association analyses were reported in a standardised Excel form by the analysts of each study and collected centrally for meta-analyses.

**Meta-analyses** Data extraction from the forms and meta-analyses was performed independently by two investigators and cross-checked for consistency. All ambiguities were clarified with the respective analysts before the final meta-analyses.

ORs and beta coefficients from the individual studies were pooled using DerSimonian and Laird random-effects meta-analyses [44]. Meta-analyses were performed of all studies combined. Because of differences in genetic background as well as in susceptibility to obesity and type 2 diabetes, meta-analyses were also stratified by East Asian and South Asian origin of the populations. Furthermore, East Asians were further stratified according to their country of origin.

Between-study heterogeneity was tested by Cochrane's Q test and quantified by the  $I^2$  index.  $I^2$  values of  $<25\%$ ,  $25\text{--}75\%$  and  $>75\%$  were defined as low, moderate and high heterogeneity, respectively [45]. To examine the sources of



heterogeneity in our meta-analyses, we performed random-effects meta-regressions, where the between-study variance was estimated with the restricted maximum likelihood approach. Meta-regressions included the following study-specific variables as covariates: year of publication, country of origin, sample size, study design, mean age and mean BMI.

A funnel plot, along with Begg's and Egger's tests, was used to test for the presence of publication bias.

Statistical analyses were performed with the Stata 9.0 software (StataCorp LP, College Station, TX, USA). Meta-analyses and meta-regressions were implemented by the *metan* and *metareg* commands of Stata, respectively.  $p < 0.05$  was considered to be significant, except for Cochrane's Q test for heterogeneity and Begg's and Egger's tests for publication bias, where a level of  $p < 0.10$  was used.

The variation in obesity-related continuous traits explained by the *FTO* variant was evaluated using the equation  $2f(1-f)a^2$ , where  $f$  is the frequency of the variant and  $a$  is its additive standardised effect [5]. Population-attributable risk (PAR) was calculated as  $PAR = (X - 1)/X$ . Assuming a multiplicative model,  $X = (1 - f)^2 + 2f(1 - f)\gamma + f^2\gamma^2$ , where  $\gamma$  is the estimated OR, and  $f$  is the frequency of risk allele [46].

## Results

*Characteristics of populations included in the meta-analyses* Analyses were conducted in Chinese Hans (China Mainland:  $n=10$ ; Singapore:  $n=2$ ), Japanese ( $n=7$ ), Indians ( $n=7$ ), Koreans ( $n=4$ ), Singapore Malays ( $n=1$ ) and Filipinos ( $n=1$ ; Table 1). Fifteen of the populations were case-control designed for obesity ( $n=3$ ) or type 2 diabetes ( $n=8$ ) or both ( $n=4$ ), whereas 17 populations were population-based. The mean age and BMI of the populations ranged from 27.9 to 66.8 years and from 20.5 to 27.1 kg/m<sup>2</sup>, respectively. The prevalence in population-based studies ranged from 3.1% to 37.9% for obesity and from 2.9% to 41.9% for type 2 diabetes.

The MAF of *FTO*-rs9939609 (or proxy) is 12–14% in Chinese Hans and Koreans, 18–20% in Japanese and Filipinos, and 30–33% in Singapore Malays and Indians (Table 1).

*Associations with obesity and overweight* A total of 24 populations ( $n_{\text{obese}}=13,032$ ;  $n_{\text{overweight}}=22,474$ ;  $n_{\text{normalweight}}=35,767$ ) were available for meta-analyses of the association between the *FTO* variant and risk of obesity and overweight.

Each additional *FTO*-rs9939609 minor (A) allele increased the odds of obesity by 1.25 ( $p=9.0 \times 10^{-19}$ ) compared with normal weight individuals (Fig. 2), and by 1.17 ( $p=7.4 \times 10^{-11}$ ) compared with non-obese individuals

(ESM Fig. 1). Each additional minor allele increased the odds of overweight by 1.13 ( $p=1.0 \times 10^{-11}$ ; ESM Fig. 2). The odds of obesity and overweight were the same in both East Asian and South Asian populations ( $p=0.18$  and 0.84, respectively; ESM Table 2). Associations were similar in men and women (ESM Table 3). The heterogeneity across all studies was low ( $13\% \leq I^2 \leq 19\%$ ).

When a dominant genetic model was used, the odds were only slightly higher than for the additive genetic model (ESM Table 4).

*Association with type 2 diabetes* In our meta-analysis of 22 populations ( $n_{\text{cases}}=33,744$ ,  $n_{\text{controls}}=43,549$ ), each additional *FTO*-rs9939609 minor allele increased the odds of type 2 diabetes by 1.15 ( $p=5.5 \times 10^{-8}$ ) when adjusted for age and sex (Fig. 3). Further adjustment for BMI attenuated, but did not abolish, the association with type 2 diabetes (OR 1.10,  $p=6.6 \times 10^{-5}$ ) (Fig. 4). Results were similar in East Asians and South Asians (ESM Table 2), in men and women (ESM Table 3), and when a dominant model was used (ESM Table 4).

The association results across all studies showed moderate heterogeneity ( $44\% \leq I^2 \leq 48\%$ ; Figs 3 and 4). Meta-regression analyses revealed that the difference in study design contributed to some of the heterogeneity. Subsequent subgroup analyses showed that the association with type 2 diabetes was more pronounced in studies with a case-control design (OR [95% CI]=1.19 [1.14, 1.23],  $p=3.7 \times 10^{-19}$ ,  $I^2=0.0\%$ ) than in cohort studies (OR [95% CI]=1.09 [0.99, 1.20],  $p=0.07$ ,  $I^2=54.4\%$ ), which showed moderate heterogeneity (ESM Table 5).

*Associations with obesity-related continuous traits* The meta-analyses of the association of *FTO*-rs9939609 with BMI, waist circumference, hip circumference, WHR and body fat percentage included 30 ( $n=71,022$ ), 22 ( $n=51,543$ ), 20 ( $n=48,508$ ), 20 ( $n=48,508$ ) and nine ( $n=19,580$ ) populations, respectively.

Each additional *FTO*-rs9939609 minor allele was associated with a 0.26 kg/m<sup>2</sup> higher BMI ( $p=2.8 \times 10^{-17}$ ; equivalent to ~750 g/allele for a person 1.7 m tall) (Fig. 5), 0.51 cm larger waist circumference ( $p=3.0 \times 10^{-9}$ ) (ESM Fig. 3), 0.36 cm larger hip circumference ( $p=0.0003$ ) (ESM Fig. 4), 0.003 greater WHR ( $p=1.2 \times 10^{-6}$ ; ESM Fig. 5), and 0.31% higher body fat percentage ( $p=0.0005$ ) (ESM Fig. 6). All associations were very similar between East and South Asians (ESM Table 2), between men and women (ESM Table 3), or when a dominant genetic model was used (ESM Table 4).

We observed moderate heterogeneity across studies in the associations with BMI and hip circumference (BMI:  $I^2=33\%$ ; hip circumference:  $I^2=51\%$ ; Fig. 5; ESM Fig. 4). Meta-regression suggested that, for BMI, the heterogeneity was

**Table 1** Descriptive information of studies included in the meta-analyses, sorted by ethnicity, study design and publication year

Paper	Study	Publication year	Ethnicity	Country	Study design	Sample size			NFG	QT analyses	Mean age (years)	Mean BMI (kg/m <sup>2</sup> )	FTO SNP	MAF
						Obese	OW	NW						
Li et al. [16]	NHAPC	2008	East Asian	China	Population based	472	1,215	1,503	423	1,893	3,190	rs9939609	0.11	
Sha et al. [55]	GSBC	2009	East Asian	China	Population based	78	326	1,223	n.a.	n.a.	1,627	rs9939609	0.12	
Hu et al. [56]	SHDS	2009	East Asian	China	Case-control <sup>b</sup>	n.a.	n.a.	n.a.	1,759	1,791	1,791	rs8050136	0.12	
Li et al. [35]	WDS	2010	East Asian	China	Case-control <sup>b, c</sup>	243	976	1,368	877	1,405	1,405	rs9939609	0.12	
Cheung et al. [24]	CRISPS	2010	East Asian	China	Case-control <sup>b</sup>	419	n.a.	691	n.a.	n.a.	691	rs8050136	0.12	
Liu et al. [18]	n.a.	2010	East Asian	China	Case-control <sup>b, c</sup>	277	794	893	1,767	1,961	1,961	rs9939609	0.12	
Ng et al. [21]	CUHK	2010	East Asian	China	Case-control <sup>b, c</sup>	1,147	2,293	2,432	5,872	583	583	rs3751812	0.12	
Shu et al. [42]	SGWAS	2010	East Asian	China	Case-control <sup>b</sup>	n.a.	n.a.	n.a.	1,043	2,170	2,170	rs9939609	0.12	
Wen et al. [57]	FLSGS	2010	East Asian	China	Case-control <sup>b</sup>	n.a.	n.a.	n.a.	1,160	1,127	1,127	rs8050136	0.12	
Chang et al. [23]	NTUH	2008	East Asian	Taiwan	Case-control <sup>b, c</sup>	737	677	719	881	1,254	1,254	rs9939609	0.14	
Cha et al. [25]	Kirin	2008	East Asian	Korea	Population based	252	304	361	n.a.	n.a.	917	rs17817449	0.14	
Cha et al. [58]	KCMS	2009	East Asian	Korea	Population based	61	261	688	n.a.	n.a.	1,010	rs8050136	0.12	
Kim et al. (unpublished data)	YangPyeong Cardiovascular Cohort Study		East Asian	Korea	Population based	339	995	1,092	194	2,061	2,426	rs9939609	0.12	
Ng et al. [34]	Korea SNUH	2008	East Asian	Korea	Case-control <sup>b</sup>	n.a.	n.a.	n.a.	758	629	629	rs8050136	0.12	
Takeuchi et al. [59]	CAGE-Amagasaki	2009	East Asian	Japan	Population based	388	1,562	3,719	n.a.	n.a.	5,660	rs9939609	0.19	
Takeuchi et al. [59]	CAGE-Fukuoka	2009	East Asian	Japan	Population based	721	3,763	8,076	n.a.	n.a.	12,560	rs9939609	0.19	
Takeuchi et al. [59]	CAGE-BMI	2009	East Asian	Japan	Population based	168	607	1,006	n.a.	n.a.	1,781	rs9939609	0.20	
Karasawa et al. [19]	Takahata	2010	East Asian	Japan	Population based	220	886	1,533	215	2,306	2,639	rs9939609	0.20	
Hotta et al. [20]	GWASJPN obesity	2008	East Asian	Japan	Case-control <sup>a</sup>	1,559	n.a.	1,541	n.a.	n.a.	1,541	rs9939609	0.18	
Omori et al. [37]	RIKEN T2D	2008	East Asian	Japan	Case-control <sup>b</sup>	n.a.	n.a.	n.a.	4,584	2,262	2,262	rs8050136	0.20	
Takeuchi et al. [59]	CAGE-T2DM	2009	East Asian	Japan	Case-control <sup>b</sup>	n.a.	n.a.	n.a.	6,781	7,307	n.a.	rs9939609	0.19	
Marville et al. [27]	CLHNS	2008	East Asian	Philippines	Population based	321	560	836	155	1,463	1,717	rs9939609	0.18	
Tan et al. [22]	SP2	2008	East Asian	Singapore (Chinese)	Population based	195	624	1,609	145	2,248	2,430	rs8050136	0.12	
Tan et al. [22]	SiMES	2008	East Asian	Singapore (Malays)	Population based	848	826	846	787	1,248	2,520	rs8050136	0.30	
Tan et al. [22]	SDCS	2008	East Asian	Singapore (Chinese)	Case-control <sup>f</sup>	426	809	757	n.a.	n.a.	n.a.	rs8050136	0.14	
Chambers et al. [6]	LOLIPOP (IA317)	2008	South Asian	India	Population based	727	858	536	434	1,651	2,247	rs8050136	0.33	
Chambers et al. [6]	LOLIPOP (IA610)	2008	South Asian	India	Population based	2,479	2,647	1,423	1,780	4,715	7,060	rs8050136	0.32	
Tan et al. [22]	SINDI	2008	South Asian	India	Population based	760	910	858	974	1,348	2,528	rs8050136	0.33	

Table 1 (continued)

Paper	Study	Publication year	Ethnicity	Country	Study design	Sample size				Mean age (years)	Mean BMI (kg/m <sup>2</sup> )	FTO SNP	MAF
						Obese	OW	NW	T2DM				
Yajnik et al. [33]	Parthenon	2009	South Asian	India	Population based	136	320	511	n.a.	967	23.76	rs9939609	0.33
Yajnik et al. [33]	PMNS	2009	South Asian	India	Population based	59	271	1,546	50	1,681	20.83	rs9939609	0.31
Sanghera et al. [40]	Sikh Diabetes Study	2008	South Asian	India	Case-control <sup>b</sup>	n.a.	n.a.	n.a.	1,138	765	26.25	rs9939609	0.31
Yajnik et al. [33]	WELLGEN	2009	South Asian	India	Case-control <sup>b</sup>	n.a.	n.a.	n.a.	1,967	1,681	20.50	rs9939609	0.31

Individuals from CAGE-T2DM study were selected from other three CAGE population-based studies


<sup>a</sup> Obese case-control study

<sup>b</sup> T2DM case-control study

<sup>c</sup> Obese case-control study conducted in T2DM cases

n.a., data not available or not used in meta-analysis; NFG, normal fasting glucose; NW, normal weight; OW, overweight; QT, quantitative trait; T2DM, type 2 diabetes





**Fig. 2** Association of *FTO*-rs9939609 (or proxy) with obesity. Study-specific association analyses assumed an additive genetic model, comparing obese with normal-weight individuals, adjusted for age and

sex. Effect sizes were combined using random-effects meta-analyses (DerSimonian–Laird method)

mainly due to difference in mean age and mean BMI among different populations. For hip circumference, the heterogeneity was mainly attributed to difference in mean BMI, i.e. the effect of the *FTO* minor allele tended to be larger in populations with a mean BMI  $\geq 24$  kg/m<sup>2</sup>, compared with those with a mean BMI  $< 24$  kg/m<sup>2</sup>.


*FTO*-rs9939609 explained 0.16% and 0.20% of the inter-individual variation in BMI in East and South Asian populations, respectively. The proportion of variation in other obesity-related continuous traits explained by *FTO*-rs9939609 was  $< 0.10\%$  (ESM Table 2).

**Publication bias** The funnel plots for the associations with obesity, type 2 diabetes, waist circumference, WHR and body fat percentage were symmetrical and the results for Begg’s and Egger’s tests were non-significant ( $p \geq 0.10$ ),

indicating that our results were not affected by publication bias (ESM Fig. 7). However, there was some evidence of publication bias and/or genetic heterogeneity for BMI (Begg’s test,  $p=0.08$ ; Egger’s test,  $p=0.07$ ) and hip circumference (Begg’s test,  $p=0.03$ ; Egger’s test,  $p=0.08$ ; ESM Fig. 7).

**Discussion**

This meta-analysis, combining data of 96,551 Asians from 32 populations, further confirms that genetic variation in *FTO* is associated with increased risk of obesity in East and South Asians. Despite differences in genetic background and obesity susceptibility between East and South Asians, the effect of *FTO* on obesity and related traits was generally



**Fig. 3** Association of *FTO*-rs939609 (or proxy) with type 2 diabetes. Study-specific association analyses assumed an additive genetic model adjusted for age and sex. Effect sizes were combined using random-effects meta-analyses (DerSimonian–Laird method)


similar to, or only somewhat smaller than, those reported for white Europeans. We furthermore confirm that variation in *FTO* is associated with increased risk of type 2 diabetes, an association that, unlike in white Europeans, is not abolished after adjustment for BMI in both East and South Asians.

Large-scale studies in individuals of white European descent have reported that each additional *FTO* minor allele increases the odds of obesity by 1.20–1.32-fold [1, 5, 7, 47]. The association with obesity observed in Asians in the present study was remarkably similar, with each additional minor allele increasing obesity risk by 1.25-fold (95% CI 1.19, 1.31), consistent with the association observed for obesity in previous literature-based meta-analyses of case–control studies in East and South Asians [18, 29, 30].

The association of the *FTO* variant with overweight was the same in East and South Asians (OR 1.13 per minor

allele) and very similar to the effects (ORs ranging from 1.13 to 1.18) that have been reported in large-scale studies of white Europeans [1, 7, 47]. While the effect sizes observed for the influence of *FTO* on obesity and overweight in Asians are very similar to those of Europeans, it should be noted that the definitions of obesity and overweight are different, as BMI cut-offs are somewhat lower in Asians than in Europeans, consistent with the association of BMI with metabolic disease [48].

The *FTO* minor allele increases BMI by 0.26 kg/m<sup>2</sup> (equivalent to ~750 g/allele for a person 1.7 m tall) in Asians, with very similar results for East and South Asians. This observation suggests that the effect of *FTO* on BMI in Asians is substantially smaller than the effect observed in a meta-analysis of more than 125,000 white Europeans (0.39 kg/m<sup>2</sup> per minor allele, or 1,130 g per minor allele) [7]. This difference may be due to the fact that BMI in




**Fig. 4** Association of *FTO*-rs9939609 (or proxy) with type 2 diabetes adjusted for BMI. Study-specific association analyses assumed an additive genetic model adjusted for age, sex, and BMI. Effect sizes were combined using random-effects meta-analyses (DerSimonian–Laird method)

Asians does not represent exactly the same adiposity phenotype as in Europeans. However, given that other large-scale studies in white Europeans have reported effects for *FTO* on BMI that range between 0.26 and 0.39 kg/m<sup>2</sup>, the comparison between Asians and Europeans should be made with caution [1, 3, 5, 15, 47]. The *FTO* variant also showed convincing association with measures of fat distribution such as waist and hip circumference and WHR in Asians. Despite the often described difference in abdominal obesity between East and South Asians, the effect sizes were very similar in the two groups. Consistent with the observations for BMI, the effect sizes tended to be somewhat smaller than those reported for white Europeans. For example, each additional *FTO* minor allele increased waist circumference by 0.51 cm in Asians, whereas large-scale studies in Europeans have reported an increase of 0.73–1.00 cm [1, 9, 47].

As the MAF of the *FTO* variant is substantially lower in Asians (East Asians, ~17%; South Asians, ~32%) than in white Europeans (~45%), and as the effect of this allele on obesity-related traits is similar or somewhat lower in Asians than in white Europeans, the overall contribution of genetic variation in *FTO* to obesity susceptibility will be lower in Asians, in particular East Asians. For example, the *FTO* variant explained less of the inter-individual variation in BMI in Asians (East Asians, 0.16%; South Asians, 0.20%) than in white Europeans (0.34%) [7]. Furthermore, the low risk allele frequency led to a lower PAR for the risk of obesity (East Asians, 8.3%; South Asians, 10.6%) and overweight (East Asians, 4.1%; South Asians, 7.8%) in Asians than in white Europeans (obesity, 20.4%; overweight, 12.7%) [1].

The *FTO* locus was first identified in a GWAS for type 2 diabetes in white Europeans, i.e. each minor allele



**Fig. 5** Association of *FTO*-rs9939609 (or proxy) with BMI. Study-specific association analyses assumed an additive genetic model adjusted for age and sex. Effect sizes were combined using random-effect meta-analyses (DerSimonian–Laird method)

increased the odds of diabetes by 1.15-fold [1]. However, after adjustment for BMI, the association between the *FTO* variant and type 2 diabetes was completely abolished (OR 1.03), suggesting that *FTO* is primarily an obesity-susceptibility locus [1]. In our meta-analysis, we observed a similar effect of *FTO* on risk of type 2 diabetes, with each minor allele increasing the odds by 1.15-fold. Interestingly, adjustment for BMI did not abolish the association, but only slightly attenuated it to a 1.10-fold increased risk of type 2 diabetes for each additional minor allele. These observations were similar in East and South Asians, suggesting that the *FTO* locus influences the risk of type 2 diabetes, at least in part, independently of its effect on BMI. The reason for the discrepancy between the original

observations in Europeans and our observations in Asians are not known, but may be due to the fact that *FTO* seems to have a smaller effect on BMI in Asians than in Europeans. It may also be due to the fact that BMI, as suggested above, represents a different adiposity phenotype in Asians than in Europeans because of differences in body composition. Although BMI is a marker for general adiposity, it does not distinguish between fat mass and fat-free mass and does not reflect regional fat distribution. Observational studies have suggested that, for a given amount of total body fat, East and South Asians have more abdominal fat and less muscle mass than white Europeans [49, 50]. However, while it has been generally believed that in white Europeans the association with type 2

diabetes is fully mediated by the effect of *FTO* on BMI, not all studies confirm this observation. A recent large-scale study in 41,504 Scandinavians found that the *FTO* minor allele indeed increased type 2 diabetes risk (OR 1.13), but this association remained present (OR 1.09) after adjustment for BMI, consistent with the observations in the present study. The biological pathways that underlie the independent association between *FTO* variation with obesity and type 2 diabetes remain unclear. However, results of gene expression studies have shown that *FTO* expression in human islets cells is not associated with BMI [51], whereas *FTO* mRNA and protein levels in muscle are increased in individuals with type 2 diabetes compared with non-diabetic obese individuals or healthy lean controls [52]. Furthermore, *FTO* overproduction in myotubes suggested a role for *FTO* in oxidative metabolism, lipogenesis and oxidative stress in muscle, a cluster of metabolic defects characteristic of type 2 diabetes [52].

Despite the fact that our meta-analyses included Asians with different genetic backgrounds, the overall heterogeneity of the association effects was generally only low to moderate. Interestingly, we found that the associations were generally very similar in East and South Asians, although these populations are known to have genetically different origins [53]. Furthermore, we found no differences between men and women, consistent with the observations in white Europeans [7]. We found some evidence that age may contribute to the heterogeneity of the association between *FTO* and BMI. Life course effects have been reported in white Europeans [54], and longitudinal analyses will be needed to establish this in Asian populations. Longitudinal studies are also more appropriate than cross-sectional studies for disentangling the intricate interplay between *FTO*, obesity and type 2 diabetes throughout life [15].

It should be noted that the association between *FTO* variation and obesity risk in Asians had been established in three earlier meta-analyses [18, 29, 30]. These meta-analyses were substantially smaller than the present ones and focused solely on case–control analyses of obesity and type 2 diabetes, while no continuous traits were studied. The meta-analysis by Liu et al [18] included individuals of East and South Asian origin, which were analysed together without comparison of effect sizes between the two populations. This study also examined the association with type 2 diabetes, but did not explore the association after adjustment for BMI [18]. Furthermore, the three previous meta-analyses were all literature-based and thus more prone to publication bias, whereas our meta-analysis was designed on the basis of a de novo analysis of data according to a standardised plan in all studies identified as having available data and agreement to participate. No

evidence of publication bias was observed except for the associations with BMI and hip circumference. The analytical consistency across studies helped minimise between-study heterogeneity. Although our results are representative of individuals of Southeast Asian, East Asian and South Asian descent, the association of *FTO* with risk of obesity and type 2 diabetes in other Asian populations remains to be examined.

In summary, we have firmly established that genetic variation in the first intron of *FTO* is associated with increased risk of obesity and type 2 diabetes in Asians, with effect sizes similar to those in Europeans. Furthermore, we confirm that the association of *FTO* with risk of type 2 diabetes is partly independent of BMI.

**Acknowledgements** A full list of acknowledgements can be found in the **ESM**. This work was supported by the following: Hong Kong Government Research Grants Council Central Allocation Scheme (CUHK 1/04C), Research Grants Council Earmarked Research Grant (CUHK4724/07M) and the Innovation and Technology Fund of the Government of the Hong Kong SAR (ITS/487/09FP); National Institutes of Health (DK078150, TW05596, HL085144, TW008288) and pilot funds (RR20649, ES10126, and DK56350); National Institutes of Health (R01 AR050496, R21 AG027110, R01 AG026564, R01 AR057049-01A1, R21 AA015973) and Specialized Center of Research (P50 AR055081) and National Science Foundation of China (81000363, 31000554) and the PhD Programs Foundation of Ministry of Education of China (20100201120058) and the Fundamental Research Funds for the Central Universities, Shanghai Leading Academic Discipline Project (S30501); National Research Foundation of Korea (NRF, no. 20100020617); the Japanese Society for the Promotion of Science (17209021) and of Priority Area ‘Applied Genomics’ (1601223, 17019028, 18018020, 20018026); the Korea Health 21 R&D Project, Korean Ministry of Health & Welfare (00-PJ3-PG6-GN07-001); the National Institute for Health Research Comprehensive Biomedical Research Centre at Imperial College Healthcare NHS Trust, the British Heart Foundation (SP/04/002), the Medical Research Council (G0700931), the Wellcome Trust (084723/Z/08/Z) and the National Institute for Health Research (RP-PG-0407-10371); the Knowledge Innovation Program of Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences (2009KIP401), the Chief Scientist Program of Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences (SIBS2008006), the Knowledge Innovation Program Project of the Chinese Academy of Sciences (KSCX1-YW-02), the National Natural Science Foundation of China (30930081), and National High Technology Research and Development Program (863 Program) (2009AA022704), National 973 Program (2011CB504002); the National Science Council (NSC 95-3112-B002-002, NSC 96-2752-B002-008-PAE) of Taiwan; major program of Shanghai Municipality for Basic Research (08dj1400601), National Natural Science Foundation of China (30800617); National Institute of Health (NIH/FIC KO1 TW006087, NIH/NIDDK R01 DK082766); the Program for the National Natural Science Foundation of China (NSFC-30872116) and New Century Excellent Talents in the University of China (NCET-04-0707); Council of Scientific and Industrial Research, Government of India (NWP0032); the National Heart, Lung, and Blood Institute, National Institutes of Health, US Department of Health and Human Services (N01WH22110, 24152, 32100-2, 32105-6, 32108-9, 32111-13, 32115, 32118-32119, 32122, 42107-26, 42129-32, 44221) and the National Institute of Diabetes and Digestive and Kidney Diseases R01 (DK062290) from the National Institutes of Health.



**Contribution statement** RJFL, HL and XL contributed to the conception and design of the study. CL and TOK performed the literature search, designed the analysis plan, performed the meta-analyses and researched the data. HL, TOK, CL and RJFL wrote the manuscript. HL, JZ, YL, CH, ZY, WZ, WB, SC, YW, TY, AS, BYC, CSY, DZ, FT, KY, JCC, KRM, LFB, MI, EN, NL, TF, SK, WW, CVJ, WL, YC, YX, YG, SL, YS, SHK, HDS, KSP, CHDF, JYK, PCS, KSL, WZ, XS, HD, HI, GVK, DKS, LC, LL, RH, YK, MD, KH, WJ, JSK, JCC, GRC, RCM, SM, RD, MY, RT, NK, XL and RJFL collected study-specific data, analysed the study-specific data according to the standardised analysis plan, and reviewed and edited the manuscript. All authors have approved the final version of the manuscript to be published.

**Duality of interest** The authors declare that there is no duality of interest associated with this manuscript.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

- Frayling TM, Timpson NJ, Weedon MN et al (2007) A common variant in the *FTO* gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* 316:889–894
- Scuteri A, Sanna S, Chen WM et al (2007) Genome-wide association scan shows genetic variants in the *FTO* gene are associated with obesity-related traits. *PLoS Genet* 3:e115
- Loos RJ, Lindgren CM, Li S et al (2008) Common variants near *MC4R* are associated with fat mass, weight and risk of obesity. *Nat Genet* 40:768–775
- Willer CJ, Speliotes EK, Loos RJ et al (2009) Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. *Nat Genet* 41:25–34
- Thorleifsson G, Walters GB, Gudbjartsson DF et al (2009) Genome-wide association yields new sequence variants at seven loci that associate with measures of obesity. *Nat Genet* 41:18–24
- Chambers JC, Elliott P, Zabaneh D et al (2008) Common genetic variation near *MC4R* is associated with waist circumference and insulin resistance. *Nat Genet* 40:716–718
- Speliotes EK, Willer CJ, Berndt SI et al (2010) Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet* 42:937–948
- Lindgren CM, Heid IM, Randall JC et al (2009) Genome-wide association scan meta-analysis identifies three loci influencing adiposity and fat distribution. *PLoS Genet* 5:e1000508
- Heard-Costa NL, Zillikens MC, Monda KL et al (2009) *NRXN3* is a novel locus for waist circumference: a genome-wide association study from the CHARGE Consortium. *PLoS Genet* 5:e1000539
- Heid IM, Jackson AU, Randall JC et al (2010) Meta-analysis identifies 13 new loci associated with waist–hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. *Nat Genet* 42:949–960
- Meyre D, Delplanque J, Chevre JC et al (2009) Genome-wide association study for early-onset and morbid adult obesity identifies three new risk loci in European populations. *Nat Genet* 41:157–159
- Scherag A, Dina C, Hinney A et al (2010) Two new Loci for body-weight regulation identified in a joint analysis of genome-wide association studies for early-onset extreme obesity in French and German study groups. *PLoS Genet* 6:e1000916
- Kring SL, Holst C, Zimmermann E et al (2008) *FTO* gene associated fatness in relation to body fat distribution and metabolic traits throughout a broad range of fatness. *PLoS One* 3:e2958
- Hotta K, Nakamura M, Nakamura T et al (2010) Polymorphisms in *NRXN3*, *TFAP2B*, *MSRA*, *LYPLAL1*, *FTO* and *MC4R* and their effect on visceral fat area in the Japanese population. *J Hum Genet* 55:738–742
- Hertel JK, Johansson S, Sonestedt E et al (2011) *FTO*, Type 2 diabetes, and weight gain throughout adult life: a meta-analysis of 41,504 subjects from the Scandinavian HUNT, MDC, and MPP studies. *Diabetes* 60:1637–1644
- Li H, Wu Y, Loos RJ et al (2008) Variants in the fat mass- and obesity-associated (*FTO*) gene are not associated with obesity in a Chinese Han population. *Diabetes* 57:264–268
- Horikoshi M, Hara K, Ito C et al (2007) Variations in the *HHEX* gene are associated with increased risk of type 2 diabetes in the Japanese population. *Diabetologia* 50:2461–2466
- Liu Y, Liu Z, Song Y et al (2010) Meta-analysis added power to identify variants in *FTO* associated with type 2 diabetes and obesity in the Asian population. *Obesity (Silver Spring)* 18:1619–1624
- Karasawa S, Daimon M, Sasaki S et al (2010) Association of the common fat mass and obesity associated (*FTO*) gene polymorphism with obesity in a Japanese population. *Endocr J* 57:293–301
- Hotta K, Nakata Y, Matsuo T et al (2008) Variations in the *FTO* gene are associated with severe obesity in the Japanese. *J Hum Genet* 53:546–553
- Ng MC, Tam CH, So WY et al (2010) Implication of genetic variants near *NEGR1*, *SEC16B*, *TMEM18*, *ETV5/DGKG*, *GNPDA2*, *LIN7C/BDNF*, *MTCH2*, *BCDIN3D/FAIM2*, *SH2B1*, *FTO*, *MC4R*, and *KCTD15* with obesity and type 2 diabetes in 7705 Chinese. *J Clin Endocrinol Metab* 95:2418–2425
- Tan JT, Dorajoo R, Seielstad M et al (2008) *FTO* variants are associated with obesity in the Chinese and Malay populations in Singapore. *Diabetes* 57:2851–2857
- Chang YC, Liu PH, Lee WJ et al (2008) Common variation in the fat mass and obesity-associated (*FTO*) gene confers risk of obesity and modulates BMI in the Chinese population. *Diabetes* 57:2245–2252
- Cheung CY, Tso AW, Cheung BM et al (2010) Obesity susceptibility genetic variants identified from recent genome-wide association studies: implications in a Chinese population. *J Clin Endocrinol Metab* 95:1395–1403
- Cha SW, Choi SM, Kim KS et al (2008) Replication of genetic effects of *FTO* polymorphisms on BMI in a Korean population. *Obesity (Silver Spring)* 16:2187–2189
- Lee HJ, Kim IK, Kang JH et al (2010) Effects of common *FTO* gene variants associated with BMI on dietary intake and physical activity in Koreans. *Clin Chim Acta* 411:1716–1722
- Marvelle AF, Lange LA, Qin L, Adair LS, Mohlke KL (2008) Association of *FTO* with obesity-related traits in the Cebu Longitudinal Health and Nutrition Survey (CLHNS) Cohort. *Diabetes* 57:1987–1991
- Cho YS, Go MJ, Kim YJ et al (2009) A large-scale genome-wide association study of Asian populations uncovers genetic factors influencing eight quantitative traits. *Nat Genet* 41:527–534
- Peng S, Zhu Y, Xu F, Ren X, Li X, Lai M (2011) *FTO* gene polymorphisms and obesity risk: a meta-analysis. *BMC Med* 9:71
- Xi B, Mi J (2009) *FTO* polymorphisms are associated with obesity but not with diabetes in East Asian populations: a meta-analysis. *Biomed Environ Sci* 22:449–457



31. Dorajoo R, Blakemore AI, Sim X et al (2011) Replication of 13 obesity loci among Singaporean Chinese, Malay and Asian-Indian populations. *Int J Obes (Lond)*. doi:10.1038/ijo.2011.86
32. Ramya K, Radha V, Ghosh S, Majumder PP, Mohan V (2010) Genetic variations in the *FTO* gene are associated with type 2 diabetes and obesity in south Indians (CURES-79). *Diabetes Technol Ther* 13:33–42
33. Yajnik CS, Janipalli CS, Bhaskar S et al (2009) *FTO* gene variants are strongly associated with type 2 diabetes in South Asian Indians. *Diabetologia* 52:247–252
34. Ng MC, Park KS, Oh B et al (2008) Implication of genetic variants near *TCF7L2*, *SLC30A8*, *HHEX*, *CDKAL1*, *CDKN2A/B*, *IGF2BP2*, and *FTO* in type 2 diabetes and obesity in 6,719 Asians. *Diabetes* 57:2226–2233
35. Li X, Song F, Jiang H et al (2010) A genetic variation in the fat mass- and obesity-associated gene is associated with obesity and newly diagnosed type 2 diabetes in a Chinese population. *Diabetes Metab Res Rev* 26:128–132
36. Shi J, Long J, Gao YT et al (2010) Evaluation of genetic susceptibility loci for obesity in Chinese women. *Am J Epidemiol* 172:244–254
37. Omori S, Tanaka Y, Takahashi A et al (2008) Association of *CDKAL1*, *IGF2BP2*, *CDKN2A/B*, *HHEX*, *SLC30A8*, and *KCNJ11* with susceptibility to type 2 diabetes in a Japanese population. *Diabetes* 57:791–795
38. Tabara Y, Osawa H, Guo H et al (2009) Prognostic significance of *FTO* genotype in the development of obesity in Japanese: the J-SHIPP study. *Int J Obes (Lond)* 33:1243–1248
39. Han X, Luo Y, Ren Q et al (2010) Implication of genetic variants near *SLC30A8*, *HHEX*, *CDKAL1*, *CDKN2A/B*, *IGF2BP2*, *FTO*, *TCF2*, *KCNQ1*, and *WFS1* in type 2 diabetes in a Chinese population. *BMC Med Genet* 11:81
40. Sanghera DK, Ortega L, Han S et al (2008) Impact of nine common type 2 diabetes risk polymorphisms in Asian Indian Sikhs: *PPARG2* (Pro12Ala), *IGF2BP2*, *TCF7L2* and *FTO* variants confer a significant risk. *BMC Med Genet* 9:59
41. Takeuchi F, Yamamoto K, Katsuya T et al (2011) Association of genetic variants for susceptibility to obesity with type 2 diabetes in Japanese individuals. *Diabetologia* 54:1350–1359
42. Shu XO, Long J, Cai Q et al (2010) Identification of new genetic risk variants for type 2 diabetes. *PLoS Genet* 6: e1001127
43. Zhou BF, Cooperative Meta-Analysis Group of the Working Group on Obesity in China (2002) Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults: study on optimal cut-off points of body mass index and waist circumference in Chinese adults. *Biomed Environ Sci* 15:83–96
44. DerSimonian R, Laird N (1986) Meta-analysis in clinical trials. *Control Clin Trials* 7:177–188
45. Higgins JP, Thompson SG (2002) Quantifying heterogeneity in a meta-analysis. *Stat Med* 21:1539–1558
46. Ng MC, Tam CH, Lam VK, So WY, Ma RC, Chan JC (2007) Replication and identification of novel variants at *TCF7L2* associated with type 2 diabetes in Hong Kong Chinese. *J Clin Endocrinol Metab* 92:3733–3737
47. Li S, Zhao JH, Luan J et al (2009) Cumulative effects and predictive value of common obesity-susceptibility variants identified by genome-wide association studies. *Am J Clin Nutr* 91:184–190
48. Consultation WHOE (2004) Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 363:157–163
49. Lear SA, Humphries KH, Kohli S, Chockalingam A, Frohlich JJ, Birmingham CL (2007) Visceral adipose tissue accumulation differs according to ethnic background: results of the Multicultural Community Health Assessment Trial (M-CHAT). *Am J Clin Nutr* 86:353–359
50. Chan JC, Malik V, Jia W et al (2009) Diabetes in Asia: epidemiology, risk factors, and pathophysiology. *JAMA* 301:2129–2140
51. Kirkpatrick CL, Marchetti P, Purrello F et al (2010) Type 2 diabetes susceptibility gene expression in normal or diabetic sorted human alpha and beta cells: correlations with age or BMI of islet donors. *PLoS One* 5:e11053
52. Bravard A, Lefai E, Meugnier E et al (2011) *FTO* is increased in muscle during type 2 diabetes, and its overexpression in myotubes alters insulin signaling, enhances lipogenesis and ROS production, and induces mitochondrial dysfunction. *Diabetes* 60:258–268
53. Consortium HP-AS, Abdulla MA, Ahmed I et al (2009) Mapping human genetic diversity in Asia. *Science* 326:1541–1545
54. Hardy R, Wills AK, Wong A et al (2009) Life course variations in the associations between *FTO* and *MC4R* gene variants and body size. *Hum Mol Genet* 19:545–552
55. Sha BY, Yang TL, Zhao LJ et al (2009) Genome-wide association study suggested copy number variation may be associated with body mass index in the Chinese population. *J Hum Genet* 54:199–202
56. Hu C, Zhang R, Wang C et al (2009) *PPARG*, *KCNJ11*, *CDKAL1*, *CDKN2A-CDKN2B*, *IDE-KIF11-HHEX*, *IGF2BP2* and *SLC30A8* are associated with type 2 diabetes in a Chinese population. *PLoS One* 4:e7643
57. Wen J, Ronn T, Olsson A et al (2010) Investigation of type 2 diabetes risk alleles support *CDKN2A/B*, *CDKAL1*, and *TCF7L2* as susceptibility genes in a Han Chinese cohort. *PLoS One* 5:e9153
58. Cha S, Koo I, Park BL et al (2009) Genetic effects of *FTO* and *MC4R* polymorphisms on body mass in constitutional types. *Evid Based Complement Alternat Med*. doi:10.1093/ecam/nep162
59. Takeuchi F, Serizawa M, Yamamoto K et al (2009) Confirmation of multiple risk loci and genetic impacts by a genome-wide association study of type 2 diabetes in the Japanese population. *Diabetes* 58:1690–1699