

## Research Article

# Jordan $*$ -Derivations on $C^*$ -Algebras and $JC^*$ -Algebras

Jong Su An,<sup>1</sup> Jianlian Cui,<sup>2</sup> and Choonkil Park<sup>3</sup>

<sup>1</sup> Department of Mathematics Education, Pusan National University, Pusan 609-735, South Korea

<sup>2</sup> Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China

<sup>3</sup> Department of Mathematics, Hanyang University, Seoul 133-791, South Korea

Correspondence should be addressed to Choonkil Park, baak@hanyang.ac.kr

Received 19 September 2008; Revised 20 October 2008; Accepted 31 October 2008

Recommended by Ferhan Merdivenci Atici

We investigate Jordan  $*$ -derivations on  $C^*$ -algebras and Jordan  $*$ -derivations on  $JC^*$ -algebras associated with the following functional inequality  $\|f(x) + f(y) + kf(z)\| \leq \|kf((x+y)/k+z)\|$  for some integer  $k$  greater than 1. We moreover prove the generalized Hyers-Ulam stability of Jordan  $*$ -derivations on  $C^*$ -algebras and of Jordan  $*$ -derivations on  $JC^*$ -algebras associated with the following functional equation  $f((x+y)/k+z) = (f(x) + f(y))/k + f(z)$  for some integer  $k$  greater than 1.

Copyright © 2008 Jong Su An et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction and preliminaries

The stability problem of functional equations originated from a question of Ulam [1] concerning the stability of group homomorphisms. Hyers [2] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' theorem was generalized by Aoki [3] for additive mappings and by Th. M. Rassias [4] for linear mappings by considering an unbounded Cauchy difference. In 1982–1994, a generalization of Hyers-Ulam stability result was proved by J. M. Rassias [5–9]. This author assumed that Cauchy-Găvruta-Rassias inequality

$$\|f(x+y) - f(x) - f(y)\| \leq \theta \|x\|^p \cdot \|y\|^q \quad (1.1)$$

is controlled by a product of different powers of norms, where  $\theta \geq 0$  and  $p, q \in \mathbb{R}$  such that  $r = p+q \neq 1$ , and retained the condition of continuity of  $f(tx)$  in  $t \in \mathbb{R}$  for each fixed  $x$ . In 1999, Găvruta [10] studied the singular case  $r = p+q = 1$ , by constructing a nice counterexample to the above pertinent Ulam stability problem. Also J. M. Rassias [5–9, 11–13] investigated other

conditions and still obtained stability results. In all these cases, the approach to the existence question was to prove asymptotic type formulas of the form

$$L(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n} \quad \text{or} \quad L(x) = \lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}\right). \quad (1.2)$$

**Theorem 1.1** (see [5–7, 9]). *Let  $X$  be a real normed vector space and  $Y$  a real complete normed vector space. Assume in addition that  $f : X \rightarrow Y$  is an approximately additive mapping for which there exist constants  $\theta \geq 0$  and  $p, q \in \mathbb{R}$  such that  $r = p + q \neq 1$  and  $f$  satisfies*

$$\|f(x + y) - f(x) - f(y)\| \leq \theta \|x\|^p \cdot \|y\|^q \quad (1.3)$$

for all  $x, y \in X$ . Then, there exists a unique additive mapping  $L : X \rightarrow Y$  satisfying

$$\|f(x) - L(x)\| \leq \frac{\theta}{|2 - 2^r|} \|x\|^r \quad (1.4)$$

for all  $x \in X$ . If, in addition,  $f : X \rightarrow Y$  is a mapping such that the transformation  $t \mapsto f(tx)$  is continuous in  $t \in \mathbb{R}$  for each fixed  $x \in X$ , then  $L$  is an  $\mathbb{R}$ -linear mapping.

**Theorem 1.2** (see [4]). *Let  $f : E \rightarrow E'$  be a mapping from a normed vector space  $E$  into a Banach space  $E'$  subject to the inequality*

$$\|f(x + y) - f(x) - f(y)\| \leq \varepsilon (\|x\|^p + \|y\|^p) \quad (1.5)$$

for all  $x, y \in E$ , where  $\varepsilon$  and  $p$  are constants with  $\varepsilon > 0$  and  $p < 1$ . Then, the limit

$$L(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n} \quad (1.6)$$

exists for all  $x \in E$ , and  $L : E \rightarrow E'$  is the unique additive mapping which satisfies

$$\|f(x) - L(x)\| \leq \frac{2\varepsilon}{2 - 2^p} \|x\|^p \quad (1.7)$$

for all  $x \in E$ . Also, if for each  $x \in E$  the mapping  $f(tx)$  is continuous in  $t \in \mathbb{R}$ , then  $L$  is linear.

Th. M. Rassias [14], during the 27th International Symposium on Functional Equations, asked the question whether such a theorem can also be proved for  $p \geq 1$ . Gajda [15] following the same approach as in Th. M. Rassias [4] gave an affirmative solution to this question for  $p > 1$ . It was shown by Gajda [15] as well as by Th. M. Rassias and Šemrl [16] that one cannot prove a Th. M. Rassias' type theorem when  $p = 1$ . The counterexamples of Gajda [15] as well as of Th. M. Rassias and Šemrl [16] have stimulated several mathematicians to invent new definitions of *approximately additive* or *approximately linear* mappings, compare Găvruta [17] and Jung [18], who among others studied the Hyers-Ulam stability of functional equations. Theorem 1.2 that was introduced for the first time by Th. M. Rassias [4] provided a lot of influence in the development of a generalization of the Hyers-Ulam stability concept. This new concept is known as *generalized Hyers-Ulam stability* of functional equations (cf. the books of Czerwik [19], Hyers et al. [20]).

Găvruta [17] provided a further generalization of Th. M. Rassias' theorem. Isac and Th. M. Rassias [21] applied the Hyers-Ulam stability theory to prove fixed point theorems

and study some new applications in nonlinear analysis. In [22], Hyers et al. studied the asymptoticity aspect of Hyers-Ulam stability of mappings. Beginning around the year 1980, the topic of approximate homomorphisms and their stability theory in the field of functional equations and inequalities was taken up by several mathematicians (see [3–18, 21–51]).

Gilányi [26] showed that if  $f$  satisfies the functional inequality

$$\|2f(x) + 2f(y) - f(x - y)\| \leq \|f(x + y)\|, \quad (1.8)$$

then  $f$  satisfies the Jordan-von Neumann functional equation

$$2f(x) + 2f(y) = f(x + y) + f(x - y). \quad (1.9)$$

See also [52]. Fechner [53] and Gilányi [27] proved the generalized Hyers-Ulam stability of the functional inequality (1.8). Park et al. [42] introduced and investigated 3-variable Cauchy-Jensen functional inequalities and proved the generalized Hyers-Ulam stability of the 3-variable Cauchy-Jensen functional inequalities.

*Definition 1.3.* Let  $A$  be a  $C^*$ -algebra. A  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow A$  is called a Jordan  $*$ -derivation if

$$\delta(a^2) = a\delta(a) + \delta(a^*)a^* \quad (1.10)$$

for all  $a \in A$ .

A  $C^*$ -algebra  $A$ , endowed with the Jordan product  $a \circ b := (ab + ba)/2$  on  $A$ , is called a  $JC^*$ -algebra (see [38, 39]).

*Definition 1.4.* Let  $A$  be a  $JC^*$ -algebra. A  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow A$  is called a Jordan  $*$ -derivation if

$$\delta(a^2) = a \circ \delta(a) + \delta(a^*) \circ a^* \quad (1.11)$$

for all  $a \in A$ .

This paper is organized as follows. In Section 2, we investigate Jordan  $*$ -derivations on  $C^*$ -algebras associated with the functional inequality

$$\|f(x) + f(y) + kf(z)\| \leq \left\| kf\left(\frac{x+y}{k} + z\right) \right\|, \quad (1.12)$$

and prove the generalized Hyers-Ulam stability of Jordan  $*$ -derivations on  $C^*$ -algebras associated with the functional equation

$$f\left(\frac{x+y}{k} + z\right) = \frac{f(x) + f(y)}{k} + f(z). \quad (1.13)$$

In Section 3, we investigate Jordan  $*$ -derivations on  $JC^*$ -algebras associated with the functional inequality (1.12), and prove the generalized Hyers-Ulam stability of Jordan  $*$ -derivations on  $JC^*$ -algebras associated with the functional equation (1.13).

Throughout this paper, let  $k$  be an integer greater than 1.

## 2. Jordan \*-derivations on $C^*$ -algebras

Throughout this section, assume that  $A$  is a  $C^*$ -algebra with norm  $\|\cdot\|$ .

**Lemma 2.1.** *Let  $f : A \rightarrow A$  be a mapping such that*

$$\|f(a) + f(b) + kf(c)\| \leq \left\| kf\left(\frac{a+b}{k} + c\right) \right\| \quad (2.1)$$

for all  $a, b, c \in A$ . Then,  $f$  is Cauchy additive, that is,  $f(a+b) = f(a) + f(b)$ .

*Proof.* Letting  $a = b = c = 0$  in (2.1), we get

$$\|(k+2)f(0)\| \leq \|kf(0)\|. \quad (2.2)$$

So  $f(0) = 0$ .

Letting  $c = 0$  and  $b = -a$  in (2.1), we get

$$\|f(a) + f(-a)\| \leq \|kf(0)\| = 0 \quad (2.3)$$

for all  $a \in A$ . Hence  $f(-a) = -f(a)$  for all  $a \in A$ .

Letting  $c = -(a+b)/k$  in (2.1), we get

$$\left\| f(a) + f(b) - kf\left(\frac{a+b}{k}\right) \right\| = \left\| f(a) + f(b) + kf\left(-\frac{a+b}{k}\right) \right\| \leq \|kf(0)\| = 0 \quad (2.4)$$

for all  $a, b \in A$ . Thus

$$kf\left(\frac{a+b}{k}\right) = f(a) + f(b) \quad (2.5)$$

for all  $a, b \in A$ . Letting  $b = 0$  in (2.5), we get  $kf(a/k) = f(a)$  for all  $a \in A$ . So

$$f(a+b) = kf\left(\frac{a+b}{k}\right) = f(a) + f(b) \quad (2.6)$$

for all  $a, b \in A$ , as desired.  $\square$

**Theorem 2.2.** *Let  $r > 1$  and  $\theta$  be a nonnegative real number, and let  $f : A \rightarrow A$  be a mapping such that*

$$\|\mu f(a) + f(b) + kf(c)\| \leq \left\| kf\left(\frac{\mu a + b}{k} + c\right) \right\|, \quad (2.7)$$

$$\|f(a^2) - af(a) - f(a^*)a^*\| \leq \theta \|a\|^{2r} \quad (2.8)$$

for all  $\mu \in \mathbb{T} := \{\lambda \in \mathbb{C} \mid |\lambda| = 1\}$  and all  $a, b, c \in A$ . Then, the mapping  $f : A \rightarrow A$  is a Jordan \*-derivation.

*Proof.* Let  $\mu = 1$  in (2.7). By Lemma 2.1, the mapping  $f : A \rightarrow A$  is Cauchy additive. So  $f(0) = 0$  and  $f(x) = \lim_{n \rightarrow \infty} 2^n f(a/2^n)$  for all  $a \in A$ .

Letting  $b = -\mu a$  and  $c = 0$ , we get

$$\|\mu f(a) + f(-\mu a)\| \leq \|kf(0)\| = 0 \quad (2.9)$$

for all  $a \in A$  and all  $\mu \in \mathbb{T}$ . So

$$\mu f(a) - f(\mu a) = \mu f(a) + f(-\mu a) = 0 \quad (2.10)$$

for all  $a \in A$ , and all  $\mu \in \mathbb{T}$ . Hence,  $f(\mu a) = \mu f(a)$  for all  $a \in A$  and all  $\mu \in \mathbb{T}$ . By the same reasoning as in the proof of [39, Theorem 2.1], the mapping  $f : A \rightarrow A$  is  $\mathbb{C}$ -linear.

It follows from (2.8) that

$$\begin{aligned} \|f(a^2) - af(a) - f(a^*)a^*\| &= \lim_{n \rightarrow \infty} 4^n \left\| f\left(\frac{a^2}{2^n \cdot 2^n}\right) - \frac{a}{2^n} f\left(\frac{a}{2^n}\right) - f\left(\frac{a^*}{2^n}\right) \frac{a^*}{2^n} \right\| \\ &\leq \lim_{n \rightarrow \infty} \frac{4^n \theta}{4^{nr}} \|a\|^{2r} = 0 \end{aligned} \quad (2.11)$$

for all  $a \in A$ . Thus

$$f(a^2) = af(a) + f(a^*)a^* \quad (2.12)$$

for all  $a \in A$ .

Hence the mapping  $f : A \rightarrow A$  is a Jordan  $*$ -derivation.  $\square$

**Theorem 2.3.** *Let  $r < 1$  and  $\theta$  be a nonnegative real number, and let  $f : A \rightarrow A$  be a mapping satisfying (2.7) and (2.8). Then, the mapping  $f : A \rightarrow A$  is a Jordan  $*$ -derivation.*

*Proof.* The proof is similar to the proof of Theorem 2.2.  $\square$

We prove the generalized Hyers-Ulam stability of Jordan  $*$ -derivations on  $C^*$ -algebras.

**Theorem 2.4.** *Suppose that  $f : A \rightarrow A$  is a mapping with  $f(0) = 0$  for which there exists a function  $\varphi : A \times A \times A \rightarrow [0, \infty)$  such that*

$$\lim_{n \rightarrow \infty} k^{-n} \varphi(k^n a, k^n b, k^n c) = 0, \quad (2.13)$$

$$\tilde{\varphi}(a) := \sum_{n=1}^{\infty} k^{-n+1} \varphi(k^n a, 0, 0) < \infty, \quad (2.14)$$

$$\left\| f\left(\frac{\lambda a + \lambda b}{k} + c^2\right) - \frac{\lambda}{k} f(a) - \frac{\lambda}{k} f(b) - cf(c) - f(c^*)c^* \right\| \leq \varphi(a, b, c) \quad (2.15)$$

for all  $\lambda \in \mathbb{T}$ , and all  $a, b, c \in A$ . Then, there exists a unique Jordan  $*$ -derivation  $\delta : A \rightarrow A$  such that

$$\|f(a) - \delta(a)\| \leq \tilde{\varphi}(a) \quad (2.16)$$

for all  $a \in A$ .

*Proof.* Putting  $b = c = 0$  and  $\lambda = 1$ , and replacing  $a$  by  $ka$  in (2.15), we get

$$\left\| f(a) - \frac{f(ka)}{k} \right\| \leq \varphi(ka, 0, 0) \quad (\forall a \in A). \quad (2.17)$$

Using the induction method, we have

$$\left\| \frac{f(k^n a)}{k^n} - \frac{f(k^m a)}{k^m} \right\| \leq \sum_{j=m+1}^n k^{-j+1} \varphi(k^j a, 0, 0) \quad (2.18)$$

for all  $n > m \geq 0$  and all  $a \in A$ . It follows that for every  $a \in A$ , the sequence  $\{f(k^n a)/k^n\}$  is Cauchy, and hence it is convergent since  $A$  is complete. Set

$$\delta(a) := \lim_{n \rightarrow \infty} \frac{f(k^n a)}{k^n} \quad (a \in A). \quad (2.19)$$

Let  $c = 0$  and replace  $a$  and  $b$  by  $k^n a$  and  $k^n b$ , respectively, in (2.15), we get

$$\left\| k^{-n} f\left(k^n \frac{\lambda a + \lambda b}{k}\right) - \frac{\lambda}{k} k^{-n} f(k^n a) - \frac{\lambda}{k} k^{-n} f(k^n b) \right\| \leq k^{-n} \varphi(k^n a, k^n b, 0). \quad (2.20)$$

Taking the limit as  $n \rightarrow \infty$ , we obtain

$$\delta\left(\frac{\lambda a + \lambda b}{k}\right) = \frac{\lambda}{k} \delta(a) + \frac{\lambda}{k} \delta(b) \quad (2.21)$$

for all  $a, b \in A$  and all  $\lambda \in \mathbb{T}$ . Letting  $b = 0$  and  $\lambda = 1$  in (2.21), we get

$$\delta\left(\frac{a}{k}\right) = \frac{1}{k} \delta(a) \quad (2.22)$$

for all  $a \in A$ . Hence,

$$\lambda \delta(a) + \lambda \delta(b) = k \delta\left(\frac{\lambda a + \lambda b}{k}\right) = \delta(\lambda a + \lambda b) \quad (2.23)$$

for all  $a, b \in A$ , in particular,  $\delta : A \rightarrow A$  is additive. Now similar to the discussion in [54, Theorem 2.1], we show that  $\delta : A \rightarrow A$  is  $\mathbb{C}$ -linear. Letting  $m = 0$  in (2.18), we get

$$\left\| f(a) - \frac{f(k^n a)}{k^n} \right\| \leq \sum_{j=1}^n k^{-j+1} \varphi(k^j a, 0, 0). \quad (2.24)$$

Taking the limit as  $n \rightarrow \infty$ , we have

$$\|f(a) - \delta(a)\| \leq \tilde{\varphi}(a) \quad (2.25)$$

for all  $a \in A$ . It follows from [55] that  $\delta : A \rightarrow A$  is unique.

Letting  $\lambda = 1$ ,  $a = b = 0$  and replacing  $c$  by  $k^n c$  in (2.15), we obtain

$$\|f(k^{2n}(c^2)) - k^n c f(k^n c) - k^n f(k^n c^*) c^*\| \leq \varphi(0, 0, k^n c). \quad (2.26)$$

So

$$\|k^{-2n} f(k^{2n}(c^2)) - k^{-n} c f(k^n c) - k^{-n} f(k^n c^*) c^*\| \leq k^{-2n} \varphi(0, 0, k^n c) \quad (2.27)$$

for all  $c \in A$ . Letting  $n$  tend to infinity, we have

$$\delta(c^2) = c \delta(c) + \delta(c^*) c^* \quad (2.28)$$

for all  $c \in A$ . Hence  $\delta : A \rightarrow A$  is a Jordan  $*$ -derivation.  $\square$

**Corollary 2.5.** Suppose that  $f : A \rightarrow A$  is a mapping with  $f(0) = 0$  for which there exist constant  $\beta \geq 0$  and  $p_1, p_2, p_3 \in (-\infty, 1)$  such that

$$\left\| f\left(\frac{\lambda a + \lambda b}{k} + c^2\right) - \frac{\lambda}{k}f(a) - \frac{\lambda}{k}f(b) - cf(c) - f(c^*)c^* \right\| \leq \beta(\|a\|^{p_1} + \|b\|^{p_2} + \|c\|^{p_3}) \quad (2.29)$$

for all  $\lambda \in \mathbb{T}$  and all  $a, b, c \in A$ . Then, there exists a unique Jordan  $*$ -derivation  $\delta : A \rightarrow A$  such that

$$\|f(a) - \delta(a)\| \leq \frac{k\beta\|a\|^{p_1}}{k^{1-p_1} - 1} \quad (2.30)$$

for all  $a \in A$ .

*Proof.* Letting  $\varphi(a, b, c) := \beta(\|a\|^{p_1} + \|b\|^{p_2} + \|c\|^{p_3})$  in Theorem 2.4, we obtain the result.  $\square$

**Theorem 2.6.** Suppose that  $f : A \rightarrow A$  is a mapping with  $f(0) = 0$  for which there exists a function  $\varphi : A \times A \times A \rightarrow [0, \infty)$  satisfying (2.15) such that

$$\begin{aligned} \lim_{n \rightarrow \infty} k^n \varphi(k^{-n}a, k^{-n}b, k^{-n}c) &= 0, \\ \tilde{\varphi}(a) &:= \sum_{n=1}^{\infty} k^{n+1} \varphi(k^{-n}a, 0, 0) < \infty \end{aligned} \quad (2.31)$$

for all  $a, b, c \in A$ . Then, there exists a unique Jordan  $*$ -derivation  $\delta : A \rightarrow A$  such that

$$\|f(a) - \delta(a)\| \leq \tilde{\varphi}(a) \quad (2.32)$$

for all  $a \in A$ .

*Proof.* Letting  $b = c = 0$  and  $\lambda = 1$  in (2.15), we get

$$\left\| kf\left(\frac{a}{k}\right) - f(a) \right\| \leq k\varphi(a, 0, 0) \quad (\forall a \in A). \quad (2.33)$$

One can apply the induction method to prove that

$$\|k^n f(k^{-n}a) - k^m f(k^{-m}b)\| \leq \sum_{j=m}^{n-1} k^{j+1} \varphi(k^{-j}a, 0, 0) \quad (2.34)$$

for all  $n > m \geq 0$  and  $a \in A$ . It follows that for every  $a \in A$ , the sequence  $\{k^n f(k^{-n}a)\}$  is Cauchy, and hence it is convergent since  $A$  is complete. Set

$$\delta(a) := \lim_{n \rightarrow \infty} k^n f(k^{-n}a) \quad (a \in A). \quad (2.35)$$

Letting  $c = 0$  and replacing  $a$  and  $b$  by  $k^{-n}a$  and  $k^{-n}b$ , respectively, in (2.15), we get

$$\left\| k^n f\left(\frac{k^{-n}(a+b)}{k}\right) - \frac{k^n}{k}f(k^{-n}a) - \frac{k^n}{k}f(k^{-n}b) \right\| \leq k^n \varphi(k^{-n}a, k^{-n}b, 0) \quad (2.36)$$

for all  $a, b \in A$ . Taking the limit as  $n \rightarrow \infty$ , we obtain

$$\delta\left(\frac{a+b}{k}\right) = \frac{\delta(a)}{k} + \frac{\delta(b)}{k} \quad (2.37)$$

for all  $a, b \in A$ . Hence  $\delta : A \rightarrow A$  is additive.

The rest of the proof is similar to the proof of Theorem 2.4.  $\square$

**Corollary 2.7.** Suppose that  $f : A \rightarrow A$  is a mapping with  $f(0) = 0$  for which there exist constant  $\beta \geq 0$  and  $p_1, p_2, p_3 \in (1, \infty)$  such that

$$\left\| f\left(\frac{\lambda a + \lambda b}{k} + c^2\right) - \frac{\lambda}{k}f(a) - \frac{\lambda}{k}f(b) - cf(c) - f(c^*)c^* \right\| \leq \beta(\|a\|^{p_1} + \|b\|^{p_2} + \|c\|^{p_3}) \quad (2.38)$$

for all  $\lambda \in \mathbb{T}$  and all  $a, b, c \in A$ . Then, there exists a unique Jordan  $*$ -derivation  $\delta : A \rightarrow A$  such that

$$\|f(a) - \delta(a)\| \leq \frac{k\beta\|a\|^{p_1}}{k^{1-p_1} - 1} \quad (2.39)$$

for all  $a \in A$ .

*Proof.* Letting  $\varphi(a, b, c) := \beta(\|a\|^{p_1} + \|b\|^{p_2} + \|c\|^{p_3})$  in Theorem 2.6, we obtain the result.  $\square$

### 3. Jordan $*$ -derivations on $JC^*$ -algebras

Throughout this section, assume that  $A$  is a  $JC^*$ -algebra with norm  $\|\cdot\|$ .

**Theorem 3.1.** Let  $r > 1$  and  $\theta$  be a nonnegative real number, and let  $f : A \rightarrow A$  be a mapping satisfying (2.7) such that

$$\|f(a^2) - a \circ f(a) - f(a^*) \circ a^*\| \leq \theta\|a\|^{2r} \quad (3.1)$$

for all  $a \in A$ . Then, the mapping  $f : A \rightarrow A$  is a Jordan  $*$ -derivation.

*Proof.* By the same reasoning as in the proof of Theorem 2.2, the mapping  $f : A \rightarrow A$  is  $\mathbb{C}$ -linear.

It follows from (3.1) that

$$\begin{aligned} \|f(a^2) - a \circ f(a) - f(a^*) \circ a^*\| &= \lim_{n \rightarrow \infty} 4^n \left\| f\left(\frac{a^2}{2^n \cdot 2^n}\right) - \frac{a}{2^n} \circ f\left(\frac{a}{2^n}\right) - f\left(\frac{a^*}{2^n}\right) \circ \frac{a^*}{2^n} \right\| \\ &\leq \lim_{n \rightarrow \infty} \frac{4^n \theta}{4^{nr}} \|a\|^{2r} = 0 \end{aligned} \quad (3.2)$$

for all  $a \in A$ . Thus,

$$f(a^2) = a \circ f(a) + f(a^*) \circ a^* \quad (3.3)$$

for all  $a \in A$ .

Hence, the mapping  $f : A \rightarrow A$  is a Jordan  $*$ -derivation.  $\square$

**Theorem 3.2.** Let  $r < 1$  and  $\theta$  be a nonnegative real number, and let  $f : A \rightarrow A$  be a mapping satisfying (2.7) and (3.1). Then, the mapping  $f : A \rightarrow A$  is a Jordan  $*$ -derivation.

*Proof.* The proof is similar to the proofs of Theorems 2.2 and 3.1.  $\square$

We prove the generalized Hyers-Ulam stability of Jordan  $*$ -derivations on  $JC^*$ -algebras.



**Theorem 3.3.** Suppose that  $f : A \rightarrow A$  is a mapping with  $f(0) = 0$  for which there exists a function  $\varphi : A \times A \times A \rightarrow [0, \infty)$  satisfying (2.13) and (2.14) such that

$$\left\| f\left(\frac{\lambda a + \lambda b}{k} + c^2\right) - \frac{\lambda}{k}f(a) - \frac{\lambda}{k}f(b) - c \circ f(c) - f(c^*) \circ c^* \right\| \leq \varphi(a, b, c) \quad (3.4)$$

for all  $\lambda \in \mathbb{T}$  and all  $a, b, c \in A$ . Then, there exists a unique Jordan  $*$ -derivation  $\delta : A \rightarrow A$  such that

$$\|f(a) - \delta(a)\| \leq \tilde{\varphi}(a) \quad (3.5)$$

for all  $a \in A$ .

*Proof.* By the same reasoning as in the proof of Theorem 2.4, there exists a unique  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow A$  such that

$$\|f(a) - \delta(a)\| \leq \tilde{\varphi}(a) \quad (3.6)$$

for all  $a \in A$ . The mapping  $\delta : A \rightarrow A$  is given by

$$\delta(a) := \lim_{n \rightarrow \infty} \frac{f(k^n a)}{k^n} \quad (a \in A). \quad (3.7)$$

Letting  $\lambda = 1$ ,  $a = b = 0$  and replacing  $c$  by  $k^n c$  in (3.4), we obtain

$$\|f(k^{2n} c^2) - k^n c \circ f(k^n c) - k^n f(k^n c^*) \circ c^*\| \leq \varphi(0, 0, k^n c). \quad (3.8)$$

So

$$\|k^{-2n} f(k^{2n} c^2) - k^{-n} c \circ f(k^n c) - k^{-n} f(k^n c^*) \circ c^*\| \leq k^{-2n} \varphi(0, 0, k^n c) \quad (3.9)$$

for all  $c \in A$ . Letting  $n$  tend to infinity, we have

$$\delta(c^2) = c \circ \delta(c) + \delta(c^*) \circ c^* \quad (3.10)$$

for all  $c \in A$ . Hence,  $\delta : A \rightarrow A$  is a Jordan  $*$ -derivation.  $\square$

**Corollary 3.4.** Suppose that  $f : A \rightarrow A$  is a mapping with  $f(0) = 0$  for which there exist constant  $\beta \geq 0$  and  $p_1, p_2, p_3 \in (-\infty, 1)$  such that

$$\left\| f\left(\frac{\lambda a + \lambda b}{k} + c^2\right) - \frac{\lambda}{k}f(a) - \frac{\lambda}{k}f(b) - c \circ f(c) - f(c^*) \circ c^* \right\| \leq \beta(\|a\|^{p_1} + \|b\|^{p_2} + \|c\|^{p_3}) \quad (3.11)$$

for all  $\lambda \in \mathbb{T}$  and all  $a, b, c \in A$ . Then, there exists a unique Jordan  $*$ -derivation  $\delta : A \rightarrow A$  such that

$$\|f(a) - \delta(a)\| \leq \frac{k\beta\|a\|^{p_1}}{k^{1-p_1} - 1} \quad (3.12)$$

for all  $a \in A$ .

*Proof.* Letting  $\varphi(a, b, c) := \beta(\|a\|^{p_1} + \|b\|^{p_2} + \|c\|^{p_3})$  in Theorem 3.3, we obtain the result.  $\square$

**Theorem 3.5.** Suppose that  $f : A \rightarrow A$  is a mapping with  $f(0) = 0$  for which there exists a function  $\varphi : A \times A \times A \rightarrow [0, \infty)$  satisfying (2.31) and (3.4). Then, there exists a unique Jordan  $*$ -derivation  $\delta : A \rightarrow A$  such that

$$\|f(a) - \delta(a)\| \leq \tilde{\varphi}(a) \quad (3.13)$$

for all  $a \in A$ .

*Proof.* The rest of the proof is similar to the proofs of Theorems 2.4 and 3.3.  $\square$

**Corollary 3.6.** Suppose that  $f : A \rightarrow A$  is a mapping with  $f(0) = 0$  for which there exist constant  $\beta \geq 0$  and  $p_1, p_2, p_3 \in (1, \infty)$  such that

$$\left\| f\left(\frac{\lambda a + \lambda b}{k} + c^2\right) - \frac{\lambda}{k}f(a) - \frac{\lambda}{k}f(b) - c \circ f(c) - f(c^*) \circ c^* \right\| \leq \beta(\|a\|^{p_1} + \|b\|^{p_2} + \|c\|^{p_3}) \quad (3.14)$$

for all  $\lambda \in \mathbb{T}$  and all  $a, b, c \in A$ . Then, there exists a unique Jordan  $*$ -derivation  $\delta : A \rightarrow A$  such that

$$\|f(a) - \delta(a)\| \leq \frac{k\beta\|a\|^{p_1}}{k^{1-p_1} - 1} \quad (3.15)$$

for all  $a \in A$ .

*Proof.* Letting  $\varphi(a, b, c, d) := \beta(\|a\|^{p_1} + \|b\|^{p_2} + \|c\|^{p_3})$  in Theorem 3.5, we obtain the result.  $\square$

### Acknowledgment

This work was supported by Korea Research Foundation Grant KRF-2007-313-C00033.

### References

- [1] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Tracts in Pure and Applied Mathematics, no. 8, Interscience, New York, NY, USA, 1960.
- [2] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, no. 4, pp. 222–224, 1941.
- [3] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.
- [4] Th. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [5] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Journal of Functional Analysis*, vol. 46, no. 1, pp. 126–130, 1982.
- [6] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Bulletin des Sciences Mathématiques. Deuxième Série*, vol. 108, no. 4, pp. 445–446, 1984.
- [7] J. M. Rassias, "Solution of a problem of Ulam," *Journal of Approximation Theory*, vol. 57, no. 3, pp. 268–273, 1989.
- [8] J. M. Rassias, "Solution of a stability problem of Ulam," *Discussiones Mathematicae*, vol. 12, pp. 95–103, 1992.
- [9] J. M. Rassias, "Complete solution of the multi-dimensional problem of Ulam," *Discussiones Mathematicae*, vol. 14, pp. 101–107, 1994.
- [10] P. Găvruta, "An answer to a question of John M. Rassias concerning the stability of Cauchy equation," in *Advances in Equations and Inequalities*, Hadronic Mathematics, pp. 67–71, Hadronic Press, Palm Harbor, Fla, USA, 1999.

- [11] J. M. Rassias, "On the stability of the Euler-Lagrange functional equation," *Chinese Journal of Mathematics*, vol. 20, no. 2, pp. 185–190, 1992.
- [12] J. M. Rassias, "On the stability of the general Euler-Lagrange functional equation," *Demonstratio Mathematica*, vol. 29, no. 4, pp. 755–766, 1996.
- [13] M. J. Rassias and J. M. Rassias, "On the Ulam stability for Euler-Lagrange type quadratic functional equations," *The Australian Journal of Mathematical Analysis and Applications*, vol. 2, no. 1, pp. 1–10, 2005.
- [14] Th. M. Rassias, "Problem 16; 2, Report of the 27th International Symposium on Functional Equations," *Aequationes Mathematicae*, vol. 39, pp. 292–293, 1990.
- [15] Z. Gajda, "On stability of additive mappings," *International Journal of Mathematics and Mathematical Sciences*, vol. 14, no. 3, pp. 431–434, 1991.
- [16] Th. M. Rassias and P. Šemrl, "On the behavior of mappings which do not satisfy Hyers-Ulam stability," *Proceedings of the American Mathematical Society*, vol. 114, no. 4, pp. 989–993, 1992.
- [17] P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431–436, 1994.
- [18] S.-M. Jung, "On the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 204, no. 1, pp. 221–226, 1996.
- [19] S. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific, River Edge, NJ, USA, 2002.
- [20] D. H. Hyers, G. Isac, and Th. M. Rassias, *Stability of Functional Equations in Several Variables*, vol. 34 of *Progress in Nonlinear Differential Equations and Their Applications*, Birkhäuser, Boston, Mass, USA, 1998.
- [21] G. Isac and Th. M. Rassias, "Stability of  $\psi$ -additive mappings: applications to nonlinear analysis," *International Journal of Mathematics and Mathematical Sciences*, vol. 19, no. 2, pp. 219–228, 1996.
- [22] D. H. Hyers, G. Isac, and Th. M. Rassias, "On the asymptoticity aspect of Hyers-Ulam stability of mappings," *Proceedings of the American Mathematical Society*, vol. 126, no. 2, pp. 425–430, 1998.
- [23] D. G. Bourgin, "Classes of transformations and bordering transformations," *Bulletin of the American Mathematical Society*, vol. 57, pp. 223–237, 1951.
- [24] G. L. Forti, "Hyers-Ulam stability of functional equations in several variables," *Aequationes Mathematicae*, vol. 50, no. 1-2, pp. 143–190, 1995.
- [25] A. Gilányi, "On the stability of monomial functional equations," *Publicationes Mathematicae Debrecen*, vol. 56, no. 1-2, pp. 201–212, 2000.
- [26] A. Gilányi, "Eine zur Parallelogrammgleichung äquivalente Ungleichung," *Aequationes Mathematicae*, vol. 62, no. 3, pp. 303–309, 2001.
- [27] A. Gilányi, "On a problem by K. Nikodem," *Mathematical Inequalities & Applications*, vol. 5, no. 4, pp. 707–710, 2002.
- [28] P. M. Gruber, "Stability of isometries," *Transactions of the American Mathematical Society*, vol. 245, pp. 263–277, 1978.
- [29] K.-W. Jun and H.-M. Kim, "On the stability of Euler-Lagrange type cubic mappings in quasi-Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 332, no. 2, pp. 1335–1350, 2007.
- [30] K.-W. Jun, H.-M. Kim, and J. M. Rassias, "Extended Hyers-Ulam stability for Cauchy-Jensen mappings," *Journal of Difference Equations and Applications*, vol. 13, no. 12, pp. 1139–1153, 2007.
- [31] S.-M. Jung, "On the Hyers-Ulam stability of the functional equations that have the quadratic property," *Journal of Mathematical Analysis and Applications*, vol. 222, no. 1, pp. 126–137, 1998.
- [32] S.-M. Jung, "On the Hyers-Ulam-Rassias stability of a quadratic functional equation," *Journal of Mathematical Analysis and Applications*, vol. 232, no. 2, pp. 384–393, 1999.
- [33] H.-M. Kim, J. M. Rassias, and Y.-S. Cho, "Stability problem of Ulam for Euler-Lagrange quadratic mappings," *Journal of Inequalities and Applications*, vol. 2007, Article ID 10725, 15 pages, 2007.
- [34] S. Kurepa, "On the quadratic functional," *Publications de l'Institut Mathématique de l'Académie Serbe des Sciences et des Arts*, vol. 13, pp. 57–72, 1961.
- [35] Y.-S. Lee and S.-Y. Chung, "Stability of an Euler-Lagrange-Rassias equation in the spaces of generalized functions," *Applied Mathematics Letters*, vol. 21, no. 7, pp. 694–700, 2008.
- [36] P. Nakmahachalasint, "On the generalized Ulam-Găvruta-Rassias stability of mixed-type linear and Euler-Lagrange-Rassias functional equations," *International Journal of Mathematics and Mathematical Sciences*, vol. 2007, Article ID 63239, 10 pages, 2007.
- [37] C.-G. Park, "Lie  $*$ -homomorphisms between Lie  $C^*$ -algebras and Lie  $*$ -derivations on Lie  $C^*$ -algebras," *Journal of Mathematical Analysis and Applications*, vol. 293, no. 2, pp. 419–434, 2004.
- [38] C.-G. Park, "Homomorphisms between Lie  $JC^*$ -algebras and Cauchy-Rassias stability of Lie  $JC^*$ -algebra derivations," *Journal of Lie Theory*, vol. 15, no. 2, pp. 393–414, 2005.

- [39] C.-G. Park, "Homomorphisms between Poisson  $JC^*$ -algebras," *Bulletin of the Brazilian Mathematical Society*, vol. 36, no. 1, pp. 79–97, 2005.
- [40] C.-G. Park, "Hyers-Ulam-Rassias stability of a generalized Euler-Lagrange type additive mapping and isomorphisms between  $C^*$ -algebras," *Bulletin of the Belgian Mathematical Society*, vol. 13, no. 4, pp. 619–632, 2006.
- [41] C. Park, "Isomorphisms between quasi-Banach algebras," *Chinese Annals of Mathematics. Series B*, vol. 28, no. 3, pp. 353–362, 2007.
- [42] C. Park, Y. S. Cho, and M.-H. Han, "Functional inequalities associated with Jordan-von Neumann-type additive functional equations," *Journal of Inequalities and Applications*, vol. 2007, Article ID 41820, 13 pages, 2007.
- [43] C.-G. Park and J. Hou, "Homomorphisms between  $C^*$ -algebras associated with the Trif functional equation and linear derivations on  $C^*$ -algebras," *Journal of the Korean Mathematical Society*, vol. 41, no. 3, pp. 461–477, 2004.
- [44] C. G. Park, J. C. Hou, and S. Q. Oh, "Homomorphisms between  $JC^*$ -algebras and Lie  $C^*$ -algebras," *Acta Mathematica Sinica*, vol. 21, no. 6, pp. 1391–1398, 2005.
- [45] C. Park, J. Park, and J. Shin, "Hyers-Ulam-Rassias stability of quadratic functional equations in Banach modules over a  $C^*$ -algebra," *Chinese Annals of Mathematics. Series B*, vol. 24, no. 2, pp. 261–266, 2003.
- [46] A. Pietrzyk, "Stability of the Euler-Lagrange-Rassias functional equation," *Demonstratio Mathematica*, vol. 39, no. 3, pp. 523–530, 2006.
- [47] Th. M. Rassias, "The problem of S. M. Ulam for approximately multiplicative mappings," *Journal of Mathematical Analysis and Applications*, vol. 246, no. 2, pp. 352–378, 2000.
- [48] Th. M. Rassias, "On the stability of functional equations in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 251, no. 1, pp. 264–284, 2000.
- [49] Th. M. Rassias, "On the stability of functional equations and a problem of Ulam," *Acta Applicandae Mathematicae*, vol. 62, no. 1, pp. 23–130, 2000.
- [50] Th. M. Rassias, *Functional Equations, Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2003.
- [51] K. Ravi and M. Arunkumar, "On the Ulam-Găvruta-Rassias stability of the orthogonally Euler-Lagrange type functional equation," *International Journal of Applied Mathematics & Statistics*, vol. 7, no. Fe07, pp. 143–156, 2007.
- [52] J. Rätz, "On inequalities associated with the Jordan-von Neumann functional equation," *Aequationes Mathematicae*, vol. 66, no. 1-2, pp. 191–200, 2003.
- [53] W. Fechner, "Stability of a functional inequality associated with the Jordan-von Neumann functional equation," *Aequationes Mathematicae*, vol. 71, no. 1-2, pp. 149–161, 2006.
- [54] M. Amyari, F. Rahbarnia, and Gh. Sadeghi, "Some results on stability of extended derivations," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 2, pp. 753–758, 2007.
- [55] C. Baak, "Cauchy-Rassias stability of Cauchy-Jensen additive mappings in Banach spaces," *Acta Mathematica Sinica*, vol. 22, no. 6, pp. 1789–1796, 2006.





# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

