

## Research Article

# Fixed Points and Stability of the Cauchy Functional Equation in $C^*$ -Algebras

**Choonkil Park**

*Department of Mathematics, Hanyang University, Seoul 133–791, South Korea*

Correspondence should be addressed to Choonkil Park, baak@hanyang.ac.kr

Received 8 December 2008; Accepted 9 February 2009

Recommended by Tomas Domínguez Benavides

Using the fixed point method, we prove the generalized Hyers-Ulam stability of homomorphisms in  $C^*$ -algebras and Lie  $C^*$ -algebras and of derivations on  $C^*$ -algebras and Lie  $C^*$ -algebras for the Cauchy functional equation.

Copyright © 2009 Choonkil Park. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction and Preliminaries

The stability problem of functional equations originated from a question of Ulam [1] concerning the stability of group homomorphisms. Hyers [2] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' Theorem was generalized by Aoki [3] for additive mappings and by Th. M. Rassias [4] for linear mappings by considering an unbounded Cauchy difference. The paper of Th. M. Rassias [4] has provided a lot of influence in the development of what we call *generalized Hyers-Ulam stability* of functional equations. A generalization of the Th. M. Rassias theorem was obtained by Găvruta [5] by replacing the unbounded Cauchy difference by a general control function in the spirit of Th. M. Rassias' approach. The stability problems of several functional equations have been extensively investigated by a number of authors, and there are many interesting results concerning this problem (see [6–19]).

J. M. Rassias [20, 21] following the spirit of the innovative approach of Th. M. Rassias [4] for the unbounded Cauchy difference proved a similar stability theorem in which he replaced the factor  $\|x\|^p + \|y\|^p$  by  $\|x\|^p \cdot \|y\|^q$  for  $p, q \in \mathbb{R}$  with  $p + q \neq 1$  (see also [22] for a number of other new results).

We recall a fundamental result in fixed point theory.

Let  $X$  be a set. A function  $d : X \times X \rightarrow [0, \infty]$  is called a *generalized metric* on  $X$  if  $d$  satisfies

$$(1) \quad d(x, y) = 0 \text{ if and only if } x = y;$$

- (2)  $d(x, y) = d(y, x)$  for all  $x, y \in X$ ;
- (3)  $d(x, z) \leq d(x, y) + d(y, z)$  for all  $x, y, z \in X$ .

**Theorem 1.1** (see [23, 24]). *Let  $(X, d)$  be a complete generalized metric space and let  $J : X \rightarrow X$  be a strictly contractive mapping with Lipschitz constant  $L < 1$ . Then for each given element  $x \in X$ , either*

$$d(J^n x, J^{n+1} x) = \infty \quad (1.1)$$

for all nonnegative integers  $n$  or there exists a positive integer  $n_0$  such that

- (1)  $d(J^n x, J^{n+1} x) < \infty, \forall n \geq n_0$ ;
- (2) the sequence  $\{J^n x\}$  converges to a fixed point  $y^*$  of  $J$ ;
- (3)  $y^*$  is the unique fixed point of  $J$  in the set  $Y = \{y \in X \mid d(J^{n_0} x, y) < \infty\}$ ;
- (4)  $d(y, y^*) \leq (1/(1-L))d(y, Jy)$  for all  $y \in Y$ .

This paper is organized as follows. In Sections 2 and 3, using the fixed point method, we prove the generalized Hyers-Ulam stability of homomorphisms in  $C^*$ -algebras and of derivations on  $C^*$ -algebras for the Cauchy functional equation.

In Sections 4 and 5, using the fixed point method, we prove the generalized Hyers-Ulam stability of homomorphisms in Lie  $C^*$ -algebras and of derivations on Lie  $C^*$ -algebras for the Cauchy functional equation.

## 2. Stability of Homomorphisms in $C^*$ -Algebras

Throughout this section, assume that  $A$  is a  $C^*$ -algebra with norm  $\|\cdot\|_A$  and that  $B$  is a  $C^*$ -algebra with norm  $\|\cdot\|_B$ .

For a given mapping  $f : A \rightarrow B$ , we define

$$D_\mu f(x, y) := \mu f(x + y) - f(\mu x) - f(\mu y) \quad (2.1)$$

for all  $\mu \in \mathbb{T}^1 := \{\nu \in \mathbb{C} \mid |\nu| = 1\}$  and all  $x, y \in A$ .

Note that a  $\mathbb{C}$ -linear mapping  $H : A \rightarrow B$  is called a *homomorphism* in  $C^*$ -algebras if  $H$  satisfies  $H(xy) = H(x)H(y)$  and  $H(x^*) = H(x)^*$  for all  $x, y \in A$ .

We prove the generalized Hyers-Ulam stability of homomorphisms in  $C^*$ -algebras for the functional equation  $D_\mu f(x, y) = 0$ .

**Theorem 2.1.** *Let  $f : A \rightarrow B$  be a mapping for which there exists a function  $\varphi : A^2 \rightarrow [0, \infty)$  such that*

$$\|D_\mu f(x, y)\|_B \leq \varphi(x, y), \quad (2.2)$$

$$\|f(xy) - f(x)f(y)\|_B \leq \varphi(x, y), \quad (2.3)$$

$$\|f(x^*) - f(x)^*\|_B \leq \varphi(x, x) \quad (2.4)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y \in A$ . If there exists an  $L < 1$  such that  $\varphi(x, y) \leq 2L\varphi(x/2, y/2)$  for all  $x, y \in A$ , then there exists a unique  $C^*$ -algebra homomorphism  $H : A \rightarrow B$  such that

$$\|f(x) - H(x)\|_B \leq \frac{1}{2-2L}\varphi(x, x) \quad (2.5)$$

for all  $x \in A$ .

*Proof.* Consider the set

$$X := \{g : A \rightarrow B\}, \quad (2.6)$$

and introduce the *generalized metric* on  $X$ :

$$d(g, h) = \inf \{C \in \mathbb{R}_+ : \|g(x) - h(x)\|_B \leq C\varphi(x, x), \forall x \in A\}. \quad (2.7)$$

It is easy to show that  $(X, d)$  is complete.

Now we consider the linear mapping  $J : X \rightarrow X$  such that

$$Jg(x) := \frac{1}{2}g(2x) \quad (2.8)$$

for all  $x \in A$ .

By [23, Theorem 3.1],

$$d(Jg, Jh) \leq Ld(g, h) \quad (2.9)$$

for all  $g, h \in X$ .

Letting  $\mu = 1$  and  $y = x$  in (2.2), we get

$$\|f(2x) - 2f(x)\|_B \leq \varphi(x, x) \quad (2.10)$$

for all  $x \in A$ . So

$$\left\| f(x) - \frac{1}{2}f(2x) \right\|_B \leq \frac{1}{2}\varphi(x, x) \quad (2.11)$$

for all  $x \in A$ . Hence  $d(f, Jf) \leq 1/2$ .

By Theorem 1.1, there exists a mapping  $H : A \rightarrow B$  such that

(1)  $H$  is a fixed point of  $J$ , that is,

$$H(2x) = 2H(x) \quad (2.12)$$

for all  $x \in A$ . The mapping  $H$  is a unique fixed point of  $J$  in the set

$$Y = \{g \in X : d(f, g) < \infty\}. \quad (2.13)$$

This implies that  $H$  is a unique mapping satisfying (2.12) such that there exists  $C \in (0, \infty)$  satisfying

$$\|H(x) - f(x)\|_B \leq C\varphi(x, x) \quad (2.14)$$

for all  $x \in A$ .

(2)  $d(J^n f, H) \rightarrow 0$  as  $n \rightarrow \infty$ . This implies the equality

$$\lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n} = H(x) \quad (2.15)$$

for all  $x \in A$ .

(3)  $d(f, H) \leq (1/(1-L))d(f, Jf)$ , which implies the inequality

$$d(f, H) \leq \frac{1}{2-2L}. \quad (2.16)$$

This implies that the inequality (2.5) holds.

It follows from (2.2) and (2.15) that

$$\begin{aligned} \|H(x+y) - H(x) - H(y)\|_B &= \lim_{n \rightarrow \infty} \frac{1}{2^n} \|f(2^n(x+y)) - f(2^n x) - f(2^n y)\|_B \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \varphi(2^n x, 2^n y) = 0 \end{aligned} \quad (2.17)$$

for all  $x, y \in A$ . So

$$H(x+y) = H(x) + H(y) \quad (2.18)$$

for all  $x, y \in A$ .

Letting  $y = x$  in (2.2), we get

$$\mu f(2x) = f(\mu 2x) \quad (2.19)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x \in A$ . By a similar method to above, we get

$$\mu H(2x) = H(2\mu x) \quad (2.20)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x \in A$ . Thus one can show that the mapping  $H : A \rightarrow B$  is  $\mathbb{C}$ -linear.

It follows from (2.3) that

$$\begin{aligned}
\|H(xy) - H(x)H(y)\|_B &= \lim_{n \rightarrow \infty} \frac{1}{4^n} \|f(4^n xy) - f(2^n x)f(2^n y)\|_B \\
&\leq \lim_{n \rightarrow \infty} \frac{1}{4^n} \varphi(2^n x, 2^n y) \\
&\leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \varphi(2^n x, 2^n y) \\
&= 0
\end{aligned} \tag{2.21}$$

for all  $x, y \in A$ . So

$$H(xy) = H(x)H(y) \tag{2.22}$$

for all  $x, y \in A$ .

It follows from (2.4) that

$$\begin{aligned}
\|H(x^*) - H(x)^*\|_B &= \lim_{n \rightarrow \infty} \frac{1}{2^n} \|f(2^n x^*) - f(2^n x)^*\|_B \\
&\leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \varphi(2^n x, 2^n x) = 0
\end{aligned} \tag{2.23}$$

for all  $x \in A$ . So

$$H(x^*) = H(x)^* \tag{2.24}$$

for all  $x \in A$ .

Thus  $H : A \rightarrow B$  is a  $C^*$ -algebra homomorphism satisfying (2.5), as desired.  $\square$

**Corollary 2.2.** *Let  $0 < r < 1/2$  and  $\theta$  be nonnegative real numbers, and let  $f : A \rightarrow B$  be a mapping such that*

$$\|D_\mu f(x, y)\|_B \leq \theta \cdot \|x\|_A^r \cdot \|y\|_A^r, \tag{2.25}$$

$$\|f(xy) - f(x)f(y)\|_B \leq \theta \cdot \|x\|_A^r \cdot \|y\|_A^r, \tag{2.26}$$

$$\|f(x^*) - f(x)^*\|_B \leq \theta \|x\|_A^{2r} \tag{2.27}$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y \in A$ . Then there exists a unique  $C^*$ -algebra homomorphism  $H : A \rightarrow B$  such that

$$\|f(x) - H(x)\|_B \leq \frac{\theta}{2 - 4^r} \|x\|_A^{2r} \tag{2.28}$$

for all  $x \in A$ .

*Proof.* The proof follows from Theorem 2.1 by taking

$$\varphi(x, y) := \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \quad (2.29)$$

for all  $x, y \in A$ . Then  $L = 2^{2r-1}$  and we get the desired result.  $\square$

**Theorem 2.3.** *Let  $f : A \rightarrow B$  be a mapping for which there exists a function  $\varphi : A^2 \rightarrow [0, \infty)$  satisfying (2.2), (2.3), and (2.4). If there exists an  $L < 1$  such that  $\varphi(x, y) \leq (1/2)L\varphi(2x, 2y)$  for all  $x, y \in A$ , then there exists a unique  $C^*$ -algebra homomorphism  $H : A \rightarrow B$  such that*

$$\|f(x) - H(x)\|_B \leq \frac{L}{2-2L} \varphi(x, x) \quad (2.30)$$

for all  $x \in A$ .

*Proof.* We consider the linear mapping  $J : X \rightarrow X$  such that

$$Jg(x) := 2g\left(\frac{x}{2}\right) \quad (2.31)$$

for all  $x \in A$ .

It follows from (2.10) that

$$\left\| f(x) - 2f\left(\frac{x}{2}\right) \right\|_B \leq \varphi\left(\frac{x}{2}, \frac{x}{2}\right) \leq \frac{L}{2} \varphi(x, x) \quad (2.32)$$

for all  $x \in A$ . Hence,  $d(f, Jf) \leq L/2$ .

By Theorem 1.1, there exists a mapping  $H : A \rightarrow B$  such that

(1)  $H$  is a fixed point of  $J$ , that is,

$$H(2x) = 2H(x) \quad (2.33)$$

for all  $x \in A$ . The mapping  $H$  is a unique fixed point of  $J$  in the set

$$Y = \{g \in X : d(f, g) < \infty\}. \quad (2.34)$$

This implies that  $H$  is a unique mapping satisfying (2.33) such that there exists  $C \in (0, \infty)$  satisfying

$$\|H(x) - f(x)\|_B \leq C\varphi(x, x) \quad (2.35)$$

for all  $x \in A$ .

(2)  $d(J^n f, H) \rightarrow 0$  as  $n \rightarrow \infty$ . This implies the equality

$$\lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}\right) = H(x) \quad (2.36)$$

for all  $x \in A$ .

(3)  $d(f, H) \leq (1/(1-L))d(f, Jf)$ , which implies the inequality

$$d(f, H) \leq \frac{L}{2-2L}, \quad (2.37)$$

which implies that the inequality (2.30) holds.

The rest of the proof is similar to the proof of Theorem 2.1.  $\square$

**Corollary 2.4.** *Let  $r > 1$  and  $\theta$  be nonnegative real numbers, and let  $f : A \rightarrow B$  be a mapping satisfying (2.25), (2.26), and (2.27). Then there exists a unique  $C^*$ -algebra homomorphism  $H : A \rightarrow B$  such that*

$$\|f(x) - H(x)\|_B \leq \frac{\theta}{4^r - 2} \|x\|_A^{2r} \quad (2.38)$$

for all  $x \in A$ .

*Proof.* The proof follows from Theorem 2.3 by taking

$$\varphi(x, y) := \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \quad (2.39)$$

for all  $x, y \in A$ . Then  $L = 2^{1-2r}$  and we get the desired result.  $\square$

### 3. Stability of Derivations on $C^*$ -Algebras

Throughout this section, assume that  $A$  is a  $C^*$ -algebra with norm  $\|\cdot\|_A$ .

Note that a  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow A$  is called a *derivation* on  $A$  if  $\delta$  satisfies  $\delta(xy) = \delta(x)y + x\delta(y)$  for all  $x, y \in A$ .

We prove the generalized Hyers-Ulam stability of derivations on  $C^*$ -algebras for the functional equation  $D_\mu f(x, y) = 0$ .

**Theorem 3.1.** *Let  $f : A \rightarrow A$  be a mapping for which there exists a function  $\varphi : A^2 \rightarrow [0, \infty)$  such that*

$$\|D_\mu f(x, y)\|_A \leq \varphi(x, y), \quad (3.1)$$

$$\|f(xy) - f(x)y - xf(y)\|_A \leq \varphi(x, y) \quad (3.2)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y \in A$ . If there exists an  $L < 1$  such that  $\varphi(x, y) \leq 2L\varphi(x/2, y/2)$  for all  $x, y \in A$ . Then there exists a unique derivation  $\delta : A \rightarrow A$  such that

$$\|f(x) - \delta(x)\|_A \leq \frac{1}{2-2L}\varphi(x, x) \quad (3.3)$$

for all  $x \in A$ .

*Proof.* By the same reasoning as the proof of Theorem 2.1, there exists a unique involutive  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow A$  satisfying (3.3). The mapping  $\delta : A \rightarrow A$  is given by

$$\delta(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n} \quad (3.4)$$

for all  $x \in A$ .

It follows from (3.2) that

$$\begin{aligned} \|\delta(xy) - \delta(x)y - x\delta(y)\|_A &= \lim_{n \rightarrow \infty} \frac{1}{4^n} \|f(4^n xy) - f(2^n x) \cdot 2^n y - 2^n x f(2^n y)\|_A \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{4^n} \varphi(2^n x, 2^n y) \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \varphi(2^n x, 2^n y) \\ &= 0 \end{aligned} \quad (3.5)$$

for all  $x, y \in A$ . So

$$\delta(xy) = \delta(x)y + x\delta(y) \quad (3.6)$$

for all  $x, y \in A$ . Thus  $\delta : A \rightarrow A$  is a derivation satisfying (3.3).  $\square$

**Corollary 3.2.** Let  $0 < r < 1/2$  and  $\theta$  be nonnegative real numbers, and let  $f : A \rightarrow A$  be a mapping such that

$$\|D_\mu f(x, y)\|_A \leq \theta \cdot \|x\|_A^r \cdot \|y\|_A^r, \quad (3.7)$$

$$\|f(xy) - f(x)y - xf(y)\|_A \leq \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \quad (3.8)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y \in A$ . Then there exists a unique derivation  $\delta : A \rightarrow A$  such that

$$\|f(x) - \delta(x)\|_A \leq \frac{\theta}{2-4^r} \|x\|_A^{2r} \quad (3.9)$$

for all  $x \in A$ .



*Proof.* The proof follows from Theorem 3.1 by taking

$$\varphi(x, y) := \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \quad (3.10)$$

for all  $x, y \in A$ . Then  $L = 2^{2r-1}$  and we get the desired result.  $\square$

**Theorem 3.3.** *Let  $f : A \rightarrow A$  be a mapping for which there exists a function  $\varphi : A^2 \rightarrow [0, \infty)$  satisfying (3.1) and (3.2). If there exists an  $L < 1$  such that  $\varphi(x, y) \leq (1/2)L\varphi(2x, 2y)$  for all  $x, y \in A$ , then there exists a unique derivation  $\delta : A \rightarrow A$  such that*

$$\|f(x) - \delta(x)\|_A \leq \frac{L}{2-2L} \varphi(x, x) \quad (3.11)$$

for all  $x \in A$ .

*Proof.* The proof is similar to the proofs of Theorems 2.3 and 3.1.  $\square$

**Corollary 3.4.** *Let  $r > 1$  and  $\theta$  be nonnegative real numbers, and let  $f : A \rightarrow A$  be a mapping satisfying (3.7) and (3.8). Then there exists a unique derivation  $\delta : A \rightarrow A$  such that*

$$\|f(x) - \delta(x)\|_A \leq \frac{\theta}{4^r - 2} \|x\|_A^{2r} \quad (3.12)$$

for all  $x \in A$ .

*Proof.* The proof follows from Theorem 3.3 by taking

$$\varphi(x, y) := \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \quad (3.13)$$

for all  $x, y \in A$ . Then  $L = 2^{1-2r}$  and we get the desired result.  $\square$

#### 4. Stability of Homomorphisms in Lie $C^*$ -Algebras

A  $C^*$ -algebra  $\mathcal{C}$ , endowed with the Lie product  $[x, y] := (xy - yx)/2$  on  $\mathcal{C}$ , is called a *Lie  $C^*$ -algebra* (see [9–11]).

*Definition 4.1.* Let  $A$  and  $B$  be Lie  $C^*$ -algebras. A  $\mathbb{C}$ -linear mapping  $H : A \rightarrow B$  is called a *Lie  $C^*$ -algebra homomorphism* if  $H([x, y]) = [H(x), H(y)]$  for all  $x, y \in A$ .

Throughout this section, assume that  $A$  is a Lie  $C^*$ -algebra with norm  $\|\cdot\|_A$  and that  $B$  is a  $C^*$ -algebra with norm  $\|\cdot\|_B$ .

We prove the generalized Hyers-Ulam stability of homomorphisms in Lie  $C^*$ -algebras for the functional equation  $D_\mu f(x, y) = 0$ .

**Theorem 4.2.** *Let  $f : A \rightarrow B$  be a mapping for which there exists a function  $\varphi : A^2 \rightarrow [0, \infty)$  satisfying (2.2) such that*

$$\|f([x, y]) - [f(x), f(y)]\|_B \leq \varphi(x, y) \quad (4.1)$$

for all  $x, y \in A$ . If there exists an  $L < 1$  such that  $\varphi(x, y) \leq 2L\varphi(x/2, y/2)$  for all  $x, y \in A$ , then there exists a unique Lie  $C^*$ -algebra homomorphism  $H : A \rightarrow B$  satisfying (2.5).

*Proof.* By the same reasoning as the proof of Theorem 2.1, there exists a unique  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow A$  satisfying (2.5). The mapping  $H : A \rightarrow B$  is given by

$$H(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n} \quad (4.2)$$

for all  $x \in A$ .

It follows from (4.1) that

$$\begin{aligned} \|H([x, y]) - [H(x), H(y)]\|_B &= \lim_{n \rightarrow \infty} \frac{1}{4^n} \|f(4^n[x, y]) - [f(2^n x), f(2^n y)]\|_B \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{4^n} \varphi(2^n x, 2^n y) \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \varphi(2^n x, 2^n y) \\ &= 0 \end{aligned} \quad (4.3)$$

for all  $x, y \in A$ . So

$$H([x, y]) = [H(x), H(y)] \quad (4.4)$$

for all  $x, y \in A$ .

Thus  $H : A \rightarrow B$  is a Lie  $C^*$ -algebra homomorphism satisfying (2.5), as desired.  $\square$

**Corollary 4.3.** Let  $r < 1/2$  and  $\theta$  be nonnegative real numbers, and let  $f : A \rightarrow B$  be a mapping satisfying (2.25) such that

$$\|f([x, y]) - [f(x), f(y)]\|_B \leq \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \quad (4.5)$$

for all  $x, y \in A$ . Then there exists a unique Lie  $C^*$ -algebra homomorphism  $H : A \rightarrow B$  satisfying (2.28).

*Proof.* The proof follows from Theorem 4.2 by taking

$$\varphi(x, y) := \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \quad (4.6)$$

for all  $x, y \in A$ . Then  $L = 2^{2r-1}$  and we get the desired result.  $\square$

**Theorem 4.4.** Let  $f : A \rightarrow B$  be a mapping for which there exists a function  $\varphi : A^2 \rightarrow [0, \infty)$  satisfying (2.2) and (4.1). If there exists an  $L < 1$  such that  $\varphi(x, y) \leq (1/2)L\varphi(2x, 2y)$  for all  $x, y \in A$ , then there exists a unique Lie  $C^*$ -algebra homomorphism  $H : A \rightarrow B$  satisfying (2.30).

*Proof.* The proof is similar to the proofs of Theorems 2.3 and 4.2.  $\square$

**Corollary 4.5.** *Let  $r > 1$  and  $\theta$  be nonnegative real numbers, and let  $f : A \rightarrow B$  be a mapping satisfying (2.25) and (4.5). Then there exists a unique Lie  $C^*$ -algebra homomorphism  $H : A \rightarrow B$  satisfying (2.38).*

*Proof.* The proof follows from Theorem 4.4 by taking

$$\varphi(x, y) := \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \quad (4.7)$$

for all  $x, y \in A$ . Then  $L = 2^{1-2r}$  and we get the desired result.  $\square$

**Definition 4.6.** A  $C^*$ -algebra  $A$ , endowed with the Jordan product  $x \circ y := (xy + yx)/2$  for all  $x, y \in A$ , is called a *Jordan  $C^*$ -algebra* (see [25]).

**Definition 4.7.** Let  $A$  and  $B$  be Jordan  $C^*$ -algebras.

- (i) A  $\mathbb{C}$ -linear mapping  $H : A \rightarrow B$  is called a *Jordan  $C^*$ -algebra homomorphism* if  $H(x \circ y) = H(x) \circ H(y)$  for all  $x, y \in A$ .
- (ii) A  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow A$  is called a *Jordan derivation* if  $\delta(x \circ y) = x \circ \delta(y) + \delta(x) \circ y$  for all  $x, y \in A$ .

**Remark 4.8.** If the Lie products  $[\cdot, \cdot]$  in the statements of the theorems in this section are replaced by the Jordan products  $\cdot \circ \cdot$ , then one obtains Jordan  $C^*$ -algebra homomorphisms instead of Lie  $C^*$ -algebra homomorphisms.

## 5. Stability of Lie Derivations on $C^*$ -Algebras

**Definition 5.1.** Let  $A$  be a Lie  $C^*$ -algebra. A  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow A$  is called a *Lie derivation* if  $\delta([x, y]) = [\delta(x), y] + [x, \delta(y)]$  for all  $x, y \in A$ .

Throughout this section, assume that  $A$  is a Lie  $C^*$ -algebra with norm  $\|\cdot\|_A$ .

We prove the generalized Hyers-Ulam stability of derivations on Lie  $C^*$ -algebras for the functional equation  $D_\mu f(x, y) = 0$ .

**Theorem 5.2.** *Let  $f : A \rightarrow A$  be a mapping for which there exists a function  $\varphi : A^2 \rightarrow [0, \infty)$  satisfying (3.1) such that*

$$\|f([x, y]) - [f(x), y] - [x, f(y)]\|_A \leq \varphi(x, y) \quad (5.1)$$

for all  $x, y \in A$ . If there exists an  $L < 1$  such that  $\varphi(x, y) \leq 2L\varphi(x/2, y/2)$  for all  $x, y \in A$ . Then there exists a unique Lie derivation  $\delta : A \rightarrow A$  satisfying (3.3).

*Proof.* By the same reasoning as the proof of Theorem 2.1, there exists a unique involutive  $\mathbb{C}$ -linear mapping  $\delta : A \rightarrow A$  satisfying (3.3). The mapping  $\delta : A \rightarrow A$  is given by

$$\delta(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^n} \quad (5.2)$$

for all  $x \in A$ .

It follows from (5.1) that

$$\begin{aligned}
& \|\delta([x, y]) - [\delta(x), y] - [x, \delta(y)]\|_A \\
&= \lim_{n \rightarrow \infty} \frac{1}{4^n} \|f(4^n[x, y]) - [f(2^n x), 2^n y] - [2^n x, f(2^n y)]\|_A \\
&\leq \lim_{n \rightarrow \infty} \frac{1}{4^n} \varphi(2^n x, 2^n y) \\
&\leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \varphi(2^n x, 2^n y) \\
&= 0
\end{aligned} \tag{5.3}$$

for all  $x, y \in A$ . So

$$\delta([x, y]) = [\delta(x), y] + [x, \delta(y)] \tag{5.4}$$

for all  $x, y \in A$ . Thus  $\delta : A \rightarrow A$  is a derivation satisfying (3.3).  $\square$

**Corollary 5.3.** *Let  $0 < r < 1/2$  and  $\theta$  be nonnegative real numbers, and let  $f : A \rightarrow A$  be a mapping satisfying (3.7) such that*

$$\|f([x, y]) - [f(x), y] - [x, f(y)]\|_A \leq \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \tag{5.5}$$

for all  $x, y \in A$ . Then there exists a unique Lie derivation  $\delta : A \rightarrow A$  satisfying (3.9).

*Proof.* The proof follows from Theorem 5.2 by taking

$$\varphi(x, y) := \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \tag{5.6}$$

for all  $x, y \in A$ . Then  $L = 2^{2r-1}$  and we get the desired result.  $\square$

**Theorem 5.4.** *Let  $f : A \rightarrow A$  be a mapping for which there exists a function  $\varphi : A^2 \rightarrow [0, \infty)$  satisfying (3.1) and (5.1). If there exists an  $L < 1$  such that  $\varphi(x, y) \leq (1/2)L\varphi(2x, 2y)$  for all  $x, y \in A$ , then there exists a unique Lie derivation  $\delta : A \rightarrow A$  satisfying (3.11).*

*Proof.* The proof is similar to the proofs of Theorems 2.3 and 5.2.  $\square$

**Corollary 5.5.** *Let  $r > 1$  and  $\theta$  be nonnegative real numbers, and let  $f : A \rightarrow A$  be a mapping satisfying (3.7) and (5.5). Then there exists a unique Lie derivation  $\delta : A \rightarrow A$  satisfying (3.12).*

*Proof.* The proof follows from Theorem 5.4 by taking

$$\varphi(x, y) := \theta \cdot \|x\|_A^r \cdot \|y\|_A^r \tag{5.7}$$

for all  $x, y \in A$ . Then  $L = 2^{1-2r}$  and we get the desired result.  $\square$

*Remark 5.6.* If the Lie products  $[\cdot, \cdot]$  in the statements of the theorems in this section are replaced by the Jordan products  $\cdot \circ \cdot$ , then one obtains Jordan derivations instead of Lie derivations.

## Acknowledgment

This work was supported by Korea Research Foundation Grant KRF-2008-313-C00041.

## References

- [1] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Tracts in Pure and Applied Mathematics, no. 8, Interscience, New York, NY, USA, 1960.
- [2] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, no. 4, pp. 222–224, 1941.
- [3] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.
- [4] Th. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [5] P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431–436, 1994.
- [6] C.-G. Park, "On the stability of the linear mapping in Banach modules," *Journal of Mathematical Analysis and Applications*, vol. 275, no. 2, pp. 711–720, 2002.
- [7] C.-G. Park, "Modified Trif's functional equations in Banach modules over a  $C^*$ -algebra and approximate algebra homomorphisms," *Journal of Mathematical Analysis and Applications*, vol. 278, no. 1, pp. 93–108, 2003.
- [8] C.-G. Park, "On an approximate automorphism on a  $C^*$ -algebra," *Proceedings of the American Mathematical Society*, vol. 132, no. 6, pp. 1739–1745, 2004.
- [9] C.-G. Park, "Lie  $*$ -homomorphisms between Lie  $C^*$ -algebras and Lie  $*$ -derivations on Lie  $C^*$ -algebras," *Journal of Mathematical Analysis and Applications*, vol. 293, no. 2, pp. 419–434, 2004.
- [10] C.-G. Park, "Homomorphisms between Lie  $JC^*$ -algebras and Cauchy-Rassias stability of Lie  $JC^*$ -algebra derivations," *Journal of Lie Theory*, vol. 15, no. 2, pp. 393–414, 2005.
- [11] C.-G. Park, "Homomorphisms between Poisson  $JC^*$ -algebras," *Bulletin of the Brazilian Mathematical Society*, vol. 36, no. 1, pp. 79–97, 2005.
- [12] C.-G. Park, "Hyers-Ulam-Rassias stability of a generalized Euler-Lagrange type additive mapping and isomorphisms between  $C^*$ -algebras," *Bulletin of the Belgian Mathematical Society. Simon Stevin*, vol. 13, no. 4, pp. 619–632, 2006.
- [13] C. Park, "Fixed points and Hyers-Ulam-Rassias stability of Cauchy-Jensen functional equations in Banach algebras," *Fixed Point Theory and Applications*, vol. 2007, Article ID 50175, 15 pages, 2007.
- [14] C. Park, Y. S. Cho, and M.-H. Han, "Functional inequalities associated with Jordan-von Neumann-type additive functional equations," *Journal of Inequalities and Applications*, vol. 2007, Article ID 41820, 13 pages, 2007.
- [15] C. Park and J. Cui, "Generalized stability of  $C^*$ -ternary quadratic mappings," *Abstract and Applied Analysis*, vol. 2007, Article ID 23282, 6 pages, 2007.
- [16] C.-G. Park and J. Hou, "Homomorphisms between  $C^*$ -algebras associated with the Trif functional equation and linear derivations on  $C^*$ -algebras," *Journal of the Korean Mathematical Society*, vol. 41, no. 3, pp. 461–477, 2004.
- [17] Th. M. Rassias, "Problem 16; 2, Report of the 27th International Symposium on Functional Equations," *Aequationes Mathematicae*, vol. 39, no. 2-3, pp. 292–293, 309, 1990.
- [18] Th. M. Rassias, "The problem of S. M. Ulam for approximately multiplicative mappings," *Journal of Mathematical Analysis and Applications*, vol. 246, no. 2, pp. 352–378, 2000.
- [19] Th. M. Rassias, "On the stability of functional equations in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 251, no. 1, pp. 264–284, 2000.
- [20] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Journal of Functional Analysis*, vol. 46, no. 1, pp. 126–130, 1982.
- [21] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Bulletin des Sciences Mathématiques*, vol. 108, no. 4, pp. 445–446, 1984.

- [22] J. M. Rassias, "Solution of a problem of Ulam," *Journal of Approximation Theory*, vol. 57, no. 3, pp. 268–273, 1989.
- [23] L. Cădariu and V. Radu, "Fixed points and the stability of Jensen's functional equation," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 1, article 4, pp. 1–7, 2003.
- [24] J. B. Diaz and B. Margolis, "A fixed point theorem of the alternative, for contractions on a generalized complete metric space," *Bulletin of the American Mathematical Society*, vol. 74, no. 2, pp. 305–309, 1968.
- [25] R. J. Fleming and J. E. Jamison, *Isometries on Banach Spaces: Function Spaces*, vol. 129 of *Chapman & Hall/CRC Monographs and Surveys in Pure & Applied Mathematics*, Chapman & Hall/CRC, Boca Raton, Fla, USA, 2003.