



Clavicle midshaft fractures should not be considered an easy surgery: reduction and prebending

Young-Hoon Jo¹, Bong Gun Lee²

¹Department of Orthopaedic Surgery, Hanyang University Guri Hospital, Guri, Korea

²Department of Orthopaedic Surgery, Hanyang University Seoul Hospital, Seoul, Korea

Clavicle midshaft fractures are common, often requiring surgical fixation when displaced [1]. While this operation seems straightforward due to the thin soft tissue coverage and easy surgical approach, clavicle midshaft fracture surgery presents several challenges [2].

First, fracture reduction is typically difficult as the lateral fragment is displaced downward by the weight of the arm, and the medial fragment is displaced upward due to the action of the sternocleidomastoid muscle. These force vectors must be carefully considered during reduction. Second, soft tissue dissection must be minimized to promote fracture healing, limiting the field of view to one side of the clavicle and assessment of the angulation of the fracture. Last, the shape of the metal plate does not always match the shape of the clavicle as the curve and the dimension of the clavicle vary by patient; in addition, the location of the fracture is variable. Therefore, optimal placement of the plate can be challenging [3-5].

To address these issues, prebending the plate before surgery can be beneficial. Kim et al. [6] reported prebending the plate using a three-dimensional (3D) printed model based on a mirror image of the contralateral uninjured clavicle. That study appears to have been conducted with a deep understanding of the unique characteristics of clavicle midshaft fractures. I agree with the results of the study, suggesting that a preoperatively prebent plate based on a 3D-printed model that matches the clavicle shape can

be beneficial for fracture reduction in terms of operative time and clinical outcome.

In a study conducted by Leroux et al. [7], secondary surgery was performed in 24.6% of operatively repaired clavicles. The majority of these reoperations were for implant removal, with a higher removal rate in women compared to men. The study hypothesized that skin irritation, caused by undergarments crossing the clavicle plate and the overlying skin incision, could be a contributing factor. The higher rate of implant removal in women may support this hypothesis. The prebending method proposed by the authors may reduce skin irritation from the plates and patient discomfort, potentially reducing the need for implant removal surgeries [8].

The image reported as an axial image of the clavicle in the study is not a true axial image. A "true axial image" is a 90° vertical image compared to the anterior view. In the study of interest, the axial image is actually a cephalad image [9,10]. The study's strength is that it provides radiologic parameters (clavicle length and clavicle angle) using bilateral clavicles to evaluate the appropriateness of clavicle midshaft fracture reduction. Additionally, it offers clinical evidence of the cost-effective utility of preoperative prebending using a 3D-printed model [6]. Considering these strengths, I believe the results of the study can greatly benefit surgeons performing clavicle midshaft fracture surgery.

Received: November 6, 2023 Revised: November 11, 2023 Accepted: November 13, 2023

Correspondence to: Bong Gun Lee

Department of Orthopaedic Surgery, Hanyang University Seoul Hospital, 222-1 Wangsimni-ro, Seongdong-gu, Seoul 04763, Korea

Tel: +82-2-2290-8493, E-mail: orthdr@naver.com, ORCID: <https://orcid.org/0000-0002-4003-5529>

Copyright© 2023 Korean Shoulder and Elbow Society.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

NOTES

ORCID

Young-Hoon Jo <https://orcid.org/0000-0002-4299-2496>
Bong Gun Lee <https://orcid.org/0000-0002-4003-5529>

Author contributions

Conceptualization: YHJ. Writing – original draft: BGL. Writing – review & editing: YHJ.

Conflict of interest

Bong Gun Lee is an editorial board member of the journal but was not involved in the peer reviewer selection, evaluation, or decision process of this article. No other potential conflict of interest relevant to this article was reported.

Funding

None.

Data availability

None.

Acknowledgments

None.

REFERENCES

1. Ahmed AF, Salameh M, AlKhatib N, Elmhiregh A, Ahmed GO. Open reduction and internal fixation versus nonsurgical treatment in displaced midshaft clavicle fractures: a meta-analysis. *J Orthop Trauma* 2018;32:e276–83.
2. van der Meijden OA, Gaskill TR, Millett PJ. Treatment of clavicle fractures: current concepts review. *J Shoulder Elbow Surg* 2012;21:423–9.
3. Daruwalla ZJ, Courtis P, Fitzpatrick C, Fitzpatrick D, Mullett H. Anatomic variation of the clavicle: a novel three-dimensional study. *Clin Anat* 2010;23:199–209.
4. Bernat A, Huysmans T, Van Glabbeek F, Sijbers J, Gielen J, Van Tongel A. The anatomy of the clavicle: a three-dimensional cadaveric study. *Clin Anat* 2014;27:712–23.
5. King PR, Scheepers S, Ikram A. Anatomy of the clavicle and its medullary canal: a computed tomography study. *Eur J Orthop Surg Traumatol* 2014;24:37–42.
6. Kim H, Jung Y, Song HS. Plate prebending using a three-dimensional-printed model affords effective anatomical reduction in clavicular shaft fractures. *Clin Shoulder Elb* 2023;26:397–405.
7. Leroux T, Wasserstein D, Henry P, et al. Rate of and risk factors for reoperations after open reduction and internal fixation of midshaft clavicle fractures: a population-based study in Ontario, Canada. *J Bone Joint Surg Am* 2014;96:1119–25.
8. Wolf S, Chitnis AS, Manoranjith A, et al. Surgical treatment, complications, reoperations, and healthcare costs among patients with clavicle fracture in England. *BMC Musculoskeletal Disord* 2022;23:135.
9. Sharr JR, Mohammed KD. Optimizing the radiographic technique in clavicular fractures. *J Shoulder Elbow Surg* 2003;12:170–2.
10. Kim HN, Liu XN, Noh KC. Use of a real-size 3D-printed model as a preoperative and intraoperative tool for minimally invasive plating of comminuted midshaft clavicle fractures. *J Orthop Surg Res* 2015;10:91.