

Original Article



Associations of chronotype and insomnia with menstrual problems in newly employed nurses at university hospitals in the Republic of Korea

Han-Na Jung ¹, Dongwhan Suh ¹, Woo Chul Jeong ¹, Jia Ryu ²,
Yu-Mi Kim ^{3,4}, Seohyun Yoon ⁵, and Hyunjoo Kim ^{1*}

¹Department of Occupational and Environmental Medicine, Ewha Womans University Mokdong Hospital, Seoul, Korea

²Department of Occupational and Environmental Medicine, Catholic Kwandong University International St. Mary's Hospital, Incheon, Korea

³Department of Preventive Medicine, College of Medicine, Hanyang University, Seoul, Korea

⁴School of Public Health, Hanyang University, Seoul, Korea

⁵Ewha Medical Research Institute, Ewha Womans University School of Medicine, Seoul, Korea



Received: Feb 16, 2023

Revised: Jun 6, 2023

Accepted: Jun 30, 2023

Published online: Aug 4, 2023

*Correspondence:

Hyunjoo Kim

Department of Occupational and Environmental Medicine, Ewha Womans University Mokdong Hospital, 1071 Anyangcheon-ro, Yangcheon-gu, Seoul 07985, Korea.
Email: hj7121@gmail.com

Copyright © 2023 Korean Society of Occupational & Environmental Medicine
This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Han-Na Jung
<https://orcid.org/0000-0002-3194-9716>
Dongwhan Suh
<https://orcid.org/0000-0001-6327-5480>
Woo Chul Jeong
<https://orcid.org/0000-0003-1057-780X>
Jia Ryu
<https://orcid.org/0000-0002-7745-564X>
Yu-Mi Kim
<https://orcid.org/0000-0003-1123-8690>

ABSTRACT


Background: Dysmenorrhea and menstrual cycle changes occur in women working shifts. Circadian rhythm disruption and sleep disturbances associated with shift work leads to health problems. We identified chronotypes and the occurrence of insomnia among newly employed university hospital nurses and investigated the association of these factors with menstrual problems.

Methods: We conducted pre-placement health examinations for shift workers using self-reported questionnaires between 2018 and 2020. A total of 463 nurses were included in the study. Sociodemographic data, shift work experience, and information on insomnia were collected from health examination data. In addition, details regarding chronotype, dysmenorrhea, irregular and abnormal menstrual cycles, amenorrhea, and contraceptive use were obtained from the questionnaire. Multiple logistic regression analysis was performed to study the association between chronotype, insomnia, and menstrual problems after controlling for age, body mass index, contraceptive use, amenorrhea, and prior shift work.

Results: The prevalence rates of dysmenorrhea, irregular menstrual cycles, and longer menstrual cycles were 23.8%, 14.9%, and 4.1%, respectively. The risk of dysmenorrhea increased in the evening-type (odds ratio [OR]: 3.209; 95% confidence interval [CI]: 1.685–6.113) and those with insomnia (OR: 1.871; 95% CI: 1.074–3.261). Additionally, the risk of an irregular menstrual cycle (OR: 2.698; 95% CI: 1.167–6.237) increased in the evening-type, and the risk of a longer menstrual cycle (OR: 4.008; 95% CI: 1.354–11.864) increased in individuals with insomnia.

Conclusions: Our findings suggest that dysmenorrhea is promoted in the evening-type and insomnia individuals. There may be an increased risk of irregular menstrual cycles among evening-type nurses and an increased risk of longer menstrual cycles among those with insomnia. Therefore, factors such as evening-type and insomnia should be considered for the prevention of menstrual problems in women performing shift work.

Keywords: Circadian rhythm; Chronotype; Insomnia; Dysmenorrhea; Menstrual cycle; Nurse

Seohyun Yoon 

<https://orcid.org/0000-0003-4355-2596>

Hyunjoo Kim 

<https://orcid.org/0000-0002-9806-981X>

Abbreviations

BMI: body mass index; CI: confidence interval; ISI: Korean version of the Insomnia Severity Index; KtCS: Korean Translation of the Composite Scale; LH: luteinizing hormone; OR: odds ratio.

Funding

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (grant number: 2018R1A2B6004110).

Competing interests

The authors declare that they have no competing interests.

Authors contributions

Conceptualization: Jung HN, Kim H. Data curation: Jung HN, Ryu J. Formal analysis: Jung HN, Suh D, Kim H. Funding acquisition: Kim H. Investigation: Jung HN, Kim H. Writing - original draft: Jung HN, Kim H. Writing - review & editing: Jung HN, Suh D, Ryu J, Jeong WC, Kim YM, Yoon S, Kim H.

BACKGROUND

Health problems among nurses are directly related to patient safety. Therefore, managing health problems is a high priority to ensure optimal public health outcomes. Nurses who perform shift work may also experience chronic health problems. The adverse health effects of shift work may be mediated by behavioral mechanisms such as altered light exposure, diet, and sleep patterns. Altered light exposure, such as light at night and darkness during the day, could change hormones regulated by light and sleep (for example, prolactin, cortisol, and melatonin). Altered sleep patterns, such as short day sleeps and early wake times can disturb sleep. Shift work also increases risk behaviors and psychosocial stress. These factors cause physiological effects such as neuroendocrine stress, cardiac metabolic stress, altered immune function, cellular stress, and psychological effects such as cognitive impairment.¹ As a result, shift work increases nurses' risk of developing chronic diseases and falling victim to medical accidents.²

Circadian rhythm refers to the 24-hour internal clock in the brain that regulates cycles of alertness and sleepiness by responding to light changes in the environment.³ The chronotype is the expression of an individual's circadian rhythm, which is generally characterized by one's preference for activities at a particular time during a 24-hour cycle.⁴ For example, the morning-type prefers to sleep at night, wake up early, and become active in the morning, whereas the evening-type prefers the opposite.⁵

It is known that the determination of chronotype was mainly influenced by unchanging factors like age, sex, and polygenetic variation.⁶ It could be also influenced by environmental factors such as shift work. There is a view that health problems such as insomnia may occur when individual circadian rhythms are inconsistent with the social schedule.⁷ Chronotype shows variability, but the mechanism and extent are not yet clear.

Sleep disturbances include disorders of excessive somnolence; disorders of the sleep-wake schedule; dysfunctions associated with sleep, sleep stages, partial arousals; and insomnia.⁸ Regarding the relationship between chronotype and sleep, there are theories that chronotype contributes to sleep time and duration,⁷ and the theory that sleep characteristics are well explained by polygenetic factors including chronotype genes.⁹

Important health issues affecting nurses who perform shift work are dysmenorrhea and changes in menstrual cycles.¹⁰ Dysmenorrhea is defined as a painful cramp of uterine origin that occurs during menstruation.¹¹ The menstrual cycle is an essential life rhythm governed by the interaction of sex hormone levels and promotes stability through a feedback mechanism involving circadian rhythm and sleep,¹² and menstrual problems are not only important indicators of female reproductive health but are also significant factors affecting quality of life.¹³⁻¹⁵

Dysmenorrhea is a debilitating condition observed in many women that can result in reduced productivity¹⁶ and increased visits to medical institutions.¹⁵ Menstrual cycle changes are associated with gynecological diseases, including infertility.¹⁷ Furthermore, menstrual cycle changes increase the risk of severe health problems, such as breast cancer,¹⁸ type 2 diabetes,¹⁹ cardiovascular disease,²⁰ and premature mortality.²¹ Circadian rhythm and sleep are associated with menstrual problems. Although there have been reports of an increased risk of dysmenorrhea in the evening-type,^{22,23} these findings were not sufficient to establish a relationship between circadian rhythm and dysmenorrhea. The results of several

studies suggest an association between sleep disturbances and menstrual problems in the general population.^{24,25} In the case of shift work, previous studies have shown increased dysmenorrhea and changes in the amount of menstruation.^{26,27} However, few studies have examined the direct relationship between sleep disturbance and menstrual problems in female workers.

Although previous studies have suggested that menstrual cycle changes may be affected by circadian rhythms, the statistical significance of this association has been mixed.^{28,29} For example, in a previous study, the evening-type had a longer menstrual cycle than others, and the morning-type had a shorter duration of menstrual bleeding.²⁸ Therefore, it is necessary to further evaluate the association between circadian rhythms and menstrual cycle. Studies have reported an association between menstrual cycle irregularity and sleep disturbances such as short sleep duration²⁵ and insomnia symptoms.²⁴ Based on these findings, we further assessed the association between insomnia and abnormalities in menstrual cycle duration.

In this study, we aimed to identify the prevalence of menstrual problems and related factors, such as chronotype and insomnia, among nurses who were newly employed in hospitals. We also explored the relationship between these factors and menstrual problems such as dysmenorrhea and irregular menstrual cycles.

METHODS

Study population

The study population was newly employed at 2 university hospitals in Seoul, Korea. They underwent a pre-placement health examination and completed a self-reported questionnaire. Also, they underwent health checks before being assigned to shift work from March 1, 2018 to October 31, 2020. They had never worked shifts at the 2 university hospitals. However, some had experience working shifts at their previous hospitals, while the rest just graduated from nursing school and got their first jobs.

Of the 537 participants who completed the questionnaire, 463 were included in the final analysis. First, 18 people without body mass index (BMI) information were removed from the health examination data. Twenty-nine male workers were excluded because menstrual problems could not be assessed. Participants who reported that they had not menstruated in the past year were also excluded. Among the remaining 489 patients, 14 patients with irregular bleeding, who had less than 9 menstrual periods for 1 year in the menstrual questionnaire, were excluded as secondary causes could be suspected. Among the 475 patients, those without Dysmenorrhea score ($n = 1$) or circadian rhythm information ($n = 12$) were also excluded.

Study variables

Independent variables

To confirm the circadian rhythm type, the Korean Translation of the Composite Scale (KtCS), a structured measurement tool, was included in the questionnaire to evaluate the preferred times for various activities. There are 13 items in the KtCS; 3 items are scored on a 1–5-point scale, and the remaining ten items on a 1–4-point scale. The total score ranged from 13 to 55 points. A lower score indicated evening-type with a high preference for evening activities, and a higher score indicated morning-type with a greater morning preference.³⁰ Referring

to previous studies, a total score of ≤ 24 points (below the 10th percentile) was defined as the evening-type, and a total score of > 37 points (exceeding the 90th percentile) as the morning-type. The remaining score ranges were intermediate types.⁵ As mentioned above, the evening-type is related to dysmenorrhea^{22,23} and menstrual cycle changes²⁸; hence, we divided the chronotypes into the evening-type and non-evening-type (morning-type or intermediate-type). The points of participants who answered more than 80% of the question items in the questionnaire were corrected by converting the ratio.³¹ Four hundred forty-seven nurses responded to 100% of the KtCS questionnaire, 16 nurses whose KtCS score was corrected by ratio conversion because they responded between 80%–100%, and 12 were excluded because they responded less than 80%. The 12 nurses excluded from the study showed no significant differences in demographic characteristics compared with the nurses included in the analysis.

To evaluate insomnia, the Insomnia Severity Index, developed by Morin that includes a preplacement health examination, was applied.³² Seven items related to insomnia were used (severity of sleep onset, sleep maintenance, early morning awakening problems, sleep dissatisfaction, interference of sleep difficulties with daytime functioning, noticeability of sleep problems by others, and distress caused by sleep difficulties). Scores are added from 0 on a 4-point scale, and the distribution ranges from 0 to 28, with higher scores indicating severe insomnia. This study classified insomnia as a score of 8.

Dependent variables

Dysmenorrhea was defined as the pain intensity of menstruation for the past year measured on a numerical pain intensity scale of 0–10, and the top 25% score was 7 points (interquartile range 4–7 points). A score of < 7 and ≥ 7 was classified as absence and presence of dysmenorrhea, respectively. To evaluate the menstrual cycle, the question “What was your menstrual cycle over the past year?” was answered using the following 6 options: within 21 days, 21–25 days, 26–31 days, 32–39 days, ≥ 40 days, and irregular.²¹ Only those who answered “irregular” were defined as having an irregular menstrual cycle. A shorter menstrual cycle was defined as a menstrual period of < 21 days, and a longer menstrual cycle was defined as a menstrual period of > 40 days.¹⁹

Confounding variables

Shift work experience, age, BMI, smoking, alcohol consumption, exercise, and alcohol consumption were identified using preplacement health examination data. Prior shift work experience information was obtained in units of months from doctor’s history taken during pre-placement health examinations for shift workers. If the past shift work history was 0 month, it was classified as “never,” and if it was more than 1 month, it was classified as “ever.” According to World Health Organization standards, alcohol consumption is classified as non-drinker, moderate drinker (within 20 g of alcohol per time for women, drinking less than once a week), and risky drinkers. The degree of physical activity was classified as low, moderate, or high according to the Korean version of the International Physical Activity Questionnaire, a measurement tool for health checkups of the National Health Insurance.³³

The questionnaire identified contraception and amenorrhea experiences over the past year. Participants answered “yes” or “no” to the questions “Have you used contraception in the past year?” and “Have you not had menstruation for more than 90 days in the past year?”

Statistical analysis

All statistical analyses were performed using Statistical Package for the Social Sciences 22.0 (IBM SPSS: IBM SPSS version 22.0; IBM Corp., Armonk, NY, USA). Nurses were categorized as having a chronotype or insomnia. For basic comparisons of categorical variables, sociodemographic characteristics and the prevalence of menstrual problems were examined using χ^2 tests, except for shift work experience, contraception, and amenorrhea; these thresholds were evaluated using Fisher's exact test. Multiple logistic regression analysis was performed to evaluate the association between chronotype and insomnia and 3 menstrual problems: dysmenorrhea, irregular menstrual cycles, and abnormal menstrual cycles. Age, BMI, contraceptive experience in the past year, amenorrhea experience in the past year, and prior shift work experience were selected as confounding variables by reviewing previous studies on risk factors for dysmenorrhea³⁴ and menstrual cycle.³⁵ Multiple logistic regression analyses were performed on 420 nurses, excluding those with amenorrhea, to identify risk factors for menstrual irregularity and on 373 nurses without menstrual irregularity to identify risk factors for menstrual cycle abnormalities. The confounders included in these models were the aforementioned variables, with the exception of amenorrhea.

Ethics statement

This study was approved by the Institutional Review Board (IRB 2018-05-017-021) of the Ewha Womans University Mokdong Hospital and conducted in accordance with the Declaration of Helsinki of the World Medical Association. The study protocol was explained to the participants who provided written informed consent. Participants voluntarily agreed to complete the questionnaire.

RESULTS

The general characteristics of the study participants are summarized in **Table 1**. Of the 463 nurses, 16.2% reported having insomnia. Participants were aged < 25 years (37.4%), 25–29 years (50.1%), and > 30 years (12.5%). According to BMI, 10.8% had a BMI below 18.5, 75.2% had a BMI between 18.5 and 24.9, and 14.0% had a BMI > 30. There were 55.9% nurses with prior shift work experience and 23.8% with dysmenorrhea. Regarding the length of the menstrual cycle, the proportion of shorter cycles (< 21 days) was 0.9%, longer cycles (\geq 40 days) was 4.1%, and irregular menstrual cycles was 14.9%. The proportion of nurses with dysmenorrhea was higher in the evening-type group ($p < 0.001$). The proportion of nurses with prior shift work experience ($p < 0.001$), dysmenorrhea ($p = 0.003$), and late periods ($p = 0.004$) was higher in the insomnia group than in the non-insomnia group. In the 2 groups divided by evening-type, the proportions of general characteristics were not significantly different, except for dysmenorrhea. There were no statistically significant differences in age, obesity, contraceptive use, or amenorrhea between the nurses with and without insomnia.

The associations between dysmenorrhea, insomnia symptoms, and circadian rhythms are shown in **Table 2**. Age, contraceptive experience, and BMI, which are the individual risk factors related to dysmenorrhea³⁴ and menstrual cycle,³⁵ were adjusted. In addition, prior shift work experience, which may affect the occurrence of insomnia symptoms and circadian rhythms, was corrected.²² The risk of dysmenorrhea was higher in the evening group (odds ratio [OR]: 3.209; 95% confidence interval [CI]: 1.685–6.113). The risk of dysmenorrhea was 1.871 times (95% CI: 1.685–6.113) higher in the group with insomnia symptoms than in the group without insomnia.

Associations of chronotype and insomnia with menstrual problems

Table 1. Prevalence of the study population according to chronotype and insomnia among 463 newly employed nurses

Variables	Total	Chronotype		Insomnia (ISI \geq 8)	
		Non-evening-type ^a	Evening-type ^b	No	Yes
Total	463	416 (89.8)	47 (10.2)	388 (83.8)	75 (16.2)
Age (years)					
< 25	173 (37.4)	157 (37.7)	16 (34.0)	153 (39.4)	20 (26.7)
25–29	232 (50.1)	204 (49.0)	28 (59.6)	191 (49.2)	141 (54.7)
\geq 30	58 (12.5)	55 (13.2)	3 (6.4)	44 (11.3)	14 (18.7)
<i>p</i> -value		0.264 ^c		0.055 ^c	
Obesity (kg/m ²)					
0 < BMI < 18.5	50 (10.8)	45 (10.8)	5 (10.6)	44 (11.3)	6 (8.0)
18.5 \leq BMI < 25.0	348 (75.2)	311 (74.8)	37 (78.7)	291 (75.0)	57 (76.0)
BMI \geq 25.0	65 (14.0)	60 (14.4)	5 (10.6)	53 (13.7)	12 (16.0)
<i>p</i> -value		0.771 ^c		0.637 ^c	
Prior shift work experience					
Never (0 month)	204 (44.1)	185 (44.5)	19 (40.4)	185 (47.7)	19 (25.3)
Ever (\geq 1 month)	259 (55.9)	231 (55.5)	28 (59.6)	203 (52.3)	56 (74.7)
<i>p</i> -value		0.356 ^d		< 0.001 ^d	
Using contraception					
Yes	172 (37.1)	151 (36.3)	21 (44.7)	150 (38.7)	22 (29.3)
No	291 (62.9)	265 (63.7)	26 (55.3)	238 (61.3)	53 (70.7)
<i>p</i> -value		0.166 ^d		0.080 ^d	
Amenorrhea ^e					
No	420 (90.9)	377 (90.8)	43 (91.5)	353 (91.2)	67 (89.3)
Yes	42 (9.1)	38 (9.2)	4 (8.5)	34 (8.8)	8 (10.7)
<i>p</i> -value		0.571 ^d		0.369 ^d	
Dysmenorrhea					
No	353 (76.2)	328 (78.8)	25 (53.2)	306 (78.9)	47 (62.7)
Yes	110 (23.8)	88 (21.2)	22 (46.8)	82 (21.1)	28 (37.3)
<i>p</i> -value		< 0.001 ^c		0.003 ^c	
The length of menstrual cycle (days)					
Shorter (< 21)	4 (0.9)	3 (0.7)	1 (2.1)	4 (1.0)	0 (0.0)
Normal (21–39)	371 (80.1)	341 (82.0)	30 (63.8)	312 (80.4)	59 (78.7)
Longer (\geq 40)	19 (4.1)	15 (3.6)	4 (8.5)	10 (2.6)	9 (12.0)
Irregular	69 (14.9)	57 (13.7)	12 (25.5)	62 (16.0)	7 (9.3)
<i>p</i> -value		0.094 ^c		0.004 ^c	

Values are presented as number (%).

BMI: body mass index; KtCS: Korean Translation of the Composite Scale; ISI: Korean version of the Insomnia Severity Index.

^aMorning-type and intermediate-type (KtCS $>$ 24); ^bEvening-type (KtCS \leq 24) ^cTested using χ^2 test; ^dTested using Fisher's exact test; ^eNo menstruation more than 90 days in the past year.

Table 2. Association of chronotype and insomnia with dysmenorrhea among 463 newly employed nurses

Variables	Total	Case (%)	Dysmenorrhea	
			Crude OR (95% CI)	Adjusted ^a OR (95% CI)
Chronotype				
Non-evening-type ^b	416	88 (21.2)	Ref.	Ref.
Evening-type ^c	9	22 (46.8)	3.280 (1.766–6.094)	3.209 (1.685–6.113)
Insomnia (ISI \geq 8)				
No	388	82 (21.1)	Ref.	Ref.
Yes	75	28 (37.3)	2.223 (1.312–3.768)	1.871 (1.074–3.261)

CI: confidence interval; OR: odds ratio; ISI: Korean version of the Insomnia Severity Index; KtCS: Korean Translation of the Composite Scale.

^aAdjusted for age, body mass index (< 18.5 kg/m²), no contraception in the past year, amenorrhea, and prior shift work experience; ^bMorning-type and intermediate-type (KtCS $>$ 24) ^cEvening-type (KtCS \leq 24).

The occurrence of irregular menstrual cycles was analyzed, except for individuals suspected of having amenorrhea (Table 3) for whom the menstrual cycle could not be evaluated. We adjusted for confounding variables such as age, BMI, previous year's contraceptive experience, and prior shift work experience. We then analyzed irregular menstrual cycles in 420 nurses and found that in the evening-type, the risk of irregular menstrual cycles was 2.698 times (95% CI: 1.167–6.237) higher than that in other groups.

Associations of chronotype and insomnia with menstrual problems

Table 3. Association of chronotype and insomnia with irregular menstrual cycle among 420^a newly employed nurses without amenorrhea

Variables	Total	Case (%)	Irregular menstrual cycle	
			Crude OR (95% CI)	Adjusted ^b OR (95% CI)
Chronotype				
Non-evening-type ^c	377	35 (9.3)	Ref.	Ref.
Evening-type ^d	43	9 (20.9)	2.587 (1.147–5.831)	2.698 (1.167–6.237)
Insomnia (ISI ≥ 8)				
No	353	40 (11.3)	Ref.	Ref.
Yes	67	4 (6.0)	0.497 (0.172–1.438)	0.442 (0.149–1.312)

CI: confidence interval; OR: odds ratio; ISI: Korean version of the Insomnia Severity Index; KtCS: Korean Translation of the Composite Scale.

^aForty-two patients with unusual amenorrhea findings were excluded; ^bAdjusted for age, body mass index (≥ 25.0 kg/m²), no contraception in the past year, and prior shift work experience; ^cMorning-type and intermediate-type (KtCS > 24); ^dEvening-type (KtCS ≤ 24).

Table 4. Association of chronotype and insomnia with abnormal menstrual cycle among 376^a newly employed nurses without amenorrhea or irregular menstrual cycle

Variables	Total	Case (%)	Longer menstrual cycle (> 40)	
			Crude OR (95% CI)	Adjusted ^b OR (95% CI)
Chronotype				
Non-evening-type ^c	330	12 (3.6)	Ref.	Ref.
Evening-type ^d	30	4 (13.3)	3.759 (1.275–11.079)	3.203 (0.910–11.267)
Insomnia (ISI ≥ 8)				
No	304	9 (3.0)	Ref.	Ref.
Yes	56	7 (12.5)	2.885 (1.102–7.550)	4.008 (1.354–11.864)

CI: confidence interval; OR: odds ratio; ISI: Korean version of the Insomnia Severity Index; KtCS: Korean Translation of the Composite Scale.

^aEighty-seven patients with unusual findings of amenorrhea or irregular menstrual cycles were excluded; ^bAdjusted for age, body mass index (≥ 25.0 kg/m²), no contraception in the past year, and prior shift work experience; ^cMorning-type and intermediate-type (KtCS > 24) ^dEvening-type (KtCS ≤ 24).

As shown in **Table 4**, 87 patients with unusual amenorrhea or irregular menstrual cycles were excluded. The remaining 376 nurses who had regular cycles were analyzed for abnormalities. For insomnia, the risk of a longer cycle was 4.008 times higher (95% CI: 1.354–11.864) than that of the other groups.

DISCUSSION

After controlling for age, BMI, contraceptive use, and prior shift work experience, we found that the risk of dysmenorrhea was associated with insomnia symptoms and evening-type. Additionally, an increased risk of an irregular menstrual cycle was independently associated with the evening-type, and an increased risk of an abnormal menstrual cycle > 40 days was independently associated with insomnia.

Two cross-sectional studies have reported an association between circadian rhythm and dysmenorrhea. These studies consistently showed that there was more dysmenorrhea in the evening-type than in other chronotypes.^{23,36} A possible explanation for the increased risk of dysmenorrhea in the evening-type was sleep disruption.^{37,38} The altered sleep timing of the evening-type might be related to the disruption of the 24-hour clock, which is key to reproductive hormone regulation. This suggestion does not support our findings as circadian rhythms are not independently associated with the risk of dysmenorrhea. Another possible explanation was that evening-type used psychoactive substances more frequently.^{39,40} However, a cross-sectional study conducted on female students in Japan reported contrasting findings. In this study, students who frequently experienced dysmenorrhea had more evening-type dysmenorrhea than those who did not. They speculated that the threshold of menstrual pain could be lowered through depressive mood symptoms in the evening-type.

The finding that insomnia was independently associated with the risk of dysmenorrhea is consistent with those of several studies. Among related factors, reduced melatonin levels might explain both insomnia- and evening-type symptoms. Insomnia and evening-type conditions commonly interfere with adequate sleep at night.⁴¹ In insomnia and evening-type insomnia, melatonin synthesized in the body by the stimulation of darkness⁴² may be present at a low concentration. Animal experiments have shown that melatonin inhibits uterine contractions and exerts analgesic effects.⁴³ Additionally, plasma melatonin levels increase during menstruation.⁴⁴ If melatonin secretion, which contributes to the control of dysmenorrhea, was reduced, it might have affected dysmenorrhea. Other factors include the perturbed actions of several hormones such as estrogen, progesterone, prolactin, and growth hormones.⁴⁵ In addition, many theories have traditionally reported that insomnia results from dysmenorrhea.^{38,46-48}

In this study, the risk of an irregular menstrual cycle and abnormal cycle (≥ 40) increased in the evening-type; however, the finding of abnormal cycle was not statistically significant. These results suggest that menstrual cycle-related problems tend to increase in the evening-type. Several studies have supported the association between the evening-type and these problems. They obtained mixed results for irregular⁴⁹ and abnormal cycles. These results can be explained by differences in reproductive hormones, an increase in melatonin due to sleep debt, and weak zeitgeber effects.

First, 2 cross-sectional studies supported the differences in reproductive hormone levels. In one study, the menstrual cycle was one day longer in the evening than in the morning. In contrast, the luteinizing hormone (LH) peak appeared one day later. If the circadian rhythm changes, mutations can occur in circadian clock genes. Alternatively, the central and peripheral oscillators are not synchronized. This may affect the regulation of reproductive hormones in ovaries and uterus. Second, we hypothesized that an imbalance in melatonin causes changes like sleep debt in the evening-type menstrual cycle. Increased daytime sleep and decreased darkness in the evening can cause an imbalance in melatonin levels.⁵⁰ One study focused on the fact that an imbalance in melatonin can suppress the secretion of gonadotropin-releasing hormones and affect ovulation. In a study in which reproductive hormones were measured during the day and night depending on the season, nighttime serum melatonin in the follicular phase increased significantly in winter compared with that in summer. Nighttime serum LH levels were higher in summer than in winter. This could affect fertility by regulating LH secretion based on the melatonin levels.⁵¹ Finally, weak zeitgebers, which could cause problems in the evening-type menstrual cycle, refer to the weakening of the signal strength that transmits 24-hour changes to the human body in the external environment. The evening-type can be in a state of weak zeitgeber, with longer artificial light exposure during the night and shorter sunlight exposure during the day. A related clinical study reported that in weak zeitgebers, an additional delay in the circadian rhythm could occur, which could increase the effect of melatonin.⁵²

Finally, our findings demonstrated the potential involvement of insomnia in long menstrual cycles. Additionally, although not statistically significant, the risk of irregular cycles reduced in the presence of insomnia. These conflicting results seem to have reduced the statistical stability because the number of cases of insomnia and irregular or long cycles was less than 1% in our study population. Although a few studies have examined the association between insomnia and long cycles, some studies have reported an association between insomnia and irregular cycles. In a 12-month follow-up study, after adjusting for confounding variables,

the probability of a new irregular cycle increased compared to those without insomnia at baseline.⁵³ Insomnia causes estradiol levels to remain elevated,^{54,55} and LH pulse frequency is associated with wakefulness.⁵⁶ The desynchronization of reproductive hormones associated with ovulation may result in sleep disturbances. Hormonal asynchrony related to successful ovulation can be caused by sleep disturbance.

From the above results, it can be seen that chronotype and insomnia are independently related to menstrual health problems. As mentioned in the introduction, it is known that chronotype can affect sleep time and sleep duration, so the interaction between chronotype and insomnia was examined in multiple logistic regression analysis. Although not presented in the result, no statistically significant interaction between chronotype and insomnia was observed for any of the 3 menstrual problems.

This study had several limitations that should be considered when interpreting our findings. First, our study was cross-sectional; hence, causal relationships could not be determined. However, chronotypes are determined by unchanging characteristics, such as genes and age.⁶ Nurses with secondary causes of menstrual problems were also excluded. We believe that chronotypes are more likely to precede menstrual disorders. Longitudinal data will be helpful in examining the associations between chronotype, insomnia, dysmenorrhea, and menstrual cycle.

Second, variables such as chronotype, insomnia, and menstrual problems were measured using self-report questionnaires. Self-reported data are susceptible to recall bias or denial of problem. Measurement tools for chronotype⁵ and insomnia³² have been validated in several studies. Menstrual problems are less likely to be affected by recall bias because menarche problems may occur monthly.⁵⁷ Studies that have reported menstrual problems have mainly used self-reports, whereas studies using hormone measurements are rare.

Finally, although randomized sampling was used to ensure the representativeness of all nurses working at a tertiary hospital, our study findings might not be generalizable. Because the sample size was large, the effect of the selection bias was thought to be less significant.

There have been studies that analyzed the relationship between chronotype and insomnia,^{58,59} or insomnia and specific menstrual problem such as dysmenorrhea³⁷ and menstrual cycles.⁵³⁻⁵⁶ But few studies had analyzed the relationship between chronotype, insomnia, and menstrual problems such as dysmenorrhea or menstrual cycle together. Our study population is a random sample extracted from newly employed nurses in a university hospital, so it is difficult to generalize it to working women. It might be possible to apply our result to young nurses similar to our study population. In terms of occupational medicine, it is important to establish measures to protect workers with vulnerable characteristics from health problems related to hazards that workers might be exposed to. Therefore our research results can contribute to protecting these workers.

CONCLUSIONS

Our study found that the risk of dysmenorrhea increased among nurses who were of evening-type or had insomnia. The risk of irregular menstrual cycles increased in the evening-type, and the risk of a longer cycle was higher in those with insomnia. Since this study is a cross-sectional study, it is not possible to establish a causal relationship. However, the association

between chronotype, insomnia and menstrual problems observed in our study may provide a hypothesis when designing studies on menstrual problems in nurses.

ACKNOWLEDGEMENTS

The authors thank the staff of the Ewha Medical Examination Center of Ewha Womans University Mokdong Hospital for their assistance in this study.

REFERENCES

1. Kecklund G, Axelsson J. Health consequences of shift work and insufficient sleep. *BMJ* 2016;355:i5210.
[PUBMED](#) | [CROSSREF](#)
2. Wickwire EM, Geiger-Brown J, Scharf SM, Drake CL. Shift work and shift work sleep disorder: clinical and organizational perspectives. *Chest* 2017;151(5):1156-72.
[PUBMED](#) | [CROSSREF](#)
3. Turek FW. From circadian rhythms to clock genes in depression. *Int Clin Psychopharmacol* 2007;22 Suppl 2:S1-8.
[PUBMED](#) | [CROSSREF](#)
4. Hurley S, Goldberg D, Von Behren J, Clague DeHart J, Wang S, Reynolds P. Chronotype and postmenopausal breast cancer risk among women in the California Teachers Study. *Chronobiol Int* 2019;36(11):1504-14.
[PUBMED](#) | [CROSSREF](#)
5. Kook SH, Yoon JS, Lee HY. Cross validation of the Korean translation of composite Scale (KtCS) to measure morningness-eveningness. *J Korean Neuropsychiatr Assoc* 1999;38(2):297-305.
6. Lee JH, Kim IS, Kim SJ, Wang W, Duffy JF. Change in individual chronotype over a lifetime: a retrospective study. *Sleep Med Rev* 2011;2(2):48-53.
[CROSSREF](#)
7. Gentry NW, Ashbrook LH, Fu YH, Ptáček LJ. Human circadian variations. *J Clin Invest* 2021;131(16):e148282.
[PUBMED](#) | [CROSSREF](#)
8. Walker HK, Hall WD, Hurst JW. The history, physical, and laboratory examinations. *Clinical Methods*. 3rd ed. Boston, MA, USA: Butterworths; 1990.
9. Perkiö A, Merikanto I, Kantöjärvi K, Paunio T, Sinnott-Armstrong N, Jones SE, et al. Portability of polygenic risk scores for sleep duration, insomnia and chronotype in 33,493 individuals. *Clocks Sleep* 2022;5(1):10-20.
[PUBMED](#) | [CROSSREF](#)
10. Chang WP, Chang YP. Meta-analysis comparing menstrual regularity and dysmenorrhea of women working rotating shifts and fixed day shifts. *J Womens Health (Larchmt)* 2021;30(5):722-30.
[PUBMED](#) | [CROSSREF](#)
11. Szmídt MK, Granda D, Sicinska E, Kaluza J. Primary dysmenorrhea in relation to oxidative stress and antioxidant status: a systematic review of case-control studies. *Antioxidants (Basel)* 2020;9(10):994.
[PUBMED](#) | [CROSSREF](#)
12. Draper CF, Duisters K, Weger B, Chakrabarti A, Harms AC, Brennan L, et al. Menstrual cycle rhythmicity: metabolic patterns in healthy women. *Sci Rep* 2018;8(1):14568.
[PUBMED](#) | [CROSSREF](#)
13. Pogodina A, Dolgikh O, Astakhova T, Klimkina J, Khramova E, Rychkova L. Health-related quality of life and menstrual problems in adolescents. *J Paediatr Child Health* 2022;58(6):1028-32.
[PUBMED](#) | [CROSSREF](#)
14. Shimamoto K, Hirano M, Wada-Hiraie O, Goto R, Osuga Y. Examining the association between menstrual symptoms and health-related quality of life among working women in Japan using the EQ-5D. *BMC Womens Health* 2021;21(1):325.
[PUBMED](#) | [CROSSREF](#)
15. Titiyayo A, Agunbiade OM, Banjo O, Lawani A. Menstrual discomfort and its influence on daily academic activities and psychosocial relationship among undergraduate female students in Nigeria. *Tanzan J Health Res* 2009;11(4):181-8.
[PUBMED](#) | [CROSSREF](#)

16. Sharma A, Taneja DK, Sharma P, Saha R. Problems related to menstruation and their effect on daily routine of students of a medical college in Delhi, India. *Asia Pac J Public Health* 2008;20(3):234-41.
[PUBMED](#) | [CROSSREF](#)
17. Small CM, Manatunga AK, Klein M, Dominguez CE, Feigelson HS, McChesney R, et al. Menstrual cycle variability and the likelihood of achieving pregnancy. *Rev Environ Health* 2010;25(4):369-78.
[PUBMED](#) | [CROSSREF](#)
18. Olsson HL, Olsson ML. The menstrual cycle and risk of breast cancer: a review. *Front Oncol* 2020;10:21.
[PUBMED](#) | [CROSSREF](#)
19. Solomon CG, Hu FB, Dunaif A, Rich-Edwards J, Willett WC, Hunter DJ, et al. Long or highly irregular menstrual cycles as a marker for risk of type 2 diabetes mellitus. *JAMA* 2001;286(19):2421-6.
[PUBMED](#) | [CROSSREF](#)
20. Solomon CG, Hu FB, Dunaif A, Rich-Edwards JE, Stampfer MJ, Willett WC, et al. Menstrual cycle irregularity and risk for future cardiovascular disease. *J Clin Endocrinol Metab* 2002;87(5):2013-7.
[PUBMED](#) | [CROSSREF](#)
21. Wang YX, Arvizu M, Rich-Edwards JW, Stuart JJ, Manson JE, Missmer SA, et al. Menstrual cycle regularity and length across the reproductive lifespan and risk of premature mortality: prospective cohort study. *BMJ* 2020;371:m3464.
[PUBMED](#) | [CROSSREF](#)
22. Rosa D, Terzoni S, Dellafiore F, Destrebecq A. Systematic review of shift work and nurses' health. *Occup Med (Lond)* 2019;69(4):237-43.
[PUBMED](#) | [CROSSREF](#)
23. Negriff S, Dorn LD. Morningness/eveningness and menstrual symptoms in adolescent females. *J Psychosom Res* 2009;67(2):169-72.
[PUBMED](#) | [CROSSREF](#)
24. Xing X, Xue P, Li SX, Zhou J, Tang X. Sleep disturbance is associated with an increased risk of menstrual problems in female Chinese university students. *Sleep Breath* 2020;24(4):1719-27.
[PUBMED](#) | [CROSSREF](#)
25. Nam GE, Han K, Lee G. Association between sleep duration and menstrual cycle irregularity in Korean female adolescents. *Sleep Med* 2017;35:62-6.
[PUBMED](#) | [CROSSREF](#)
26. Chung FF, Yao CC, Wan GH. The associations between menstrual function and life style/working conditions among nurses in Taiwan. *J Occup Health* 2005;47(2):149-56.
[PUBMED](#) | [CROSSREF](#)
27. Uehata T, Sasakawa N. The fatigue and maternity disturbances of night workwomen. *J Hum Ergol (Tokyo)* 1982;11 Suppl:465-74.
[PUBMED](#)
28. Toffol E, Merikanto I, Lahti T, Luoto R, Heikinheimo O, Partonen T. Evidence for a relationship between chronotype and reproductive function in women. *Chronobiol Int* 2013;30(6):756-65.
[PUBMED](#) | [CROSSREF](#)
29. Michels KA, Mendola P, Schliep KC, Yeung EH, Ye A, Dunietz GL, et al. The influences of sleep duration, chronotype, and nightwork on the ovarian cycle. *Chronobiol Int* 2020;37(2):260-71.
[PUBMED](#) | [CROSSREF](#)
30. Yun JA, Ahn YS, Jeong KS, Joo EJ, Choi KS. The relationship between chronotype and sleep quality in Korean firefighters. *Clin Psychopharmacol Neurosci* 2015;13(2):201-8.
[PUBMED](#) | [CROSSREF](#)
31. McDowell I. *Measuring Health: A Guide to Rating Scales and Questionnaires*. 3rd ed. Oxford, UK: Oxford University Press; 2006.
32. Bastien CH, Vallières A, Morin CM. Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Med* 2001;2(4):297-307.
[PUBMED](#) | [CROSSREF](#)
33. Chun MY. Validity and reliability of Korean version of international physical activity questionnaire short form in the elderly. *Korean J Fam Med* 2012;33(3):144-51.
[PUBMED](#) | [CROSSREF](#)
34. Ju H, Jones M, Mishra G. The prevalence and risk factors of dysmenorrhea. *Epidemiol Rev* 2014;36(1):104-13.
[PUBMED](#) | [CROSSREF](#)
35. Bae J, Park S, Kwon JW. Factors associated with menstrual cycle irregularity and menopause. *BMC Womens Health* 2018;18(1):36.
[PUBMED](#) | [CROSSREF](#)

36. Takeuchi H, Oishi T, Harada T. Association between morningness-eveningness preference and mental/physical premenstrual symptoms in Japanese females 12 to 31 years of age. *Chronobiol Int* 2005;22(6):1055-68.
[PUBMED](#) | [CROSSREF](#)
37. Labyak S, Lava S, Turek F, Zee P. Effects of shiftwork on sleep and menstrual function in nurses. *Health Care Women Int* 2002;23(6-7):703-14.
[PUBMED](#) | [CROSSREF](#)
38. Baker FC, Driver HS, Rogers GG, Paiker J, Mitchell D. High nocturnal body temperatures and disturbed sleep in women with primary dysmenorrhea. *Am J Physiol* 1999;277(6):E1013-21.
[PUBMED](#) | [CROSSREF](#)
39. Harlow SD, Park M. A longitudinal study of risk factors for the occurrence, duration and severity of menstrual cramps in a cohort of college women. *Br J Obstet Gynaecol* 1996;103(11):1134-42.
[PUBMED](#) | [CROSSREF](#)
40. Teperi J, Rimpelä M. Menstrual pain, health and behaviour in girls. *Soc Sci Med* 1989;29(2):163-9.
[PUBMED](#) | [CROSSREF](#)
41. Giannotti F, Cortesi F, Sebastiani T, Ottaviano S. Circadian preference, sleep and daytime behaviour in adolescence. *J Sleep Res* 2002;11(3):191-9.
[PUBMED](#) | [CROSSREF](#)
42. Srinivasan V, Spence WD, Pandi-Perumal SR, Zakharia R, Bhatnagar KP, Brzezinski A. Melatonin and human reproduction: shedding light on the darkness hormone. *Gynecol Endocrinol* 2009;25(12):779-85.
[PUBMED](#) | [CROSSREF](#)
43. Hertz-Eshel M, Rahamimoff R. Effect of melatonin on uterine contractility. *Life Sci* 1965;4(14):1367-72.
[PUBMED](#) | [CROSSREF](#)
44. Barron ML. Light exposure, melatonin secretion, and menstrual cycle parameters: an integrative review. *Biol Res Nurs* 2007;9(1):49-69.
[PUBMED](#) | [CROSSREF](#)
45. Arafa A, Mahmoud O, Abu Salem E, Mohamed A. Association of sleep duration and insomnia with menstrual symptoms among young women in Upper Egypt. *Middle East Current Psychiatry* 2020;27(1):2.
[CROSSREF](#)
46. Woosley JA, Lichstein KL. Dysmenorrhea, the menstrual cycle, and sleep. *Behav Med* 2014;40(1):14-21.
[PUBMED](#) | [CROSSREF](#)
47. Driver HS, Werth E, Dijk DJ, Borbély AA. The menstrual cycle effects on sleep. *Sleep Med Clin* 2008;3(1):141.
[CROSSREF](#)
48. LeRoux A, Wright L, Perrot T, Rusak B. Impact of menstrual cycle phase on endocrine effects of partial sleep restriction in healthy women. *Psychoneuroendocrinology* 2014;49:34-46.
[PUBMED](#) | [CROSSREF](#)
49. Sveum K. Delayed sleep phase syndrome linked to irregular menstrual cycles, premenstrual symptoms in women. [https://aasm.org/delayed-sleep-phase-syndrome-linked-to-irregular-menstrual-cycles-premenstrual-symptoms-in-women/#\\$%](https://aasm.org/delayed-sleep-phase-syndrome-linked-to-irregular-menstrual-cycles-premenstrual-symptoms-in-women/#$%). Updated 2012. Accessed June 10, 2008.
50. Roenneberg T, Wirz-Justice A, Mrosovsky M. Life between clocks: daily temporal patterns of human chronotypes. *J Biol Rhythms* 2003;18(1):80-90.
[PUBMED](#) | [CROSSREF](#)
51. Kivelä A, Kauppila A, Ylöstalo P, Vakkuri O, Leppäluoto J. Seasonal, menstrual and circadian secretions of melatonin, gonadotropins and prolactin in women. *Acta Physiol Scand* 1988;132(3):321-7.
[PUBMED](#) | [CROSSREF](#)
52. Emens JS, Lewy AJ, Lefler BJ, Sack RL. Relative coordination to unknown "weak zeitgebers" in free-running blind individuals. *J Biol Rhythms* 2005;20(2):159-67.
[PUBMED](#) | [CROSSREF](#)
53. Kang W, Jang KH, Lim HM, Ahn JS, Park WJ. The menstrual cycle associated with insomnia in newly employed nurses performing shift work: a 12-month follow-up study. *Int Arch Occup Environ Health* 2019;92(2):227-35.
[PUBMED](#) | [CROSSREF](#)
54. Baumgartner A, Dietzel M, Saletu B, Wolf R, Campos-Barros A, Gräf KJ, et al. Influence of partial sleep deprivation on the secretion of thyrotropin, thyroid hormones, growth hormone, prolactin, luteinizing hormone, follicle stimulating hormone, and estradiol in healthy young women. *Psychiatry Res* 1993;48(2):153-78.
[PUBMED](#) | [CROSSREF](#)
55. Merklinger-Gruchala A, Ellison PT, Lipson SF, Thune I, Jasienska G. Low estradiol levels in women of reproductive age having low sleep variation. *Eur J Cancer Prev* 2008;17(5):467-72.
[PUBMED](#) | [CROSSREF](#)

56. Hall JE, Sullivan JP, Richardson GS. Brief wake episodes modulate sleep-inhibited luteinizing hormone secretion in the early follicular phase. *J Clin Endocrinol Metab* 2005;90(4):2050-5.
[PUBMED](#) | [CROSSREF](#)
57. Sequeira ME, Lewis SJ, Bonilla C, Smith GD, Joinson C. Association of timing of menarche with depressive symptoms and depression in adolescence: Mendelian randomisation study. *Br J Psychiatry* 2017;210(1):39-46.
[PUBMED](#) | [CROSSREF](#)
58. Li SX, Chan NY, Man Yu MW, Lam SP, Zhang J, Yan Chan JW, et al. Eveningness chronotype, insomnia symptoms, and emotional and behavioural problems in adolescents. *Sleep Med* 2018;47:93-9.
[PUBMED](#) | [CROSSREF](#)
59. Chan NY, Zhang J, Tsang CC, Li AM, Chan JW, Wing YK, et al. The associations of insomnia symptoms and chronotype with daytime sleepiness, mood symptoms and suicide risk in adolescents. *Sleep Med* 2020;74:124-31.
[PUBMED](#) | [CROSSREF](#)