



Correction to: Misclassification of Alcohol Use Disorder in MASLD and MetALD: Prevalence, Clinical Characteristics, and Outcomes

Jun-Hyuk Lee¹, Sung-Ho Ahn², Jimin Park³, So Young Jeon⁴, Eileen L. Yoon^{5,6}, Hye Sun Lee⁴, Dae Won Jun^{5,6}

¹Department of Family Medicine, Nowon Eulji Medical Center, Eulji University School of Medicine, Seoul, Korea; ²Department of Family Medicine, Gangnam Severance Hospital, Yonsei University College of Medicine, Seoul, Korea; ³Department of Medicine, Eulji University School of Medicine, Daejeon, Korea; ⁴Biostatistics Collaboration Unit, Department of Research Affairs, Yonsei University College of Medicine, Seoul, Korea; ⁵Hanyang Institute of Bioscience and Biotechnology, Hanyang University, Seoul, Korea; ⁶Department of Internal Medicine, Hanyang University College of Medicine, Seoul, Korea

<https://doi.org/10.5009/gnl250072>
Gut Liver 2025;19(5):735-745.

In the article entitled “Misclassification of Alcohol Use Disorder in MASLD and MetALD: Prevalence, Clinical Characteristics, and Outcomes,”¹ errors were identified in the legends of Figure 2 and Figure 3. The figures have been revised for clarity and accuracy. This article has been corrected online.

REFERENCE

1. Lee J, Ahn S, Park J, et al. Misclassification of alcohol use disorder in MASLD and MetALD: prevalence, clinical characteristics, and outcomes. Gut and Liver 2025;19:735-745.

