

Open

Besifovir Dipivoxil Maleate 144-Week Treatment of Chronic Hepatitis B: An Open-Label Extensional Study of a Phase 3 Trial

Hyung Joon Yim, MD¹, Won Kim, MD², Sang Hoon Ahn, MD³, Jin Mo Yang, MD⁴, Jae Young Jang, MD⁵, Yong Oh Kweon, MD⁶, Yong Kyun Cho, MD⁷, Yoon Jun Kim, MD⁸, Gun Young Hong, MD⁹, Dong Joon Kim, MD¹⁰, Young Kul Jung, MD¹, Soon Ho Um, MD¹, Joo Hyun Sohn, MD¹¹, Jin Woo Lee, MD¹², Sung Jae Park, MD¹³, Byung Seok Lee, MD¹⁴, Ju Hyun Kim, MD¹⁵, Hong Soo Kim, MD¹⁶, Seung Kew Yoon, MD¹⁷, Moon Young Kim, MD¹⁸, Kwan Sik Lee, MD³, Young Suk Lim, MD¹⁹, Wan Sik Lee, MD²⁰ and Kwang-Hyub Han, MD³

INTRODUCTION: Chronic hepatitis B (CHB) remains a major worldwide public health concern. Besifovir dipivoxil maleate (BSV) is a new promising treatment for CHB. However, long-term efficacy and safety have not yet been evaluated. Therefore, the goal of the study is to determine the antiviral efficacy and safety of BSV treatment over a 144-week duration (BSV-BSV) in comparison with those of a sequential treatment with tenofovir disoproxil fumarate (TDF) followed by a 96-week duration BSV administration (TDF-BSV).

METHODS: After 48 weeks of a double-blind comparison between BSV and TDF treatments, patients continued the open-label BSV study. We evaluated antiviral efficacy and drug safety up to 144 weeks for BSV-BSV and TDF-BSV groups. The primary endpoint was a virological response (hepatitis B virus DNA < 69 IU/mL).

RESULTS: Among the 197 patients enrolled, 170 and 158 patients entered the second-year and third-year open-label phase extensional study, respectively, whereas 153 patients completed the 144-week follow-up. The virological response rate over the 144-week period was 87.7% and 92.1% in BSV-BSV and TDF-BSV groups, respectively ($P = 0.36$). The rates of ALT normalization and HBeAg seroconversion were similar between the groups. No drug-resistant mutations to BSV were noted. Bone mineral density and renal function were well preserved in the BSV-BSV group and were significantly improved after switching therapy in TDF-BSV patients.

DISCUSSION: This extensional study of a phase 3 trial (NCT01937806) suggests that BSV treatment is efficacious and safe for long-term use in treatment-naïve and TDF-experienced patients with CHB.

SUPPLEMENTARY MATERIAL accompanies this paper at <http://links.lww.com/AJG/B485>, <http://links.lww.com/AJG/B491>

Am J Gastroenterol 2020;115:1217–1225. <https://doi.org/10.14309/ajg.0000000000000605>

INTRODUCTION

Chronic hepatitis B (CHB) remains a major worldwide public health problem despite the presence of effective vaccines and potent antivirals (1,2). Although the prevalence of chronic

hepatitis B virus (HBV) infection was significantly reduced after the introduction of a universal HBV vaccination program in many Asian countries, new cases of HBV infection are still reported, even in low-prevalence areas (3–5). In South Korea,

¹Department of Internal Medicine, Korea University College of Medicine, Seoul, Korea; ²Department of Internal Medicine, Seoul National University College of Medicine, Seoul Metropolitan Government Boramae Medical Center, Seoul, Korea; ³Department of Internal Medicine, Yonsei University College of Medicine, Seoul, Korea; ⁴Department of Internal Medicine, Catholic University Medical College St. Vincent, Suwon, Korea; ⁵Department of Internal Medicine, College of Medicine, Soonchunhyang University, Seoul, Korea; ⁶Department of Internal Medicine, Kyungpook National University College of Medicine, Daegu, Korea; ⁷Department of Internal Medicine, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul, Korea; ⁸Department of Internal Medicine, Liver Research Institute, Seoul National University College of Medicine, Seoul, Korea; ⁹Department of Internal Medicine, Kwangju Christian Hospital, Gwangju, Korea; ¹⁰Department of Internal Medicine, Hallym University College of Medicine, Chuncheon, Korea; ¹¹Department of Internal Medicine, Hanyang University College of Medicine, Seoul, Korea; ¹²Department of Internal Medicine, Inha University College of Medicine, Incheon, Korea; ¹³Department of Internal Medicine, Paik Hospital, Inje University, Busan, Korea; ¹⁴Department of Gastroenterology and Hepatology, Chungnam National University School of Medicine, Daejeon, Korea; ¹⁵Department of Internal Medicine, Gacheon University College of Medicine, Incheon, Korea; ¹⁶Department of Internal Medicine, Soonchunhyang University College of Medicine, Cheonan, Korea; ¹⁷Department of Internal Medicine, Catholic University of Korea, Seoul, Korea; ¹⁸Department of Internal Medicine, Yonsei University Wonju College of Medicine, Wonju, Korea; ¹⁹Department of Gastroenterology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea; ²⁰Department of Internal Medicine, Chonnam University Medical School, Gwangju, Korea. **Correspondence:** Kwang-Hyub Han, MD. E-mail: gihankhys@yuhs.ac

Received October 25, 2019; accepted February 26, 2020; published online May 1, 2020

Table 1. Baseline characteristics of the study subjects

	BSV-BSV (n = 81)	TDF-BSV (n = 76)	P Value
Male	54 (66.7)	49 (64.5)	0.77
Age, yr	46.00 (10.9)	43.99 (9.7)	0.22
HBeAg positive	51 (63.0)	43 (56.6)	0.87
HBV genotype			0.41
A	1 (1.2)	0 (0.0)	
C	79 (97.5)	75 (98.7)	
D	0 (0.0)	1 (1.3)	
Not determined	1 (1.2)	0 (0.0)	
HBV DNA (log ₁₀ IU/mL)	6.3 (1.7)	6.5 (1.5)	0.62
ALT (U/L)	106.15 (101.6)	128.41 (141.8)	0.36
ALT > ULN by central lab criteria ^a	78 (96.3)	71 (93.4)	0.48
ALT normalization by central lab criteria	3 (3.7)	5 (6.6)	
eGFR by MDRD (mL/min)	89.74 (14.8)	91.97 (13.7)	0.36
Creatinine (mg/dL)	0.86 (0.2)	0.84 (0.1)	0.43
Phosphate (mg/dL)	3.54 (0.5)	3.44 (0.6)	0.29
Total hip bone mineral density clinical status, n ^b	70	58	0.80
Normal (T-score ≥ -1.0)	58/70 (82.9)	49/58 (84.5)	
Osteopenia (-2.5 ≤ T-score < -1.0)	12/70 (17.1)	9/58 (15.5)	
Osteoporosis (T-score < -2.5)	0 (0.0)	0 (0.0)	
Data not collected	11	18	
Spine bone mineral density clinical status, n ^c	73	62	0.67
Normal (T-score ≥ -1.0)	50/73 (68.5)	40/62 (64.5)	
Osteopenia (-2.5 ≤ T-score < -1.0)	19/73 (26.0)	20/62 (32.3)	
Osteoporosis (T-score < -2.5)	4/73 (5.5)	2/62 (3.2)	
Data not collected	8	14	
Concurrent history	64 (79.0)	50 (65.8)	0.06
Mild renal impairment ^c	48 (59.3)	40 (52.6)	0.40
Cirrhosis	19 (23.5)	14 (18.4)	0.44
Cardiovascular disease	0 (0.0)	0 (0.0)	—
Diabetes mellitus	12 (12.8)	4 (4.3)	0.05
Hypertension	11 (11.7)	10 (10.8)	0.94
Hyperlipidemia	2 (2.1)	2 (2.2)	1.00
Prior antiviral therapy	0 (0.0)	1 (1.1) ^a	0.48

Data are number (%), n/N (%), or mean (SD) unless otherwise stated. P value: 2-sample *t* test or Wilcoxon rank-sum test for continuous variables and χ^2 test or Fisher exact test for categorical variables.

ALT, alanine aminotransferase; BMD, bone mineral density; BMD, bone mineral density; BSV, besifovir dipivoxil maleate; eGFR, estimated glomerular filtration rate; HBeAg, hepatitis B e antigen; HBV, hepatitis B virus; MDRD, modification of diet in renal disease; TDF, tenofovir disoproxil fumarate; ULN, upper limit of normal.

^aThirty-three U/L for women, 41 U/L for men.

^bFor BMD, some data were not collected.

^c50 ≤ eGFR < 90 (mL/min).

CHB still has a significant impact on public health, considering that the hepatitis B surface antigen (HBsAg) is detected in 60%–70% of patients with liver cirrhosis (5) and 65%–75% of patients with hepatocellular carcinoma (HCC) (6,7).

Currently, second-generation nucleos(t)ide analogs (NAs), such as entecavir, tenofovir disoproxil fumarate (TDF), and

tenofovir alafenamide fumarate (TAF), which have high genetic barriers to resistance and potent viral suppression activity, are used as primary treatments of CHB (8).

Besifovir dipivoxil maleate (BSV), an acyclic nucleotide phosphonate (a guanosine monophosphate), has been tested in several phase Ia/Ib studies and has shown potent suppression of




Figure 1. Patient disposition. BSV, besifovir dipivoxil maleate; CPK, creatinine phosphokinase; FAS, full analysis set; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; IP, investigational product; PPS, per-protocol set; TDF, tenofovir disoproxil fumarate.

HBV replication (9–12). BSV was approved for the treatment of chronic HBV infection in May 2017 in South Korea after we performed a randomized, multicenter, double-blind phase 3 study comparing the efficacy and safety of once-daily treatment with BSV vs TDF for 48 weeks in naïve patients with CHB (13). We reported that BSV elicited minor side effects on the kidneys and bones, with antiviral efficacy comparable with that of TDF over 48 weeks of treatment (13). This study was planned as an open-label follow-up study in which patients receive BSV for 7 years after initial treatment with BSV or TDF.

The aim of this study was to compare the antiviral efficacy and safety of a BSV solo treatment (BSV-BSV) over a 144-week period with a 48-week TDF treatment, followed by a 96-week BSV sequential treatment (TDF-BSV).

METHODS

Study design

This experimental study took over a 144-week period of treating patients with hepatitis B in South Korea. The first 48 weeks corresponded to a double-blind, randomized, multicenter,

noninferiority, controlled trial comparing BSV (Ildong Pharmaceutical, Seoul, Korea) with TDF (Gilead Sciences, Foster City, CA) treatments. Patients from that study who were suited to continue a single-arm, open-label trial received BSV for an additional 96 weeks. Written informed consent for both the 48-week initial phase and the open-label phase was obtained from all study participants. This study was conducted with the approval of the Human Research Ethics Committee in the 22 participating hospitals in the Republic of Korea, following the Declaration of Helsinki and all other applicable guidelines, laws, and regulations. Detailed descriptions of the study design, sample size estimates, eligibility criteria, and overall study population have been previously reported (13).

Bone mineral density, noninvasive fibrosis test, and resistance surveillance were regularly performed (see Supplementary Material, Supplementary Digital Content 1, <http://links.lww.com/AJG/B491>). Particularly, FIB-4 and aspartate aminotransferase-to-platelet ratio indexes (APRI) were annually estimated to assess liver fibrosis (see Table, Supplementary Digital Content 2 for formula used, <http://links.lww.com/AJG/B485>).

Table 2. Virological, serological, and biochemical responses

	48 wk			144 wk		
	BSV (n = 97)	TDF (n = 95)	P Value	BSV-BSV (n = 81)	TDF-BSV (n = 76)	P Value
HBV DNA <69 IU/mL	76 (78.4)	80 (84.2)	0.30	71 (87.7)	70 (92.1)	0.36
HBV DNA <20 IU/mL	60 (61.9)	65 (68.4)	0.34	65 (80.3)	65 (85.5)	0.38
HBeAg loss, n/N (%) ^a	5/53 (9.4)	3/50 (6.0)	0.72	10/50 (20.0)	13/41 (31.7)	0.20
HBeAg seroconversion, n/N (%) ^a	3/53 (5.7)	1/50 (2.0)	0.62	4/50 (8.0)	5/41 (12.2)	0.73
HBsAg loss, n/N (%) ^b	0/87 (0.0)	1/87 (1.1)	1.00	0/79 (0.0)	1/71 (1.4)	0.47
HBsAg seroconversion, n/N (%) ^b	0/87 (0.0)	0/87 (0)	—	0/79 (0.0)	1/71 (1.4)	0.47
ALT normalization, n (%) ^c	69 (71.1)	71 (74.7)	0.57	66 (81.5)	67 (88.2)	0.25

Data are expressed as n (%), unless stated otherwise. P value: Pearson χ^2 test/Fisher exact test.

ALT, alanine aminotransferase; BSV, besifovir dipivoxil maleate; HBeAg, hepatitis B e antigen; HBsAg, hepatitis B surface antigen; TDF, tenofovir disoproxil fumarate.

^aAmong HBeAg-seropositive patients at baseline.

^bAmong HBsAg-seropositive patients at baseline.

^cAmong patients with baseline ALT levels above the central lab normal range (0–41 U/L for males and 0–33 U/L for females).

Subjects

Patients with CHB younger than 20 years of age and HBsAg-positive for at least 6 months before screening were eligible. Subjects with HBV DNA levels >17,241 IU/mL (1×10^5 copies/mL) and > 1,724 IU/mL (1×10^4 copies/mL), who tested positive and negative for hepatitis B e-antigen (HBeAg), respectively, were randomized 1:1 into BSV or TDF treatment groups for 48 weeks. To maintain blinding of investigators and subjects, subjects in the BSV group were given BSV 150 mg with L-carnitine 660 mg as a supplement and TDF placebo, whereas those in the TDF group were given TDF 300 mg with L-carnitine as a BSV placebo. After 48 weeks, the patients continued on BSV or were switched to BSV 150 mg with L-carnitine 660 mg during the open-label phase for up to 144 weeks.

Endpoints

The efficacy endpoint was the proportion of subjects with a virological response, defined as HBV DNA <69 IU/mL at week 144. Other efficacy and safety endpoints, including the proportions of subjects with undetectable HBV DNA levels (<20 IU/mL),

changes of serum HBV DNA levels, HBsAg or HBeAg seroconversion, alanine aminotransferase (ALT) normalization, drug resistance, adverse events (AEs), bone mineral density (BMD), renal function-related parameters, fibrosis parameters, incidences of HCC, and other abnormalities, were assessed.

Laboratory tests and assessments

All laboratory data during the trial were collected at 12-week intervals via the Central Lab (GC abs, Yongin, Korea). HBV DNA quantification was performed using a COBAS AmpliPrep/TaqMan test (Roche Diagnostics, Indianapolis, IN) with a lower detection limit of 20 IU/mL. Other laboratory methods were followed as previously reported (13).

Bone mineral density, non-invasive fibrosis test, and resistance surveillance were regularly performed (see Supplementary Material, Supplementary Digital Content 1, <http://links.lww.com/AJG/B491>). Particularly, FIB-4 and aspartate aminotransferase-to-platelet ratio indexes (APRI) were annually estimated to assess liver




Figure 2. Viral suppression by study visit. (a) Proportions of patients with HBV DNA <69 IU/mL as determined by the FAS. (b) Proportions of patients with HBV DNA <20 IU/mL as determined by FAS. Bars represent 95% confidence intervals. BSV, besifovir dipivoxil maleate; FAS, full analysis set; HBV, hepatitis B virus; TDF, tenofovir disoproxil fumarate.




Figure 3. Antiviral responses by study visit (weeks). (a) Degree of HBV DNA reduction (log IU/mL). (b) Mean HBV DNA level (log IU/mL). ANOVA, analysis of variance; BSV, besifovir dipivoxil maleate; HBV, hepatitis B virus; TDF, tenofovir disoproxil fumarate.

fibrosis (see Table, Supplementary Digital Content 2 for formula used, <http://links.lww.com/AJG/B485>).

Statistical analysis

For primary efficacy analysis, proportions of patients with HBV DNA <69 IU/mL were determined using the full analysis set (FAS), which included all patients who received at least one investigational product and completed at least one efficacy evaluation. Per-protocol analysis was performed after excluding consent withdrawals and major protocol violators from the FAS population.

Differences in the baseline characteristics and secondary endpoints between the groups were tested using an independent 2-sample *t* test or Wilcoxon rank-sum test for continuous variables and a χ^2 test or Fisher exact test for categorical variables. We followed the CONSORT guidelines for randomized controlled trials.

RESULTS

Study population

Of 197 patients who were initially randomly assigned, 170 (86%) entered the second year, 158 (80%) entered the third year extensional study, and finally 153 completed all 144 weeks of follow-up (Figure 1). Of 157 patients (81 in BSV-BSV and 76 in TDF-BSV) who qualified for the FAS at 144 weeks, 145 were included for the PPS analysis. The baseline characteristics for both treatment groups were well-balanced (Table 1). Considering both the groups, 65.6% patients enrolled were men, with a mean age of 45 years. Subjects positive for HBeAg represented 63.0% in the BSV-BSV group and 56.6% in the TDF-BSV group, with the most common HBV genotype being C. For BSV-BSV and TDF-BSV groups, the means of HBV DNA at the baseline were 6.3 and 6.5 log₁₀ IU/mL, respectively. More than half of the patients showed a normal T-score in the hip and/or spine at the baseline (64.5%–84.5%). Although the BSV-BSV group was more likely to have concurrent histories, including cirrhosis and diabetes mellitus, at the baseline, no statistically significant differences were noted between the groups.

Efficacy

Virological response. The rate of virological response (HBV DNA < 69 IU/mL) in BSV-BSV was 87.7% (71/81), whereas 92.1% (70/76) of patients who switched the treatment from TDF to BSV responded up to 144 weeks, indicating that the BSV-BSV group met the efficacy endpoint of noninferiority with pre-determined –15% margin to BSV-TDF according to the FAS (lower bound 95% CI = –13.8%, *P* = 0.36) (Table 2 and Figure 2a). The complete virological response with the lower limit of HBV DNA quantification (<20 IU/mL) was 80.3% (65/81) among the BSV-BSV patients and 85.5% (65/76) in the TDF-BSV group at week 144 (*P* = 0.38) (Table 2 and Figure 2b). Similar results were observed in the PPS (see Figure, Supplementary Digital Content 3A, 3B, <http://links.lww.com/AJG/B485>) (HBV DNA < 69 IU/mL, 92.1% vs 95.7%, *P* = 0.50; HBV DNA < 20 IU/mL, 84.2% vs 88.4%, *P* = 0.46; in BSV-BSV vs TDF-BSV, respectively).

Antiviral responses from the baseline during every follow-up visit until week 144 are shown in Figure 3. During the first 48 weeks, the degree of HBV DNA reduction was not significantly different between the 2 groups (Figure 3a), except for at the end of the first month. Mean HBV DNA decreased to <2 log₁₀ IU/mL

Table 3. Improvements in noninvasive fibrosis markers FIB-4 and APRI indexes at 144 weeks

	BSV-BSV (n = 81)	TDF-BSV (n = 76)	P value ^a	P value ^b
FIB-4				
Baseline	2.12 (2.07)	1.92 (1.14)	—	—
144 wk	1.32 (0.76)	1.27 (0.83)	—	—
Change	–0.80 (1.77)	–0.65 (0.92)	0.79	0.54
P value ^c	<0.0001	<0.0001		
APRI				
Baseline	1.22 (1.31)	1.35 (1.32)	—	—
144 wk	0.40 (0.30)	0.39 (0.38)	—	—
Change	–0.82 (1.33)	–0.97 (1.31)	0.30	0.29
P value ^c	<0.0001	<0.0001		

Data are expressed as mean (standard deviation), unless otherwise stated. APRI, aspartate aminotransferase-to-platelet ratio index; BSV, besifovir dipivoxil maleate; TDF, tenofovir disoproxil fumarate.

^aBetween-group: Wilcoxon rank-sum test.

^bBetween-group: mixed model with repeated measures (every 12 weeks measurement).

^cWithin-group: Wilcoxon signed-rank test.

Table 4. Safety data

Variables	0–144 wk		96–144 wk	
	BSV (n = 81)	TDF (n = 76)	BSV-BSV (n = 81)	TDF-BSV (n = 76)
Adverse events, n (%), [cases]	63 (77.8), [253]	56 (73.7), [209]	54 (66.7), [125]	42 (55.3), [95]
Adverse drug reactions	31 (38.3)	26 (34.2)	23 (28.4)	15 (19.7)
Serious adverse events	7 (8.6)	7 (9.2)	1 (1.2)	2 (2.6)
Serious adverse drug reactions	1 (1.2)	0	0	0
Death	0	0	0	0
Adverse events recorded in ≥3% of all patients				
Nasopharyngitis	14 (17.3)	14 (18.4)	3 (3.7)	4 (5.3)
Dyspepsia	8 (9.9)	10 (13.2)	1 (1.2)	1 (1.3)
Headache	6 (7.4)	3 (4.0)	1 (1.2)	1 (1.3)
Back pain	6 (7.4)	4 (5.3)	3 (3.7)	3 (4.0)
Fatigue	5 (6.2)	5 (6.6)	1 (1.2)	2 (2.6)
Pruritus	3 (3.7)	4 (5.3)	0	1 (1.3)
Gastritis	2 (2.5)	9 (11.8)	1 (1.2)	8 (10.5)
Nausea	5 (6.2)	1 (1.3)	0	0
ALT elevation	6 (7.4)	2 (2.6)	2 (2.5)	0
Diarrhea	3 (3.7)	5 (6.6)	2 (2.5)	0
Urticaria	1 (1.2)	5 (6.6)	0	0
Dizziness	8 (9.9)	0	3 (3.7)	0
Renal-related adverse events				
Proteinuria	0	1 (1.3)	0	1 (1.3)
Phosphate < 2.5 mg/dL	9 (11.1)	16 (21.1)	6 (7.4)	5 (6.6)
eGFR < 50 mL/min	1 (1.2)	2 (2.6)	0	1 (1.3)
Bone related adverse events				
Fracture ^a	2 (2.5)	1 (1.3)	1 (1.2)	0
Total-carnitine level at week 144				
Decrease	1 (1.2)	1 (1.3)	—	—
Increase	31 (38.3)	28 (36.8)	—	—

Data are expressed as n (%), unless stated otherwise.
 ALT, Alanine aminotransferase; BSV, besifovir dipivoxil maleate; eGFR, estimated glomerular filtration rate; TDF, tenofovir disoproxil fumarate.
^aAdverse event due to trauma or accident.

levels at week 24 and remained stable for 144 weeks (Figure 3b). No significant differences between the 2 groups were observed during the 144 weeks (Figure 3b).

Biochemical and serological responses. Among HBeAg-positive patients, the cumulative incidences of HBeAg loss at 144 weeks were 20.0% (10/50) in the BSV-BSV group and 31.7% (13/41) in the TDF-BSV group ($P = 0.20$). The incidences of HBeAg seroconversion were 8.0% (4/50) and 12.2% (5/41) in the BSV-BSV and TDF-BSV groups, respectively ($P = 0.73$). There was no significant difference in HBeAg serological responses between both groups (Table 2); only one patient (1.4%, 1/71) in the TDF-BSV group showed HBsAg loss during the 144 weeks (Table 2). ALT levels normalized in 81.5% (66/81) and 88.2% (67/76) subjects in the BSV-BSV and TDF-BSV groups after 144 weeks, with no significant difference between the groups ($P = 0.25$) (Table 2).

Subgroup analysis. Biochemical, serological, and virological responses were separately evaluated in HBeAg-positive and HBeAg-negative patients according to the baseline HBeAg status for FAS and PPS (see Tables, Supplementary Digital Content 4 and 5, <http://links.lww.com/AJG/B485>; see Figure, Supplementary Digital Content 6, <http://links.lww.com/AJG/B485>).

In the FAS, all HBeAg-negative patients reached HBV DNA <69 IU/mL at week 144. Only one patient in each group did not achieve a complete virological response (HBV DNA < 20 IU/mL) by week 144. There was no difference in virological response rates among HBeAg-negative patients.

Among HBeAg-positive patients, the proportions who achieved HBV DNA <69 IU/mL were 80.4% (41/51) in the BSV-BSV and 86.1% (37/43) in the TDF-BSV group ($P = 0.47$). The rates of HBV DNA <20 IU/mL were 70.6% (36/51) and 76.7% (33/43), respectively ($P = 0.50$). There was no difference in virological




Figure 4. Median changes from baseline in eGFR (MDRD) by study week. Data are presented as median (Q1, Q3) values (mL/min). BSV, besifovir dipivoxil maleate; eGFR, estimated glomerular filtration rate; MDRD, modification of diet in renal disease; TDF, tenofovir disoproxil fumarate.

response rates between both the groups when considering HBsAg-positive patients.

Virological breakthrough and resistance surveillance. Both groups showed no antiviral resistance for 144 weeks. The proportions of patients who experienced virological breakthrough between 96 and 144 weeks were 4.9% (4/81) in the BSV-BSV group and 7.9% (6/76) in the TDF-BSV group, with no significant difference between the groups. However, serum HBV DNA became undetectable in all these patients without modifying the therapy during follow-up. DNA sequencing was performed for patients who experienced a virological breakthrough, confirming that new mutations that caused by amino acid substitutions were not detected in the HBV conserved sites.

Noninvasive fibrosis markers: FIB-4 and APRI. The FIB-4 and APRI indexes significantly improved at 144 weeks vs baseline in both groups (Table 3; see Table, Supplementary Digital Content 7, <http://links.lww.com/AJG/B485>). FIB-4 improved from 2.12 to 1.32 and APRI from 2.12 to 0.4 in the BSV-BSV group. Likewise, FIB-4 improved from 1.92 to 1.27 and APRI from 1.35 to 0.39 in the TDF-BSV group, respectively (all $P < 0.001$). According to

these results, the degree of hepatic fibrosis significantly decreased in both the BSV-BSV and TDF-BSV groups over the duration of drug administration. There was no significant difference in FIB-4 and APRI values between both the groups.

Safety

Adverse events. After reporting AEs during the 96-week study (13), 3 serious AEs occurred: HCC in one patient from the BSV-BSV group and HCC and type 2 diabetes mellitus in 2 patients of the TDF-BSV group (Table 4). However, the investigator confirmed that all 3 patients could continue the study. No serious adverse drug reactions or death was newly reported during 96–144 weeks in the study. L-carnitine levels were measured during every follow-up visit. These levels were stably maintained, with only one patient in each group exhibiting a decreased level.

The median changes in eGFR from the baseline were -0.5 , -0.7 , and -0.5 mL/min at 48, 96, and 144 weeks, respectively, for the BSV-BSV group, indicating no effect of BSV on renal function. On the contrary, for the TDF-BSV group, the median changes in eGFR levels from baseline were -7.8 , -0.4 , and -0.8 mL/min at 48, 96, and 144 weeks, respectively; therefore, the recovered renal function after switching to BSV at 49 weeks was maintained up to 144 weeks (Figure 4). There was no incidence of acute kidney injury in the 2 groups.

The mean percent change in spine BMD from the baseline to week 144 was -0.02% (95% CI: -2.25 to 2.22) for the BSV-BSV ($P = 0.59$) and -0.31% (95% CI: -2.37 to 1.74) for the TDF-BSV ($P = 0.93$) group. This difference was not significant (Figure 5a). The mean percent change in hip BMD from the baseline to week 144 was -0.42% (95% CI: -1.53 to 0.7) for the BSV-BSV ($P = 0.10$) and 0.78% (95% CI: -1.21 to 2.77) for the TDF-BSV group. This difference was not significant (Figure 5b).

DISCUSSION

The incidence of HBV infection is decreasing worldwide because of improved socioeconomic status, universalization of vaccine programs, and effective antiviral treatments (1,4). However, population movements are now affecting the occurrence and prevalence of HBV in several low-endemic countries




Figure 5. Changes in BMD. (a) Mean percentage changes in the spine at weeks 48, 96, and 144 of treatment. Bars represent 95% confidence interval. (b) Mean percentage changes in the hip at weeks 48, 96, and 144 of treatment. Bars represent 95% confidence intervals. BMD, bone mineral density; BSV, besifovir dipivoxil maleate; TDF, tenofovir disoproxil fumarate.

(14–17). Unlike most infectious diseases, deaths from HBV-related liver cirrhosis and/or HCC increased between 1990 and 2013 (18).

BSV, the newest approved agent in the NA class in Korea, is currently recommended as the first-line oral agent for CHB treatment by the Korean Association for the Study of the Liver (19). Our study followed the CHB treatment with BSV for 144 weeks, with findings consistent with those of our previous 96-week study (13). In total, 89.8% and 82.8% of patients had HBV DNA <69 and <20 IU/mL, respectively, indicating that the antiviral effect of BSV was maintained up to 144 weeks. Reduction in hip and spine BMD was observed in patients treated with TDF for 48 weeks, but it returned to the baseline level after switching to BSV, and it was maintained up to 144 weeks. BSV treatment also showed a favorable safety profile regarding renal parameters. No viral resistance was detected during the course of the 144-week treatment. We also confirmed the improvement of fibrosis applying the FIB-4 and APRI tests.

A recent 3-year study showed that TAF was associated with a lower risk of BMD decline and nephrotoxicity (20). The mean BMDs decreased from the baseline to week 144 of therapy (hip: -2.49 vs -0.41 , $P < 0.001$; spine: -2.01 vs -0.51 , $P < 0.001$; in TDF and TAF groups, respectively). We also observed a significant decrease in BMD in the TDF group relative to that in the BSV group at 48 weeks. However, in the group of patients receiving BSV for 144 weeks, the BMD mean changes from the baseline were -0.42 (hip) and 0.02 (spine), similar to those results obtained with TAF. Treatment with BSV and TAF also showed similar trends in eGFR changes over 3 years (eGFR_{MDRD} for BSV, -0.5 vs eGFR_{CG} for TAF, -1.2). Although it is difficult to perform a direct comparison between BSV and TAF treatments, they seem to be similar for preserving renal function and bone density.

Improvements in fibrosis after antiviral treatment have been previously reported by histology (21,22). Although liver biopsy is the accepted standard method for the assessment of hepatic fibrosis, repeated evaluations are difficult to perform because of cost, the risk of life-threatening complications, and a lack of available expertise (23–25). Therefore, we analyzed the extent of liver fibrosis using FIB-4 and APRI as biomarkers, as indicated by previous studies (26–28). Importantly, in 2 noninvasive fibrosis tests, our results showed a clear fibrosis improvement on BSV administration.

Carnitine depletion was the most common side effect of BSV (11,12). Therefore, BSV was supplemented with 660 mg of L-carnitine. Carnitine concentrations were closely monitored during the study period and at week 144. Except for one patient in each group who evidenced an increase in carnitine levels, patients were generally stably maintained with the supplementation and none experienced clinical features of carnitine deficiency, such as hypoglycemia, hypoketosis, and encephalopathy.

This study has several limitations. Although the general safety concerns related to nephrotoxicity can be ignored in the BSV group considering the rate of decline of eGFR, additional markers related to renal function, such as urinary albumin, fractional excretion of uric acid, and urinary beta-2 microglobulin, were not evaluated. Moreover, no assessments of bone turnover markers for the estimation of bone formation or resorption were performed, although it is recommended for

patients receiving these NAs. Although noninvasive fibrosis markers were evaluated to demonstrate the improvement of liver fibrosis, histological analysis did not confirm such improvement; this calls for further evaluation. However, to our knowledge, this is the first study reporting antiviral effects of BSV providing safe renal and bone profile for an extended treatment period.

In summary, BSV treatment maintained antiviral efficacy over 144-week period without any viral resistance. BSV administration was safe, well-tolerated, and effective for treatment-naïve patients with CHB as well as those who switched from TDF to BSV treatment.

ACKNOWLEDGMENTS

We would like to thank Editage (www.editage.com) for English language editing and publication support.

CONFLICTS OF INTEREST

Guarantor of the article: Kwang-Hyub Han, MD.

Specific author contributions: H.J.Y. wrote the manuscript and performed the formal analysis and critical revision. K.H.H.: was involved in the study conception, design, and supervision. All the other authors contributed to data acquisition and revision of the manuscript.

Financial support: This work was supported by Ildong Pharmaceutical. This sponsor had a partial role in study design, collection, analysis, and interpretation of data.

Potential competing interests: K.H.H. received grants from the clinical trial sponsor Ildong Pharmaceutical. All the other authors declare no conflicts of interest.

Study Highlights

WHAT IS KNOWN

- ✓ BSV is a new treatment of chronic hepatitis B virus infection.
- ✓ BSV has an antiviral efficacy comparable with that of TDF over 48 weeks.
- ✓ BSV has a better safety profile than TDF in terms of bone and renal outcomes.

WHAT IS NEW HERE

- ✓ BSV demonstrated durable and potent antiviral activity over 144 weeks.
- ✓ Approximately 88% of subjects had HBV DNA <69 IU/mL without antiviral resistance mutations.
- ✓ Long-term BSV treatment for chronic hepatitis B is safe and improves fibrosis.

REFERENCES

1. European Association for the Study of the Liver. EASL 2017 clinical practice guidelines on the management of hepatitis B virus infection. *J Hepatol* 2017;67:370–98.
2. Schweitzer A, Horn J, Mikolajczyk RT, et al. Estimations of worldwide prevalence of chronic hepatitis B virus infection: A systematic review of data published between 1965 and 2013. *Lancet* 2015;386:1546–55.
3. Nelson NP, Easterbrook PJ, McMahon BJ. Epidemiology of hepatitis B virus infection and impact of vaccination on disease. *Clin Liver Dis* 2016; 20:607–28.

4. Cho EJ, Kim SE, Suk KT, et al. Current status and strategies for hepatitis B control in Korea. *Clin Mol Hepatol* 2017;23:205–11.
5. Chae HB, Kim JH, Kim JK, et al. Current status of liver diseases in Korea: Hepatitis B. *Korean J Hepatol* 2009;15:S13–24.
6. Cheon JH, Park JW, Park KW, et al. The clinical report of 1,078 cases of hepatocellular carcinomas: National Cancer Center experience. *Korean J Hepatol* 2004;10:288–97.
7. Kim SR, Kudo M, Hino O, et al. Epidemiology of hepatocellular carcinoma in Japan and Korea. A review. *Oncology* 2008;75:13–6.
8. Buti M, Riveiro-Barciela M, Esteban R. Long-term safety and efficacy of nucleos(t)ide analogue therapy in hepatitis B. *Liver Int* 2018;38(Suppl 1): 84–9.
9. Yuen MF, Kim J, Kim CR, et al. A randomized placebo controlled, dose-finding study of oral LB80380 in HBeAg-positive patients with chronic hepatitis B. *Antivir Ther* 2006;11:977–83.
10. Yuen MF, Han KH, Um SH, et al. Antiviral activity and safety of LB80380 in hepatitis B e antigen-positive chronic hepatitis B patients with lamivudine-resistant disease. *Hepatology* 2010;51:767–76.
11. Lai CL, Ahn SH, Lee KS, et al. Phase IIb multicentred randomised trial of besifovir (LB80380) versus entecavir in Asian patients with chronic hepatitis B. *Gut* 2014;63:996–1004.
12. Yuen MF, Ahn SH, Lee KS, et al. Two-year treatment outcome of chronic hepatitis B infection treated with besifovir vs. entecavir: Results from a multicentre study. *J Hepatol* 2015;62:526–32.
13. Ahn SH, Kim W, Jung YK, et al. Efficacy and safety of besifovir dipivoxil maleate compared with tenofovir disoproxil fumarate in treatment of chronic hepatitis B virus infection. *Clin Gastroenterol Hepatol* 2019;17: 1850–9.
14. Coppola N, Alessio L, Gualdieri L, et al. Hepatitis B virus, hepatitis C virus and human immunodeficiency virus infection in undocumented migrants and refugees in southern Italy, January 2012 to June 2013. *Euro Surveill* 2015;20:30009.
15. Hampel A, Solbach P, Cornberg M, et al. Current seroprevalence, vaccination and predictive value of liver enzymes for hepatitis B among refugees in Germany. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2016;59:578–83.
16. Levy V, Yuan J, Ruiz J, et al. Hepatitis B sero-prevalence and risk behaviors among immigrant men in a population-based household survey in low-income neighborhoods of northern California. *J Immigr Minor Health* 2010;12:828–33.
17. Coppola N, Alessio L, Gualdieri L, et al. Hepatitis B virus infection in undocumented immigrants and refugees in southern Italy: Demographic, virological, and clinical features. *Infect Dis Poverty* 2017;6:33.
18. Stanaway JD, Flaxman AD, Naghavi M, et al. The global burden of viral hepatitis from 1990 to 2013: Findings from the global burden of disease study 2013. *Lancet* 2016;388:1081–8.
19. Korean Association for the Study of the Liver (KASL). KASL clinical practice guidelines for management of chronic hepatitis B. *Clin Mol Hepatol* 2019;25:93–159.
20. Chan HLY, Lim YS, Seto WK, et al. 3-year efficacy and safety of tenofovir alafenamide compared with tenofovir disoproxil fumarate in HBeAg-negative and -positive patients with chronic hepatitis B. *Hepatology* 2018; 68(Suppl 1):227–8A.
21. Lai CL, Shouval D, Lok AS, et al. Entecavir versus lamivudine for patients with HBeAg-negative chronic hepatitis B. *N Engl J Med* 2006;354: 1011–20.
22. Marcellin P, Gane E, Buti M, et al. Regression of cirrhosis during treatment with tenofovir disoproxil fumarate for chronic hepatitis B: A 5-year open-label follow-up study. *Lancet* 2013;381:468–75.
23. Spinzi G, Terruzzi V, Minoli G. Liver biopsy. *N Engl J Med* 2001;344:2030.
24. Maharaj B, Maharaj RJ, Leary WP, et al. Sampling variability and its influence on the diagnostic yield of percutaneous needle biopsy of the liver. *Lancet* 1986;1:523–5.
25. Seeff LB, Everson GT, Morgan TR, et al. Complication rate of percutaneous liver biopsies among persons with advanced chronic liver disease in the HALT-C trial. *Clin Gastroenterol Hepatol* 2010;8:877–83.
26. Kim WR, Loomba R, Berg T, et al. Impact of long-term tenofovir disoproxil fumarate on incidence of hepatocellular carcinoma in patients with chronic hepatitis B. *Cancer* 2015;121:3631–8.
27. Lin CL, Liu CH, Wang CC, et al. Serum biomarkers predictive of significant fibrosis and cirrhosis in chronic hepatitis B. *J Clin Gastroenterol* 2015;49:705–13.
28. Sonneveld MJ, Brouwer WP, Chan HL, et al. Optimisation of the use of APRI and FIB-4 to rule out cirrhosis in patients with chronic hepatitis B: Results from the SONIC-B study. *Lancet Gastroenterol Hepatol* 2019;4:538–44.
29. Kim BK, Kim DY, Park JY, et al. Validation of FIB-4 and comparison with other simple noninvasive indices for predicting liver fibrosis and cirrhosis in hepatitis B virus-infected patients. *Liver Int* 2010;30:546–53.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.